

2nd Global Summit on

ONCOLOGY & CANCER

March 12-14, 2018 Singapore

Role of post mastectomy radiotherapy in T1, T2 lesions with 1-3 positive axillary lymph nodes: Study of 101 cases

Nikhil Garg

The Gujarat Cancer & Research Institute, India

Statement of the Problem: Post mastectomy radiotherapy (PMRT) reduces loco-regional recurrence (LRR) and improves overall survival. There is international consensus to recommend PMRT for patients with tumor size more than 5 cm, tumor invasion of the skin, pectoral muscle or chest wall and patients with >4 positive lymph nodes. However, the role of PMRT for patients with T1, T2 disease with 1-3 positive LN is still controversial. The side effects of radiotherapy and its associated morbidity must be considered in the risk benefit ratio, thus difficult to arrive at consensus in early breast cancer.

Methods: 101 patients treated between 2012 to 2015 were studied retrospectively, the inclusion criteria for this analysis were: Female patients with unilateral breast cancer and no distant metastasis at initial diagnosis who underwent mastectomy and axillary lymph node dissection; postoperative pathology indicated T1-2 and 1-3 positive axillary lymph nodes (T1-2N1M0) disease, at least 10 lymph nodes removed by axillary dissection; complete surgical resection of the tumor and negative margins; complete estrogen receptor (ER), progesterone receptor (PR) and human epithelial growth factor receptor family 2 (Her2) status; no neo-adjuvant chemotherapy was administered before surgery and endocrine therapy was performed based on the hormone receptor status. To study the research questions, we formulated hypotheses as follows: Radiotherapy does not have any impact on recurrence post mastectomy and there is no influence of peri nodal extension on recurrence. The above hypotheses were tested using chi-square test.

Results: Recurrences were obtained in 9 amongst radiotherapy and without radiotherapy in 16. When chi-square was applied, the value was highly significant. Hence, our hypothesis was rejected. Also in case of PNE with recurrence and radiotherapy, 8 had PNE with radiotherapy and recurrence and 27 had no recurrence, p value was 0.013% and hence highly significant.

Conclusion: Radiotherapy should be strongly considered in patients with 1-3 nodes post mastectomy as it decreases the chances of recurrence

nikhilgarg9490@gmail.com