

5th International Conference on **Biomarkers & Clinical Research**

April 15-17, 2014 St. Hilda's College - University of Oxford, UK

From chemoprevention to identification of potential therapeutic targets and biomarkers for breast cancer

Somya Shanmuganathan¹, Allal Ouhtit¹, Ishita Gupta¹, Hamad Al-Riyami¹ and Madhwa Raj²
¹Sultan Qaboos University, Sultanate of Oman
²Louisiana State University Health Science Center, USA

Ostudies, individually and often at higher doses to kill cancer cells. Based on the combination and synergism theory, we had previously demonstrated that a combination of Resveratrol and Indole-3-Carbinol synergized and killed a maximum number of Breast Cancer cells. In the present study, we have tested various combinations of 10 well known phytochemicals, used at bioavailablelevels, for their effect on cell growth and proliferation of the MDA-MB-231; breast cancer (BC) cell line and MCF-10A; normal breast epithelium as control cell line. The results revealed a super combination of 7phyto-compounds(7SC), that synergized and induced 100% clearance of the BC cells but did notaffect the normal breast epithelial cells. Next, in order to understand the underlying molecular mechanisms of this 'synergism' effect, microarray analysis will be conducted on them RNA collected from the 7SC treated and control cells at 12 and 24 hour time points. Ongoing in vivo experiments aim to evaluate the efficacy of the 7SC phyto-compound treatment in preventing tumor growth using xenograft mouse BC model, and further validate the functional relevance of these genes in BC cell growth and survival. The data supports our hypothesis that the 7SC could be used in CAM as a dietary supplement approach against BC, and further identify genes that have the potential to serve as biomarkers and gene candidate to guide the design of appropriate anti-BC therapeutic strategies.

somya.2003@gmail.com