

13<sup>th</sup> International Conference on

# Tissue Engineering & Regenerative Medicine

July 12-13, 2018 Paris, France

## **The efficacy of platelet gel derived umbilical cord blood on diabetic foot ulcers: A double -blind randomized clinical trial**

**Seyedeh Esmat Hosseini<sup>1,2</sup>**<sup>1</sup>Royal Institute for Stem Cell Biology and Technology, Iran<sup>2</sup>Shahid Beheshti University of Medical Sciences, Iran

**Introduction:** Type 2 diabetes is one of the most prevalent diseases throughout the world. The foot ulcers are severe complications of this disease. Foot ulcers induce the high rate of morbidity, impair quality of life and bring about extreme costs to health service providers and give rise to waste of time. Recently, platelet-rich plasma (PRP) and platelet gel (PG) have been used for the treatment of chronic wounds.

**Materials & Methods:** In the present study, randomized, double-blind, placebo-controlled study, platelet gel derived from umbilical cord blood (UCB) was used to heal the diabetic foot. The patients were randomly divided into three groups, under the categories of platelet gel (PG), platelet-poor plasma (PPP) and lubricant gel (placebo) with (ratio 1:1:1). The processes of gels application were launched for the subject of each group twice per week with 3-4 days of interval. This mechanism protracted for eight weeks. After completion of eight weeks, the patients were followed up after two weeks and then continued for three months with one-month interval. Thirty patients underwent statistical analysis. Except for diastolic blood pressure which was significant between groups, there were no statistically significant differences in patients' baseline demographics. No significant differences were detected between groups at baseline of wounds ( $P=0.09$ ).

**Results:** The results revealed that there is no statistically significant interaction among three groups during follow- up time. The present study provides evidence that there are no significant differences in the size of wound among PG, PPP, and placebo group.