

2nd International Conference on

PHARMACEUTICAL CHEMISTRY

October 02-04, 2017 Barcelona, Spain

An optimized process for supercritical CO₂ extraction of high-value components from *Artemisia annua* L for cosmetic applications

Rosana M. Alberici Oliveira
University of Campinas, Brazil

Artemisia annua L. is a plant used as traditional Chinese herb medicines. The plant shows a strong antimalarial activity due to the presence in its aerial parts of artemisinin and its derivative compounds. A typical plant content of artemisinin is between 0.7– 1.5% w/w depending on the growth conditions and species. Besides being the only commercial source of artemisinin, the plant contains a pleasantly aromatic essential oil valued in perfumery and cosmetics. Recent published literatures reported that artemisinin and its derivatives present a wide range of biological activities, exhibiting therapeutic potential as anti-inflammatory and antimicrobial. These high-value components could be used to generate new sustainable multifunctional cosmetic products. However, the main process for obtaining artemisinin has been carried out by traditional organic solvent extraction that has residual solvent and low purity problems. Therefore, it is desirable to develop a greener extraction technique. Supercritical carbon dioxide extraction (SFE) can be very selective, presents high extraction yields and high purity advantages. The aims of this study were to evaluate the multi-use of leaf biomass for production of essential oil, artemisinin and residual extract using supercritical CO₂ fractional extraction. Thin layer chromatography (TLC) and liquid chromatography coupled with mass spectrometry (LC-MS/MS) were used for qualifying and quantifying artemisinin and its derivative compounds in the extracts while the essential oil was characterized by gas chromatography coupled with mass spectrometry (GC-MS). The influence of the extraction pressure and temperature on the extract composition was systematically measured, analyzing the recovered fractions at different extraction times during the extraction process. Under optimized experimental conditions it was possible to obtain the fractionation of the desired products.

Biography

Rosana M. Alberici is a Chemistry from the Federal University of São Carlos (UFSCar, Brazil). She received her Ph.D. from the University of Campinas (UNICAMP, Brazil). She has been a visiting research scientist at the School of Chemistry and Biochemistry, Georgia Institute of Technology, USA. She was also a postdoctoral fellow at the ThoMSon Mass Spectrometry Laboratory, Institute of Chemistry (UNICAMP, Brazil) where she is currently a Researcher Collaborator. Her research focuses on the area of analytical mass spectrometry applied to food, biofuels, natural products and cosmetics analysis.

rmaalberici@hotmail.com

Notes: