

International Conference on

PHYSICAL EDUCATION, SPORTS MEDICINE AND DOPING STUDIES

August 08-09, 2018 Osaka, Japan

Utilization of memory foam insole as shoe modification to decrease foot pain of a track and field athlete in Saint Dominic College of Asia**Charen Rabe**

St. Dominic College of Asia, Philippines

Introduction: Memory foam molds to the body in response to heat and pressure, distributing body weight evenly. It conforms to the body and distributes the weight in way that allows the entire body to be supported. The shoe insole is removable shoes insert that is worn in a shoe for warmth to improve the fit. Foot pain is a condition due to inflammation involving any of the bones, ligaments or tendons in the foot and also due to prolong standing and walking. The utilization of Memory Foam Insole (MFI) as a shoe modification for people suffering for foot pain, when using it as an insoles it returns to its original shapes when you apply a force since its elastic and quickly molds to your unique contours. Your weight distribution on your foot speed more evenly and the pressure is relieved on places as well.

Objective: To decrease foot pain for athletes that engage in track and field events improve performance and prevent further foot debilitating condition that hinder their activities of daily livings.

Methodology: The method used is true-experimental and purposive sampling. The T-test is used to test the significant difference between two variable means. The first group will be wearing the (MFI) as insole the second group used an Ordinary Foam Insole (OFI) and both groups also use regular Commercially Available Insoles (CAI) for another 10 days for atleast four hours a day. The collection of data is done every other day. The researchers use the Modified Foot Function Index (MFFI) for evaluation of foot pain.

Results: The MFFI demonstrates that MFI, OFI and CAI has mean average of 1.5, 1.95 and 2.12, respectively with the one-tailed critical t-value of 1.83 at a 0.05 level of significance and a degree of freedom of 9.

Conclusion: There is a significant difference on using the MFI to CAI with a T-stat values for foot pain (2.43), therefore the null hypothesis is rejected and for MFI as to OFI a T-stat value of foot pain (1.71). The null hypothesis is accepted. This shows evidence that MFI is effective in decreasing foot pain as to CAI.

Biography

Charen Rabe from St. Dominic College of Asia, Philippines

raberain@gmail.com

Notes: