

13th Annual Conference on

NEPHROLOGY & RENAL CARE

May 24-25, 2018 Tokyo, Japan

Ten-year outcome study in a cohort of non-metastatic renal cell carcinoma treated with curative intent at SKCH and RC: Survival and progression after radical nephrectomy in 344 patients**Nouman Khan, Arshad Irshad Khalil, Sidra Manzoor, Komal, Azfar Ali, Usman Ahmad, Irfan Ahmad and Khurram Mir**
Shaukat Khanum Memorial Cancer Hospital & Research Centre, Pakistan

Background: Radical Nephrectomy (RN) is a standard treatment of cure for Non-Metastatic Renal Cell Carcinoma (NMRCC). Long-term outcome data are limited for Pakistani population. Our aim was to assess the long-term outcomes of RCC treated with curative intent with Radical Nephrectomy (RN).

Objective: To study the 5 and 10 years outcomes in patients with NMRCC who underwent radical nephrectomy.

Method: This is a retrospective analysis of prospectively collected data between February 2006 and December 2016. We included all the adult patients (age \geq 18 years) with NMRCC from both genders irrespective of their histologic subtypes who underwent radical nephrectomy (RN) with a curative intent. The data were analyzed for overall survival and recurrence rates at 5 and 10 years using Kaplan-Meier survival analysis. Multivariate analysis was done using Cox-regression to identify risk factors associated with poor overall outcome in terms of recurrence and mortality.

Result: 344 patients with 195 (55.5%) males and 149 (44.2%) females with a mean age of 53.5 \pm 14.1 years were followed for a mean follow-up of 31.1 \pm 26.77 months (range: 3-132 months), with 46 (13.4%) deaths. 49 (14.2%) cases had disease recurrence with 33 (9.5%) deaths from disease progression. The 5-year progression-free survival was 37% (95% CI: 49.04-72.76) with the median time to recurrence was 33 months (95% CI: 27.6-38.4) and the median overall survival was 103.7 months (95% CI: 95.7-111.7). The 5-year overall survival was 76.1% (95% CI: 75.2-77) while 10-year survival was 70.8%. There was a significant median survival difference for cases with and without recurrence (log-rank χ^2 : 117.5, $p < 0.001$, T stage, Fuhrman's grade and early postoperative recurrence).

Conclusion: Radical nephrectomy offers the best survival for non-metastatic renal cell carcinoma patients with excellent post-operative survival and progression-free profile.

Biography

Nouman Khan is a Clinical Urooncology Fellow at Shaukat Khanum Memorial Cancer Hospital and Research Centre. He has worked as a Medial Specialist at Institute of Kidney Diseases and Transplant Hospital for a span of 7 months. His skills and areas of interest include surgery, general surgery and healthcare.

dr_noumank@hotmail.com

Notes: