J Nurs Care 2018, Volume 7 DOI: 10.4172/2167-1168-C3-071

3rd World Congress on

NURSING EDUCATION, PRACTICE & RESEARCH

May 16-17, 2018 | Montreal, Canada

Nutritional status of women of reproductive age in a selected char of Rangpur district

Md. Zahid Hasan Khan

Medical Institute, Gaibandha, Bangladesh

An observational cross-sectional study was carried out at Rangpur district in Bangladesh to assess nutritional status of reproductive aged women residing in char area with a sample size 200. Face to face interview was carried out with the semi-structured questionnaire. Convenient sampling technique was used to collect data based on inclusion and exclusion criteria and written consent was taken prior to interview. Nutritional status was determined according to BMI cut off value for Asian population. Descriptive as well as inferential statistics were used to present data. Mean \pm SD age of respondents was 34.27 ± 8.60 . More than half (67%) of the respondents were illiterate and housewife (84%). Mean \pm SD income of respondents was 5700.71 ± 282.89 per month. Underweight, normal and overweight were 67%, 30% and 3% respectively. Most respondents took rice 2-3 times/day. Vegetables and soyabean were taken randomly. Lentil was taken daily. Arthritis, headache, skin disease was more common. Statistical significant association was found between nutritional status and age group (p<0.05), education (p<0.05), occupation (p<0.05) and monthly income (p \pm 0.05). Half of the respondents suffered from underweight and most of them income was very low. Income generating capacity should be increased as well effective nutrition education programme must be instituted.