

29th World Cardiology Conference

November 19-20, 2018 | Edinburgh, Scotland

Comparison of the survival between CABG vs. PCI procedures in patients with poor left ventricular function (ejection fraction <30%): A propensity-matched analysis

Shaneel Shah

University of Bristol, UK

Background: Existing evidence comparing the outcomes of coronary artery bypass graft surgery (CABG) vs. percutaneous coronary intervention (PCI) in patients with poor left ventricular function (LVF) is sparse and flawed. This is largely due to patients with poor LVF being under-represented in major research trials and the outdated nature of some studies which do not consider drug-eluting stent (DES) PCI.

Methods: Following strict inclusion criteria 717 patients who underwent revascularisation by CABG or PCI between 2002 and 2015 were enrolled. 100% of the patients had poor LVF (defined by ejection fraction <30%). By employing a propensity score analysis, 186 suitable matches (93 CABG, 93 PCI) were identified. Several outcomes were evaluated, in the matched population, using data extracted from national registry databases.

Results: CABG patients required a longer length of hospital stay post-revascularisation compared to PCI, 8.91 ± 1.38 and 4.96 ± 1.38 days respectively ($p < 0.0001$). Cox-regression proportional-hazards analysis found that PCI had a higher adjusted 5-year mortality rate (HR 1.752, 95% CI 0.998-3.078, $p = 0.05$). This trend was consistent amongst urgent cases of revascularisation, patients with three or more vessels with coronary artery disease, and cases where, complete revascularisation was achieved. Sub-analysis found the cumulative 5-year survival distribution for PCI with DES to be significantly higher than PCI without DES, but still lower than CABG (log-rank $p = 0.037$; CABG $67.6 \pm 5.3\%$, PCI with DES $54.6 \pm 3.3\%$, PCI without DES is $46.2 \pm 4.7\%$).

Conclusion: Despite a longer length of hospital stay, CABG patients experience a greater post-procedural survival benefit compared to PCI patients. We have demonstrated this at 30 days, 90 days, 1 year, 3 years and 5 years following revascularization. At present, CABG remains a superior revascularisation modality to PCI in patients with poor LVF.

shaneelshah94@gmail.com