8th International Conference and Exhibition on

LASERS, OPTICS & PHOTONICS

November 15-17, 2017 | Las Vegas, USA

Parity detection achieving Heisenberg limit in an SU(1,1) interferometer with coherent and squeezed vacuum input states

Dong Li

Microsystem and Terahertz Research Center, China

One common tool for precision measurement is interferometer. Compared with the conventional SU(2) interferometer, the SU(1,1) interferometer utilizes parametric amplifiers for wave splitting and recombination. Due to parametric amplification process, SU(1,1) interferometers have a better phase sensitivity than SU(2) ones under the same condition of input states. With squeezed vacuum input, the phase measurement sensitivity of SU(1,1) interferometers with coherent mixed with squeezed vacuum input states. Parity detection counts the evenness or oddness of the photon number in one output mode. Our work shows that parity detection reaches below Heisenberg limit when the input coherent and squeezed vacuum light are mixed in roughly equal proportions with a strong parametric amplifier strength. Compared with homodyne detection, parity detection has a slightly better phase sensitivity with coherent and squeezed vacuum inputs and parity detection is more suitable than homodyne detection in some certain situations. Lastly, we also investigate the Quantum Cramer-Rao bound for SU(1,1) interferometers, showing that phase measurement sensitivity does not surpass Quantum Cramer-Rao bound even though it surpasses Heisenberg limit. Parity detection invades SU(1,1) interferometers.

Biography

Dong Li has completed his PhD from East China Normal University. He is the Assistant Researcher of Microsystem and Terahertz Research Center. His research interests include quantum metrology and quantum interferometry.

lidong@mtrc.ac.cn

Notes: