

12th International Conference and Exhibition on **Pharmacovigilance & Drug Safety**
&
22nd International Conference and Exhibition on **Pharmaceutical Formulations**
&
21st Euro-Global Summit on **Toxicology and Applied Pharmacology**

July 04-06, 2019 Valencia, Spain

Pesticides exposure of pregnant women and the metabolism of vitamin D

Lhilali I¹, Zouine N¹, Menouni A^{1,2}, Duca R², Godderris L² and E L Jaafari S¹

¹Moulay Ismail University, Morocco

²Katholieke Universiteit Leuven, Belgium

Introduction: The maternal vitamin D deficiency is a real public health problem associated with adverse effects on the mother and the health of the child throughout life. In Morocco, hypovitaminosis D affects about 90.1% of pregnant women. As a result, the factors that regulate vitamin D metabolism are of paramount importance especially during pregnancy. In fact, prolonged and chronic exposure to phytosanitary products which are known to bioaccumulate in ecosystems and living organisms and their cocktail effect can disrupt this metabolism. However, the relationship between pesticide exposure and hypovitaminosis D in humans and more particularly in pregnant women is rarely studied, as the mechanism of action of pesticides on vitamin D metabolism. This study aims to correlate prenatal exposure to pesticides with the status of vitamin D and to evaluate gene expression of vitamin D metabolism in the placenta.

Methods: In a pilot study among 300 mother-child pairs in Meknes, exposure biomarkers of 30 pesticides (parent compounds and metabolites) belonging to three chemical families of pesticides like organochlorines, organophosphorus compounds and pyrethroids will be measured in urine and blood by HPLC/MS- MS and GC/MS and correlated with serum levels 25(OH)D and 1,25-(OH)₂D the results will be completed by a food frequency questionnaire (FFQ). The analysis of the gene expression of *CYP2R1*, *CYP27B1*, *CYP24A1* and *CTP3A4* in the placenta will be measured using RT-PCR.

Expected Results: Pesticide exposure would alter 25-hydroxyvitamin D concentration and these metabolites in pregnant women and phosphocalcium status in neonates by acting on genes related to vitamin D metabolism via activation of nuclear receptors of xenobiotic metabolism.

Conclusion: This study is first done on the subjects in Morocco. The data will be used to assess whether pesticide exposure is associated with vitamin D deficiency in pregnant women and newborns by taking into account lifestyle and chronic dietary exposure to pesticide residues. Our study aims to explore the genetic and epigenetic regulation of vitamin D in the understanding of action mechanisms of pesticides.

Biography

Lhilali I 2nd year PhD in Biology at the Faculty of Science Moulay Ismail Meknes University, I am a midwife and graduate of the 2nd cycle of paramedical teaching. I currently practice as a paramedical teacher at the Higher Institute of Nursing and Technical Health Professions Lalla Meryem in Meknes. I am also a regional trainer in safe motherhood, GESTA (Labor and Delivery Management, SONU (emergency obstetric and neonatal care) and cardiopulmonary resuscitation of the newborn.

Notes: