

Psychometric Properties of Dynamic Balance, Functional Gait and Dual task walking Outcome Measures used in Stroke Rehabilitation. A Systematic Review.

Kawthar Ajaj

King Saud University, King's College London, UK



Received: November 29, 2021, Editor Assigned: November 30-2021, Reviewed: December 01, 2021, QC No. COS- AA0002;
Proceeding No: Volume: 10, 2022 Published: April 05, 2022, Invoice No. RSTRKGLNB22-1

Abstract

Statement of the Problem: Mobility and balance problems are among the most frequently reported to have disabling effects after stroke. Rehabilitation goals to optimize post stroke functional outcomes and regain normal walking in home and in the community. The use of appropriate, valid, and reliable clinical outcome measures (OMs) is a quality requirement in rehabilitation. This review aims to identify and review psychometric properties of clinical OMs of dynamic balance, functional gait and dual task walking used in stroke rehabilitation to support decision making about appropriate measures for a clinical research study.

Methods: This systematic review was conducted in accordance with PRISMA guidelines. Studies were included if meet the following criteria (1) adults' participants with stroke; (2) assessed dynamic balance and functional walking (walk at a functional level, i.e., crossing the road); (3) used measures that can be applied in clinic; (4) psychometric validation studies (reliability, validity, and responsiveness) of patient performance OMs; (5) published in English with full text available. A systematic literature search of databases was performed. COSMIN checklist was used to determine methodological quality and statistical outcomes.

Results: 43 studies were included, a total of 23 different OMs were identified. Validity and reliability were the psychometric found for most OMs, however data on responsiveness was insufficient. The best psychometric properties for dynamic balance OMs were for the balance evaluation system test (BESTest) and mini-BESTest; for functional walking OMs were the dynamic gait index, functional gait assessment, and timed-up and go test. In the dual task walk OMs, there were a lack of consistency in testing procedure, however all demonstrated high reliability for motor related tasks, slightly less for cognitive related tasks. Conclusion: this review included reliable and valid OMs for dynamic balance, functional gait and dual task walk which help clinician and researchers in selection.

Biography

Kawthar Ajaj, lecturer in King Saud University, Riyadh, research fellow PhD candidate in King's college London. Interested in neurological rehabilitation, specifically in assessment and rehabilitation for regain balance and normal walking to minimize risk of falls.

About University

King's College London is an internationally renowned university delivering exceptional education and world-leading research. We are dedicated to driving positive and sustainable change in society and realizing our vision of making the world a better place.

Our Strategic Vision 2029 looks forward to our 200th anniversary in 2029 and sets out our ambitious plans in five key areas:

Educating the next generation of change-makers;

- Challenging ideas and driving change through research;
- Giving back to society through meaningful service;
- Working with our local communities in London;
- Fostering global citizens with an international perspective.