

Aggressive and Traditional Postoperative Rehabilitation Protocols for Rotator Cuff Repair: A Meta-analysis

Haren Nandapalan*

Department of Neurology, University of Navarra, Pamplona, Spain

Abstract

Introduction: A well-planned rehabilitation programme is just as crucial to a patient's rotator cuff repair procedure's successful completion of tendon healing and the best possible shoulder functional outcome as the size/location of the tear, the surgical approach and the fixation techniques (RCR). There is ongoing debate over the best order to carry out rehabilitation.

Purpose: This meta-objective analysis compares the results of an intense post-operative care regimen versus a traditional rehabilitation programme. The Cochrane Library, CINAHL, PubMed, Ovid MEDLINE and CEPS databases were also searched. We ultimately included six publications that met our criterion for selection.

Results: The traditional strategy improves ROM and shoulder function more than the intensive postoperative rehabilitation method, but it also involves a larger risk of the rotator cuff tendon failing to heal or rupturing again.

Conclusion: Despite the fact that patients with RCR benefit from the rigorous postoperative rehabilitation routine, additional research on the variables impacting the risk of tendon un-healing/re-tearing is needed. These elements must be taken into account while designing a post-operative programme for RCR patients.

Keywords: Rotator cuff repair • Rehabilitation • Aggressive • Range of motion • Function

Introduction

With a prevalence of 13% in adults 50 years and older and a prevalence of 50% at the age of 80, rotator cuff tears are a prevalent cause of shoulder pain and dysfunction. Patients with minor incomplete tears may receive nonoperative care. Surgical repair of the rotator cuff is advised when nonsurgical treatments have failed. The surgical method has a drawback in that different tendon to bone healing capacities exist. In young participants with non-retracted tears, the rate of subsequent tears following repair ranged from 16 percent to 94 percent. The importance and difficulty of providing the best possible rehabilitation after rotator cuff surgery have increased due to the quick development of surgical procedures such as open repair, mini-open repair and arthroscopic repair. The size and location of the rupture, the surgical approach and fixation techniques, as well as a well-planned rehabilitation routine, are crucial elements for optimal tendon healing and a positive functional outcome for the shoulder [1,2].

The length of immobilisation, passive versus active motion and vigorous post-operative treatment are still up for debate when it comes to postoperative rehabilitation programmes. Early use of passive or active range-of-motion exercises, higher dose of a rehabilitation protocol, expedited or intensive rehabilitation protocols, or integrated pre-operative rehabilitation are all examples of aggressive rehabilitation protocols. To avoid postoperative stiffness, it is advised to move passively as soon as possible following surgery. Early continuous passive motion following repair led to a faster recovery in range of motion (ROM) and pain alleviation, as shown by Raab et al. in

1996. However, according to other writers, tendon healing may be aided by immobilisation for 4 to 6 weeks following rotator cuff repair. Early motion following repair also hampered rotator cuff healing, leading to greater rates of unhealed injuries and even rotator cuff re-tearing. However, a recent study demonstrated no adverse effects from early vigorous motion in the fourth postoperative week. The likelihood of an early or aggressive post-operative treatment strategy has recently enhanced because to advancements in surgical procedures [2-4].

Methods

Independently, two physical therapists carried out the initial review. Searches were conducted in a number of databases, including PubMed, Ovid MEDLINE, CINAHL, Cochrane Library and Chinese electronic periodical service (CEPS). Additionally, a manual search of the publications' references was done. Initially, the phrase "rotator cuff repair" was used to search PubMed. The words "aggressive," "early," "progressive," or "accelerated," all of which relate to the meaning of the word "aggressive," and "physical therapy," "rehabilitation," "exercise," or "motion," all of which relate to the meaning of the word "intervention," were used with that statement. Then, the same procedure was followed for the other databases. All searches were restricted to human research [3,5].

Studies that met the following requirements and were published between 2000 and December 2012 were included: (1) They compared the outcomes of an aggressive post-operative rehabilitation protocol with those of a conventional protocol in randomised controlled trials. (2) The study subjects had rotator cuff repairs. (3) The studies used at least one of the following outcome measures: pain, shoulder range of motion, shoulder function, or status of anatomic structure of the rotator cuff tendon. (4) The studies were published in English with the full text. Studies were disregarded if at least one of the following conditions existed in the participants: cervical radiculopathy, autoimmune illness, or metastatic disease; or if the studies lacked the mean, standard deviations, or 95 percent confidence interval for any of these conditions [6-8].

*Address for Correspondence: Haren Nandapalan, Department of Neurology, University of Navarra, Pamplona, Spain, E-mail: nandapalan@gmail.com

Copyright: © 2022 Nandapalan H. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Date of Submission: 01 September, 2022, Manuscript No. ijn-22-81208; **Editor assigned:** 02 September, 2022, PreQC No. P-81208; **Reviewed:** 15 September, 2022, QC No. Q-81208; **Revised:** 22 September, 2022, Manuscript No. R-81208; **Published:** 29 September, 2022, DOI: 10.37421/2376-0281.2022.9.485

Methodological quality evaluation

Two physical therapists each scored each article included in the study using the Physiotherapy Evidence Database (PEDro) scale. The PEDro is frequently used to evaluate the effectiveness of intervention-type randomised controlled trials with sufficient reliability and validity. It consists of 11 questions, each of which receives a yes or no response and awards one point. The maximum score is 10 points because the first question, which is meant to assess the internal validity, is not factored towards the final score. Only when the information is specifically presented in the studies do evaluators respond in the affirmative. The ranges listed below were utilised to categorise the methodological quality: Scores between 9 and 10 indicated outstanding study, 6 to 8 indicated good study, 4 to 5 indicated fair study and less than 4 indicated bad study [9].

Discussion

Six RCTs that examined the outcomes of aggressive rehabilitation programmes with conventional rehabilitation protocols in patients following rotator cuff restoration were included in this meta-analysis. The outcomes of overall ROM at 6 months and 1 year after repair revealed that the aggressive rehabilitation approach was first superior to the conventional treatment and that it also resulted in a larger recovery in shoulder function. Previous research has shown that postoperative shoulder stiffness, the most common consequence following arthroscopic rotator cuff repair, is associated with lower quality of life, higher discomfort and worse shoulder function. The development of soft tissue rigidity, tightness and adhesion is also associated with adhesive capsulitis, pseudotenodesis of the deltoid and complex regional pain syndrome. After surgery, early motion can help prevent stiffness caused by immobilisation.

Although intensive rehabilitation following surgery has many benefits, it should be emphasised that this protocol may carry higher risks of rotator cuff un-healing and re-tear rate than the conventional approach. The disparities between the two methods in our review were almost considerable. Immobilization decreases tendon load-to-failure/stiffness and promotes better tendon to bone repair, according to a number of animal model studies. With smaller biomechanical stresses, overactivity can cause inflammation and boost the formation of scar tissue. Therefore, a number of studies suggested that delaying early motion would be advantageous in an effort to safeguard repaired tendons. There are a few restrictions with our analysis. First, only four studies at most could be pooled to examine each outcome because there were so few articles with varying outcome measures and time points of follow-up. Second, the software was unable to calculate the results of some research because the data did not fit normal distributions. Third, all of the articles were only of fair quality in terms of methodology. This might compromise the accuracy of our analysis [8-10].

Conclusion

According to this meta-analysis, the rigorous postoperative rehabilitation regimen results in better improvements in range of motion (ROM) and shoulder

function than the conventional protocol, but it also carries a higher risk of the rotator cuff tendon failing to repair or rupturing again. The various aspects that affect the rehabilitation protocol must be carefully taken into account when designing any post-operative programme for patients who have had rotator cuff repairs.

Acknowledgement

None.

Conflict of Interest

None

References

1. Arndt, J., P. Clavert, P. Mielcarek, J. Bouchaib and N. Meyer, et al. "Immediate passive motion versus immobilization after endoscopic supraspinatus tendon repair: A prospective randomized study." *Orthop Traumatol Surg Res* 98 (2012): S131-S138.
2. Bassett, Rick W. and Robert H. Cofield. "Acute tears of the rotator cuff. The timing of surgical repair." *Clin Orthop Relat Res* 175 (1983): 18-24.
3. Boileau, Pascal, Nicolas Brassart, Duncan J. Watkinson and Michel Carles, et al. "Arthroscopic repair of full-thickness tears of the supraspinatus: Does the tendon really heal?" *J Bone Joint Surg Am* 87 (2005): 1229-1240.
4. Cuff, Derek J. and Derek R. Pupello. "Prospective randomized study of arthroscopic rotator cuff repair using an early versus delayed postoperative physical therapy protocol." *J Shoulder Elbow Surg* 21 (2012): 1450-1455.
5. Flurin, P. H., P. Landreau, T. Gregory and P. Boileau, et al. "Arthroscopic repair of full-thickness cuff tears: A multicentric retrospective study of 576 cases with anatomical assessment." *Rev Chir Orthop Reparatrice Appar Mot* 91 (2005): 31-42.
6. Ghodadra, Neil S., Matthew T. Provencher, Nikhil N. Verma and Kevin E. Wilk, et al. "Open, mini-open and all-arthroscopic rotator cuff repair surgery: Indications and implications for rehabilitation." *J* 39 (2009): 81-A6.
7. Ghodadra, Neil S., Matthew T. Provencher, Nikhil N. Verma and Kevin E. Wilk, et al. "Open, mini-open and all-arthroscopic rotator cuff repair surgery: Indications and implications for rehabilitation." *J Biomech Eng* 39 (2009): 81-A6.
8. Gimbel, J. A., J. P. Van Kleunen, G. R. Williams and S. Thomopoulos, et al. "Long durations of immobilization in the rat result in enhanced mechanical properties of the healing supraspinatus tendon insertion site." *J Biomech Eng* 129 (2007): 400-404.
9. Parsons, Bradford O., Konrad I. Gruson, Darwin D. Chen and Alicia K. Harrison, et al. "Does slower rehabilitation after arthroscopic rotator cuff repair lead to long-term stiffness?" *J Shoulder Elbow Surg* 19 (2010): 1034-1039.
10. Peltz, Cathryn D., Joseph J. Sarver, LeAnn M. Dourte and Carola C. Würgler Hauri, et al. "Exercise following a short immobilization period is detrimental to tendon properties and joint mechanics in a rat rotator cuff injury model." *J Orthop Res* 28 (2010): 841-845.

How to cite this article: Nandapalan, Haren. "Aggressive and Traditional Postoperative Rehabilitation Protocols for Rotator Cuff Repair: A Meta-analysis." *Int J Neurorehabilitation Eng* 9 (2022): 485.