

ALK Folic Acid Transport and Integrin Signal Induced-Angiogenesis Network in Human Hepatocellular Carcinoma (HCC) by Systems-Theoretical Analysis

Juxiang Huang, Lin Wang*, Minghu Jiang and Hong Lin

Biomedical Center, School of Electronics Engineering, Beijing University of Posts and Telecommunications, Beijing, 100876, China

Abstract

We constructed the high-expression ALK activated transport and signal network in human hepatocellular carcinoma (HCC) compared with low-expression (fold change ≥ 2) no-tumor hepatitis/cirrhotic tissues (HBV or HCV infection) in GEO data set, by using integration of gene regulatory activated and inhibited network inference method with gene ontology (GO) analysis. Our result showed that ALK transport and signal upstream network ECT2, FOLR1, GNAZ, GRM1, ITGA2, LEF1, NR5A1, PTHLH, RIMS3, SORT1, SOX2 activated ALK, and downstream ALK-activated BAP1, CAD, CDH13, CNTNAP2, GRM1, ITGA2, LAPTM4B, MAP2K6, NR5A1, STMN1 in HCC. We obtained that the different biological processes of ALK activated network consisted of folic acid transport, cell surface receptor linked signal transduction, cell-cell signaling, G-protein coupled receptor protein signaling pathway, integrin-mediated signaling pathway, intracellular signaling cascade, low density lipoprotein mediated signaling, Rac protein signal transduction, Rho protein signal transduction in HCC compared with the activated network of no-tumor hepatitis/cirrhotic tissues, as a result of inducing folic acid transport and integrin signal induced-angiogenesis in HCC. Our hypothesis was verified by the different and the same biological processes of ALK activated transport and signal network of HCC compared with the corresponding inhibited network of no-tumor hepatitis/cirrhotic tissues and HCC, respectively.

Keywords: ALK; Human hepatocellular carcinoma (HCC); Folic acid transport and integrin signal induced-angiogenesis network; Systems-theoretical analysis

Introduction

ALK is our identified significant high-expression gene in human hepatocellular carcinoma (HCC) compared with low-expression (fold change ≥ 2) no-tumor hepatitis/cirrhotic tissues (HBV or HCV infection) from GEO data set GSE10140-10141 (<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE10140>, <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE10141>). ALK is related to integral to plasma membrane, membrane; nucleotide binding, transmembrane receptor protein tyrosine kinase activity, receptor signaling protein tyrosine kinase activity, receptor activity, protein binding, ATP binding, transferase activity; protein amino acid phosphorylation, protein amino acid N-linked glycosylation, transmembrane receptor protein tyrosine kinase signaling pathway, nervous system development, brain development (GO database).

Folic acid transport, integrin signal and lipoprotein are involved in angiogenesis. Such as, Folic-acid-mediated inhibition of human colon-cancer cell growth [1]; The Foxc2 transcription factor modulates angiogenesis via induction of integrin beta3 expression [2]; alphavbeta3 integrin and a moody integrin angiogenesis in a changing environment [3]; Active tumor targeting of nonmaterial's using folic acid, integrin receptors and transferring [4]; Role of tetraspanin CD151-alpha3/alpha6 integrin complex in angiogenesis [5]; Integrin affinity modulation in angiogenesis [6]; RECK function of ss1-integrin-dependent in physiologic and tumor angiogenesis [7]; Activation of Ras/MAP kinase is required in high density lipoprotein-induced angiogenesis in human coronary artery endothelial cells [8]; Collateral formation Impairment in lipoprotein(a) transgenic mice therapeutic angiogenesis induced by human hepatocyte growth factor gene [9]; Homocysteine and folic acid effects on angiogenesis and VEGF expression during chicken vascular development [10]; Pharmacological inhibition of integrin alphavbeta3 aggravates experimental liver fibrosis and inhibits hepatic angiogenesis [11]; Lipoprotein contributes to angiogenesis on the chick embryo chorioallantoic membrane [12]; Interaction of alpha9beta1 integrin with thrombospondin-1 promotes angiogenesis [13]; Reconstituted

high-density lipoprotein enhances differentiation of endothelial progenitor cells and stimulates ischemia-induced angiogenesis [14]; Angiogenesis requires Beta1 integrin expression on endothelial cells but not vasculogenesis [15]; Integrin alpha9beta1 directly binds to vascular endothelial growth factor (VEGF)-A and induces VEGF-A-induced angiogenesis [16]; A novel mediator of ovarian angiogenesis follicular fluid high density lipoprotein-associated sphingosine 1-phosphate [17]; Integrin-linked kinase modulates melanoma angiogenesis by activating NF-kappaB/interleukin-6 signaling pathway [18]; Relationship between oxidized lipoprotein, human coronary atherosclerotic plaque stabilization and angiogenesis [19]; Possible roles for folic acid in the modulation of trophoblast invasion and placental development in normal early human pregnancy [20]; Alpha(5)beta(1) integrin ligand PHSRN contributes invasion and alpha(5) mRNA in endothelial cells to stimulate angiogenesis [21]. Yet the distinct high-expression ALK folic acid transport and integrin signal induced-angiogenesis network in HCC remains to be elucidated. Here we constructed the high-expression ALK activated transport and signal network in HCC from GEO data set by gene regulatory network inference method based on linear programming and decomposition procedure.

In this study, we constructed ALK up- and down-stream activated and inhibited transport and signal network in no-tumor hepatitis/cirrhotic tissues and HCC. The biological process and data analysis of

***Corresponding author:** Lin Wang, Biomedical Center, School of Electronics Engineering, Beijing University of Posts and Telecommunications, Beijing, 100876, China, Tel: 0086-13240981826; Fax: 8610-62785736 (Minghu Jiang c/o Lin Wang); E-mail: wanglin98@tsinghua.org.cn

Received November 08, 2011; **Accepted** February 23, 2012; **Published** February 25, 2012

Citation: Huang J, Wang L, Jiang M, Lin H (2012) ALK Folic Acid Transport and Integrin Signal Induced-Angiogenesis Network in Human Hepatocellular Carcinoma (HCC) by Systems-Theoretical Analysis. Mol Biol 1:102. doi:10.4172/2168-9547.1000102

Copyright: © 2012 Huang J, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

the low- and high-expression ALK transport and signal network was done in no-tumor hepatitis/cirrhrotic tissues (HBV or HCV infection) and HCC by GO database. By comparison with the same and different gene ontology (GO) of ALK activated and inhibited transport and signal network between no-tumor hepatitis/cirrhrotic tissues and HCC, we put forwards hypothesis of ALK activated transport and signal network of inducing folic acid transport and integrin signal induced-angiogenesis in HCC.

Materials and Methods

We used microarrays containing 6,144 genes from 25 no-tumor hepatitis/cirrhrotic tissues and 25 HCC patients in GEO data set GSE10140-10141 (<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE10140>, <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE10141>). The raw microarray data was preprocessed by log base 2.

Significant expressed genes of ALK transport and signal network were identified using significant analysis of microarrays (SAM) (<http://wwwstat.stanford.edu/~tibs/SAM/>) [22]. We selected two classes unpaired and minimum fold change ≥ 2 and chose the significant highly expressed value genes of HCC compared with that of no-tumor hepatitis/cirrhrotic tissues (HBV or HCV infection) under the false-discovery rate and q-value were 0%. The q-value is like the well-known P-value, but adapted to multiple-testing situations.

ALK transport and signal network was constructed based on GRNInfer and GVedit tools (<http://www.graphviz.org/About.php>). GRNInfer is a novel mathematic method called GNR (Gene Network Reconstruction tool) based on linear programming and a decomposition procedure for inferring gene network [23]. We established ALK activated network of HCC based on the fold change ≥ 2 distinguished genes and selected parameters as lambda 0.0 because we used one data set. Lambda was a positive parameter which balanced the matching and sparsity terms in the objective function. Using different thresholds, we could predict various networks with the different edge density. The threshold parameters make the edge whose strength of link is smaller than threshold not shown in the network graph. The smaller this parameter, the more edges in the network graph. We selected threshold $1.0e-7$.



ALK transport and signal network of HCC was analyzed using Molecule Annotation System, MAS (CapitalBio Corporation, Beijing, China; <http://bioinfo.capitalbio.com/mas3/>). MAS is a Web-based software toolkit for a whole data mining and function annotation solution to extract and analyze biological molecules relationships from public databases. The primary databases of MAS integrated various well-known biological resources, such as Gene Ontology (<http://www.geneontology.org>), KEGG (<http://www.genome.jp/kegg/>), BioCarta (<http://www.biocarta.com/>), GenMapp (<http://www.genmapp.org/>), HPRD (<http://www.hprd.org/>), MINT (<http://mint.bio.uniroma2.it/mint/Welcome.do>), BIND (<http://www.blueprint.org/>), Intact (<http://www.ebi.ac.uk/intact/>), UniGene (www.ncbi.nlm.nih.gov/UniGen), OMIM (<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=OMIM>) and disease (<http://bioinfo.capitalbio.com/mas3/>). MAS offers various query entries and graphics. The algorithm is P, Q value in GO and pathway of module was presented in reference [24].

Results

We constructed ALK up- and down-stream activated and inhibited transport and signal network in no-tumor hepatitis/cirrhrotic tissues and HCC from our total network of 225 significant high-expression

molecules (fold change ≥ 2) from 6,144 genes of 25 HCC compared with 25 no-tumor hepatitis/cirrhrotic tissues (HBV or HCV infection) by GRNInfer, respectively.

We extracted the biological process of GO terms and did numbers data analysis of the different biological processes of ALK activated transport and signal network in HCC compared with activated network of no-tumor hepatitis/cirrhrotic tissues, the same biological processes of ALK activated transport and signal network in HCC compared with



| The Different ALK Activated Transport and Signal Network of HCC compared with Activated Network of No-tumor Hepatitis/cirrhotic Tissues | | | |
|---|---------|--|---------|
| Terms | Numbers | Terms | Numbers |
| folic acid transport | 1 | Rho protein signal transduction | 1 |
| cell-cell signaling | 3 | low density lipoprotein mediated signaling | 1 |
| integrin-mediated signaling pathway | 2 | cell surface receptor linked signal transduction | 1 |
| intracellular signaling cascade | 3 | G-protein coupled receptor protein signaling pathway | 3 |
| Rac protein signal transduction | 1 | | |
| The Same ALK Activated Transport and Signal Network of HCC compared with Inhibited Network of No-tumor Hepatitis/cirrhotic Tissues | | | |
| Terms | Numbers | Terms | Numbers |
| transport | 2 | cell surface receptor linked signal transduction | 1 |
| G-protein signaling | 1 | signal transduction | 7 |
| cell-cell signaling | 3 | | |

Table 1: GO Terms and numbers data analysis of the different biological processes of ALK activated transport and signal network of HCC compared with activated network of no-tumor hepatitis/cirrhotic tissues, the same biological processes of ALK activated transport and signal network of HCC compared with inhibited network of no-tumor hepatitis/cirrhotic tissues.

| The Different ALK Activated Transport and Signal Network of HCC compared with Inhibited Network of HCC | | | |
|--|---------|---|---------|
| Terms | Numbers | Terms | Numbers |
| endosome to lysosome transport | 1 | intracellular signaling cascade | 3 |
| folic acid transport | 1 | integrin-mediated signaling pathway | 2 |
| Golgi to endosome transport | 1 | Rac protein signal transduction | 1 |
| plasma membrane to endosome transport | 1 | Rho protein signal transduction | 1 |
| endosome transport via multivesicular body sorting pathway | 1 | low density lipoprotein mediated signaling | 1 |
| cell surface receptor linked signal transduction | 1 | induction of apoptosis by extracellular signals | 1 |

Table 2: GO Terms and numbers data analysis of the different biological processes of ALK activated transport and signal network of HCC compared with inhibited network of HCC.

inhibited network of no-tumor hepatitis/cirrhotic tissues, as shown in Table 1. GO terms and numbers data was analyzed the different biological processes of ALK activated compared with inhibited transport and signal network in HCC, as shown in Table 2.

ALK activated transport and signal network was constructed in HCC. Our result showed that upstream ECT2, FOLR1, GNAZ, GRM1, ITGA2, LEF1, NR5A1, PTHLH, RIMS3, SORT1, SOX2 activated ALK, and the downstream ALK-activated BAP1, CAD, CDH13, CNTNAP2, GRM1, ITGA2, LAPTM4B, MAP2K6, NR5A1, STMN1 in HCC, as shown in Figure 1 and Figure 2.

Discussion

Our aim is to construct, interpret, verify and predict the function of novel high-expression ALK folic acid transport and integrin signal induced-angiogenesis network in HCC. We have already constructed and analyzed some novel molecular network from different databases presented in our articles [24-37]. In this study, we constructed ALK up- and down-stream activated and inhibited transport and signal network in no-tumor hepatitis/cirrhotic tissues and HCC. The biological process and data analysis of the low- and high-expression ALK transport and signal network was done in no-tumor hepatitis/cirrhotic tissues (HBV or HCV infection) and HCC by GO database. By comparison with the same and different gene ontology (GO) of ALK activated and inhibited transport and signal network between no-tumor hepatitis/cirrhotic tissues and HCC, we put forwards hypothesis of ALK activated transport and signal network of inducing folic acid transport and integrin signal induced-angiogenesis in HCC.

We extracted the biological process of GO terms and did numbers data analysis of the different biological processes of ALK activated transport and signal network in HCC compared with activated network of no-tumor hepatitis/cirrhotic tissues, the same biological processes of ALK activated transport and signal network in HCC compared with inhibited network of no-tumor hepatitis/cirrhotic tissues (Table

1 and Table 2). We constructed the high-expression ALK activated transport and signal network in human hepatocellular carcinoma (HCC) compared with low-expression (fold change ≥ 2) no-tumor hepatitis/cirrhotic tissues (HBV or HCV infection) in GEO data set using integration of gene regulatory network inference method. Our result showed that ALK transport and signal upstream network ECT2, FOLR1, GNAZ, GRM1, ITGA2, LEF1, NR5A1, PTHLH, RIMS3, SORT1, SOX2 activated ALK, and downstream ALK-activated BAP1, CAD, CDH13, CNTNAP2, GRM1, ITGA2, LAPTM4B, MAP2K6, NR5A1, STMN1 (Figure 1 and Figure 2) in HCC.

By further comparison with the same and different gene ontology (GO) of ALK activated and inhibited transport and signal network between no-tumor hepatitis/cirrhotic tissues and HCC, we obtained that the different biological processes of ALK activated network consisted of folic acid transport, cell surface receptor linked signal transduction, cell-cell signaling, G-protein coupled receptor protein signaling pathway, integrin-mediated signaling pathway, intracellular signaling cascade, low density lipoprotein mediated signaling, Rac protein signal transduction, Rho protein signal transduction in HCC compared with activated network of no-tumor hepatitis/cirrhotic tissues, as a result of inducing folic acid transport and integrin signal induced-angiogenesis in HCC.

The same biological processes of ALK activated network included transport, cell surface receptor linked signal transduction, cell-cell signaling, G-protein signaling, signal transduction in HCC compared with inhibited network of no-tumor hepatitis/cirrhotic tissues. It is consistent with the different biological processes of ALK activated transport and signal network of HCC compared with activated network of no-tumor hepatitis/cirrhotic tissues.

The different biological processes of ALK activated network consisted of endosome to lysosome transport, endosome transport via multivesicular body sorting pathway, folic acid transport, Golgi to

endosome transport, plasma membrane to endosome transport, cell surface receptor linked signal transduction, induction of apoptosis by extracellular signals, integrin-mediated signaling pathway, intracellular signaling cascade, low density lipoprotein mediated signaling, Rac protein signal transduction, Rho protein signal transduction in HCC compared with inhibited network of HCC. It is consistent with the different biological processes of ALK activated transport and signal network of HCC compared with activated network of no-tumor hepatitis/cirrhotic tissues, the same biological processes of ALK activated transport and signal network of HCC compared with inhibited network of no-tumor hepatitis/cirrhotic tissues, respectively.

Therefore, our ALK activated folic acid transport and integrin signal induced-angiogenesis hypothesis was verified by the different and the same biological processes of ALK activated transport and signal network of HCC compared with the corresponding inhibited network of no-tumor hepatitis/cirrhotic tissues and HCC, respectively.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (No.60871100) and (No.61171114), the Returned Overseas Chinese Scholars for Scientific research Foundation of State Education Ministry, Significant Science and Technology Project for New Transgenic Biological Species (2009ZX08012-001B), Automatical Scientific Planning of Tsinghua University (20111081023 and 20111081010), State Key Lab of Pattern Recognition Open Foundation.

References

- Duthie SJ (2001) Folic-acid-mediated inhibition of human colon-cancer cell growth. *Nutrition* 17: 736-737.
- Hayashi, H, Sano H, Seo S, Kume T (2008) The Foxc2 transcription factor regulates angiogenesis via induction of integrin beta3 expression. *J Biol Chem* 283: 23791-23800.
- Hodivala-Dilke K (2008) alphavbeta3 integrin and angiogenesis: a moody integrin in a changing environment. *Curr Opin Cell Biol* 20: 514-519.
- Kolhatkar R, Lote A, Khambati H (2011) Active tumor targeting of nanomaterials using folic acid, transferrin and integrin receptors. *Curr Drug Discov Technol* 8: 197-206.
- Liu, W F, Zuo HJ, Chai BL, Peng D, Fei YJ, et al. (2011) Role of tetraspanin CD151- α 3/ α 6 integrin complex: Implication in angiogenesis CD151-integrin complex in angiogenesis. *Int J Biochem Cell Biol* 43: 642-650.
- Mahabeleshwar GH, Chen J, Feng W, Somanath PR, Razorenova, et al. (2008) Integrin affinity modulation in angiogenesis. *Cell Cycle* 7: 335-347.
- Miki T, Shamma A, Kitajima S, Takegami Y, Noda M, et al. (2010) The β 1-integrin-dependent function of RECK in physiologic and tumor angiogenesis. *Mol Cancer Res* 8: 665-676.
- Miura S, Fujino M, Mastsuo Y, Kawamura A, Tanigawa H, et al. (2003) High density lipoprotein-induced angiogenesis requires the activation of Ras/MAP kinase in human coronary artery endothelial cells. *Arterioscler Thromb Vasc Biol* 23: 802-808.
- Morishita R, Sakaki M, Yamamoto K, Iguchi S, Aoki M, et al. (2002) Impairment of collateral formation in lipoprotein(a) transgenic mice: therapeutic angiogenesis induced by human hepatocyte growth factor gene. *Circulation* 105: 1491-1496.
- Oosterbaan AM, Steegers EA, Ursem NT (2012) The effects of homocysteine and folic acid on angiogenesis and VEGF expression during chicken vascular development. *Microvasc Res* 83: 98-104.
- Patsenker E, Popov Y, Stickel F, Schneider V, Ledermann M, et al. (2009) Pharmacological inhibition of integrin alphavbeta3 aggravates experimental liver fibrosis and suppresses hepatic angiogenesis. *Hepatology* 50: 1501-1511.
- Ribatti D, Vacca A, Giachetta F, Cesaretti S, Anichini M, et al. (1998) Lipoprotein (a) induces angiogenesis on the chick embryo chorioallantoic membrane. *Eur J Clin Invest* 28: 533-537.
- Staniszewska I, Zaveri S, Del Valle L, Oliva I, Rothman VL, et al. (2007) Interaction of alpha9beta1 integrin with thrombospondin-1 promotes angiogenesis. *Circ Res* 100: 1308-1316.
- Sumi M, Sata M, Miura S, Rye KA, Toya N, et al. (2007) Reconstituted high-density lipoprotein stimulates differentiation of endothelial progenitor cells and enhances ischemia-induced angiogenesis. *Arterioscler Thromb Vasc Biol* 27: 813-818.
- Tanjore H, Zeisberg EM, Gerami-Naini B, Kalluri R (2008) Beta1 integrin expression on endothelial cells is required for angiogenesis but not for vasculogenesis. *Dev Dyn* 237: 75-82.
- Vlahakis NE, Young BA, Atakilit A, Hawkrigde AE, Issaka RB, et al. (2007) Integrin alpha9beta1 directly binds to vascular endothelial growth factor (VEGF)-A and contributes to VEGF-A-induced angiogenesis. *J Biol Chem* 282: 15187-15196.
- von Otte S, Paletta JR, Becker S, König S, Fobker M, et al. (2006) Follicular fluid high density lipoprotein-associated sphingosine 1-phosphate is a novel mediator of ovarian angiogenesis. *J Biol Chem* 281: 5398-5405.
- Wani AA, Jafarnejad SM, Zhou J, Li G (2011) Integrin-linked kinase regulates melanoma angiogenesis by activating NF- κ B/interleukin-6 signaling pathway. *Oncogene* 30: 2778-2788.
- Wi, LX, Tang QH, Sun L, Shi HY, Guo AT, et al. (2006) [Relationship between oxidized lipoprotein, angiogenesis and human coronary atherosclerotic plaque stabilization]. *Zhonghua Bing Li Xue Za Zhi* 35: 138-141.
- Williams PJ, Bulmer JN, Innes BA, Broughton Pipkin F (2011) Possible roles for folic acid in the regulation of trophoblast invasion and placental development in normal early human pregnancy. *Biol Reprod* 84: 1148-1153.
- Zeng ZZ, Yao H, Staszewski ED, Rockwood KF, Markwart SM, et al. (2009) alpha(5)beta(1) Integrin Ligand PHSRN Induces Invasion and alpha(5) mRNA in Endothelial Cells to Stimulate Angiogenesis. *Transl Oncol* 2: 8-20.
- Storey JD (2002) A direct approach to false discovery rates. *J Roy Stat Soc Ser B* 64: 479-498.
- Wang Y, Joshi T, Zhang XS, Ku D, Chen L, et al. (2006) Inferring gene regulatory networks from multiple microarray datasets. *Bioinformatics* 22: 2413-2420.
- Wang L, Sun L, Huang J, Jiang M (2011) Cyclin-dependent kinase inhibitor 3 (CDKN3) novel cell cycle computational network between human non-malignancy associated hepatitis/cirrhosis and hepatocellular carcinoma (HCC) transformation. *Cell Prolif* 44: 291-299.
- Huang J, Wang L, Jiang M, Zheng X (2010) Interferon α -Inducible Protein 27 Computational Network Construction and Comparison between the Frontal Cortex of HIV Encephalitis (HIVE) and HIVE-Control Patients The Open Genomics Journal 3: 1-8.
- Huang JX, Wang L, Jiang MH (2010) TNFRSF11B computational development network construction and analysis between frontal cortex of HIV encephalitis (HIVE) and HIVE-control patients. *J Inflamm (Lond)* 7: 50.
- Sun L, Wang L, Jiang M, Huang J, Lin H (2011) Glycogen debranching enzyme 6 (AGL), enolase 1 (ENOSF1), ectonucleotide pyrophosphatase 2 (ENPP2_1), glutathione S-transferase 3 (GSTM3_3) and mannosidase (MAN2B2) metabolism computational network analysis between chimpanzee and human left cerebrum. *Cell Biochem Biophys* 61: 493-505.
- Sun Y, Wang L, Jiang M, Huang J, Liu Z, et al. (2010) Secreted phosphoprotein 1 upstream invasive network construction and analysis of lung adenocarcinoma compared with human normal adjacent tissues by integrative biocomputation. *Cell Biochem Biophys* 56: 59-71.
- Sun Y, Wang L, Lui L (2008) Integrative decomposition procedure and Kappa statistics set up ATF2 ion binding module in Malignant Pleural Mesothelioma (MPM). *Frontiers of Electrical and Electronic Engineering in China* 3: 381-387.
- Wang L, Huang J, Jiang M (2011) CREB5 computational regulation network construction and analysis between frontal cortex of HIV encephalitis (HIVE) and HIVE-control patients. *Cell Biochem Biophys* 60: 199-207.
- Wang L, Huang J, Jiang M (2010) RRM2 computational phosphoprotein network construction and analysis between no-tumor hepatitis/cirrhotic liver tissues and human hepatocellular carcinoma (HCC). *Cell Physiol Biochem* 26: 303-310.
- Wang L, Huang J, Jiang M, Lin H (2012) Signal Transducer and Activator of Transcription 2 (STAT2) Metabolism Coupling Postmitotic Outgrowth to Visual and Sound Perception Network in Human Left Cerebrum by Biocomputation. *J Mol Neurosci*.
- Wang, L, Huang J, Jiang M, Sun L (2011) Survivin (BIRC5) cell cycle computational network in human no-tumor hepatitis/cirrhosis and hepatocellular carcinoma transformation. *J Cell Biochem* 112: 1286-1294.

-
34. Wang L, Huang J, Jiang M, Sun L (2011) MYBPC1 computational phosphoprotein network construction and analysis between frontal cortex of HIV encephalitis (HIVE) and HIVE-control patients. *Cell Mol Neurobiol* 31: 233-241.
35. Wang L, Huang J, Jiang M, Zheng X (2010) AFP computational secreted network construction and analysis between human hepatocellular carcinoma (HCC) and no-tumor hepatitis/cirrhotic liver tissues. *Tumour Biol* 31: 417-425.
36. Wang L, Sun Y, Jiang M, Zhang S, Woll S (2009) FOS proliferating network construction in early colorectal cancer (CRC) based on integrative significant function cluster and inferring analysis. *Cancer Invest* 27: 816-824.
37. Wang, L, Sun Y, Jiang M, Zheng X (2009) Integrative decomposition procedure and Kappa statistics for the distinguished single molecular network construction and analysis. *J Biomed Biotechnol* 2009: 726728.