

An Efficient and Operationally Simple Synthesis of Some New Chalcones by Using Grinding Technique

S Zangade, S Mokle, A Vibhute, Y Vibhute*

Laboratory of Organic Synthesis, PG Department of Chemistry, Yeshwant Mahavidyalaya, Nanded-431602 (MS), India.

*Correspondence to: Yeshwant Vibhute, drybv@rediffmail.com

Accepted: October 21, 2010; Published: February 2, 2011

Abstract

An efficient and operationally simple reaction is shown between substituted 2-acetyl-1-naphthol/2-acetyl-1-naphthol and different substituted benzaldehydes in presence of base afford chalcones in quantitative yield using grindstone technique. Mild reaction condition, no need of catalyst, non-hazardous and environmentally safer, giving excellent yield in short reaction time, are notable advantages of this method.

Keywords: Substituted benzaldehydes; 2-acetyl-1-naphthol/substituted 2-acetyl-1-naphthol; Claisen-Schmidt; chalcones; grindstone chemistry.

1. Introduction

The utility of chalcones due to their usefulness as in synthesis of various heterocyclic compounds, as plant origin [1] and exhibit antimalarial [2], antibacterial [3], antifibrogenic [4], anticancer [5], antitrichomonal [6], antiinflammatory [7], antileishmanial[8], cytotoxic and anti- *Trypanosoma cruzi* [9] activities.

A classical method for synthesis of chalcones is claisen-schmidt condensation in which aldehyde reacted with acetophenone in the presence of aqueous alkaline bases [10], Ba(OH)₂ / LiOH [11]. Chalcones also synthesized by using microwave irradiation, ultrasound irradiation [12] and by Suzuki reaction [13]. Recently various modified methods for synthesis of chalcones has been reported using different catalyst such as SOCl₂ [14], natural phosphate lithium nitrate [15], KF / natural phosphate [16], acyclic acidic ionic liquid [17], Na₂CO₃ [18], high temperature water [19], silica-sulphuric acid [20], ZrCl₄ and ionic liquid [21], silica-sulphuric acid [22], NaOH-Al₂O₃ [23] and silica chloride [24].

A classical synthesis of these compounds involves the condensation of acetophenones and aldehydes to give chalcones. The combination of solvents and long reaction time, costly chemicals / catalyst makes this method environmentally hazardous. This provided the stimulus to synthesize some new chalcones using grindstone technique [25]. In grindstone technique, reaction occurs through generation of local heat by grinding of crystals of substrate and reagent by mortar and pestle. Reactions are initiated by grinding, with the small amount of energy through friction. In some cases, a mixture and reagents turns to a glassy material. Such reaction are simple to handle, reduce pollution, comparatively cheaper to operate and may be regarded as more economical and ecologically favorable procedure in chemistry [26]. Solid-state reaction occurs more efficiently and more selectively than does the solution reaction, since molecules in the crystal are arranged tightly and regularly [27].

In present work, grindstone technique was used for the synthesis of titled compounds. This method is superior to conventional method; since it is eco-friendly, high yielding, requires no special apparatus, non-hazardous, operationally simple and convenient.

Newly synthesized chalcones were prepared by grinding together equivalent amounts of the appropriate 2-acetyl-1-naphthol / substituted 2-acetyl-1-naphthol and different substituted benzaldehydes in presence of solid KOH in a porcelain mortar under solvent-free conditions for 4-8 min. On completion of reaction (TLC), the reaction mixture was diluted with cold water, neutralized by dilute HCl and recrystallised from acetic acid.

2. Methods

2.1 Chemistry

Melting points were determined in an open capillary tube and are uncorrected. IR spectra were recorded in KBr on a Perkin-Elmer spectrometer. ¹H NMR spectra were recorded on a Gemini 300-MHz instrument in DMSO as solvent and TMS as an internal standard. The mass spectra were recorded on EISHIMADZU-GC-MS spectrometer. Elemental analysis was carried out on a Carlo Erba 1108 analyzer. The purity of products was checked by Thin Layer Chromatography (TLC) on silica gel.

2.2 General procedure for synthesis of chalcones

2-acetyl-1-naphthol / substituted 2-acetyl-1-naphthol (0.01mmol), different substituted benzaldehydes (0.01mmol) and solid palletete of KOH (0.02mmol) were taken in mortar and grind for several minute (Table-1). On completion of grinding as monitored by TLC, the obtained solid mixture was diluted with cold water, neutralized by dil HCl and recrystlized from acetic acid to give the corresponding chalcone derivatives.

3-(3,4-Dimethoxy-phenyl)-1-(1-hydroxy-naphthalen-2-yl)-propenone (3a): IR (ν max) cm^{-1} : 3255 (OH), 1629 (C=O), 1569, 1495 (ring C=C), $^1\text{H NMR}$ (300 MHz, DMSO) δ 3.73 (s, 6H, two- OCH₃), δ 6.90 (d, J=16 Hz 1H, H_a), δ 7.56 (d, J=16 Hz 1H, H_b), δ 7.35-8.08 (m, 9H, Ar-H), δ 12.8 (s, 1H, OH). MS m/z: 334 (M⁺). Anal. Calc for C₂₁H₁₈O₄: C, 75.44; H, 5.38. Found: C, 75.64; H, 5.54.

1-(1-Hydroxy-naphthalen-2-yl)-3-(4-methoxy-phenyl)-propenone (3b): IR (ν max) cm^{-1} : 3252 (OH), 1626 (C=O), 1567, 1490 (ring C=C), $^1\text{H NMR}$ (300 MHz, DMSO) δ 3.75 (s, 3H, - OCH₃), δ 6.82 (d, J= 15.5 Hz 1H, H_a), δ 7.52 (d, J= 15.5 Hz 1H, H_b), δ 7.30-8.10 (m, 10H, Ar-H), δ 13.1 (s, 1H, OH). MS m/z: 304 (M⁺). Anal. Cacl for C₂₀H₁₆O₃: C, 78.94; H, 5.26. found: C, 79.10; H, 5.45.

1-(1-Hydroxy-4-iodo-naphthalen-2-yl)-3-(4-methoxy-phenyl)-propenone (3c): IR (ν max) cm^{-1} : 3250 (OH), 1625 (C=O), 1560, 1480 (ring C=C), $^1\text{H NMR}$ (300 MHz, DMSO) δ 3.70 (s, 3H, - OCH₃), δ 6.80 (d, J= 17 Hz 1H, H_a), δ 7.60 (d, J= 17 Hz 1H, H_b), δ 7.30-8.12 (m, 8H, Ar-H), δ 7.70 (s, 1H, Ar-H), δ 13.2 (s, 1H, OH). MS m/z: 430 (M⁺). Anal. Cacl for C₂₀H₁₅O₃I: C, 55.81; H, 3.48; I, 29.53. found: C, 56.00; H, 3.60; I, 29.65.

1-(4-Bromo-1-hydroxy-naphthalen-2-yl)-3-(3,4-dimethoxy-phenyl)-propenone (3d): IR (ν max) cm^{-1} : 3258 (OH), 1629 (C=O), 1557, 1485 (ring C=C), $^1\text{H NMR}$ (300 MHz, DMSO) δ 3.70 (s, 6H, two - OCH₃), δ 6.92 (d, J= 16 Hz 1H, H_a), δ 7.54 (d, J= 16 Hz 1H, H_b), δ 7.35-8.10 (m, 7H, Ar-H), δ 7.73 (s, 1H, Ar-H), δ 13.0 (s, 1H, OH). MS m/z: 413 (M⁺). Anal. Cacl for C₂₁H₁₇O₄Br: C, 61.01; H, 4.11; Br, 19.37. found: C, 61.10; H, 4.24; I, 19.51.

1-(4-Bromo-1-hydroxy-naphthalen-2-yl)-3-(3,4,5-trimethoxy-phenyl)-propenone (3e): IR (ν max) cm^{-1} : 3260 (OH), 1626 (C=O), 1557, 1490 (ring C=C), $^1\text{H NMR}$ (300 MHz, DMSO): δ 3.71 (s, 9H, 3 X -OCH₃), δ 6.86 (d, J= 15.5 Hz 1H, H_a), δ 7.56 (d, J= 15.5 Hz 1H, H_b), δ 7.36-8.13 (m, 7H, Ar-H), δ 13.0 (s, 1H, OH). MS m/z: 443 (M⁺). Anal. Cacl for C₂₂H₁₉O₅Br: C, 55.59; H, 4.28; Br, 18.0. found: C, 55.70; H, 4.45; Br, 18.08.

3-(4-Chloro-phenyl)-1-(1-hydroxy-naphthalen-2-yl)-propenone (3f): IR (ν max) cm^{-1} : 3257 (OH), 1625 (C=O), 1572, 1490 (C=C). $^1\text{H NMR}$ (300 MHz, DMSO): δ 6.84 (d, J= 17 Hz 1H, H_a), δ 7.52 (d, J= 17 Hz 1H, H_b), δ 7.36-8.10 (m, 10H, Ar-H), δ 12.9 (s, 1H, OH). MS m/z: 308 (M⁺). Anal. Cacl for C₁₉H₁₃O₂Cl: C, 74.02; H, 4.22; Cl, 11.36. found: C, 74.10; H, 4.45; Cl, 11.48.

3-(4-Chloro-phenyl)-1-(1-hydroxy-4-iodo-naphthalen-2-yl)-propenone (3g): IR (ν max) cm^{-1} : 3250 (OH), 1628 (C=O), 1574, 1488 (C=C). $^1\text{H NMR}$ (300 MHz, DMSO): δ 6.88 (d, J= 16 Hz 1H, H_a), δ 7.58 (d, J= 16 Hz 1H, H_b), δ 7.38-8.10 (m, 8H, Ar-H), δ 7.78 (s, 1H, Ar-H), δ 13.1 (s, 1H, OH). MS m/z: 434 (M⁺). Anal. Cacl for C₁₉H₁₂O₂ClI: C, 52.53; H, 2.76; X (I+Cl), 37.32. found: C, 52.70; H, 2.88; X (I+Cl), 37.47.

1-(4-Bromo-1-hydroxy-naphthalen-2-yl)-3-(4-fluoro-phenyl)-propenone (3h): IR (ν max) cm^{-1} : 3255 (OH), 1629 (C=O), 1570, 1492 (C=C). $^1\text{H NMR}$ (300 MHz, DMSO): δ 6.86 (d, J= 15.5 Hz 1H, H_a), δ 7.54 (d, 15.5 Hz 1H, H_b), δ 7.37-8.12 (m, 9H, Ar-H), δ 13.2 (s, 1H, OH). MS m/z: 371 (M⁺). Anal. Cacl for C₁₉H₁₂O₂BrF: C, 61.45; H, 3.23; X (Br+F), 26.68. found: C, 61.54; H, 3.38; X, (Br+F), 26.78.


3-(4-Fluoro-phenyl)-1-(1-hydroxy-naphthalen-2-yl)-propenone (3i): IR (ν max) cm^{-1} : 3260 (OH), 1628 (C=O), 1574, 1486 (C=C). $^1\text{H NMR}$ (300 MHz, DMSO): δ 6.82 (d, J= 17 Hz 1H, H_a), δ 7.52 (d, J= 17 Hz 1H, H_b), δ 7.35-8.09 (m, 10H, Ar-H), δ 13.2 (s, 1H, OH). MS m/z: 292 (M⁺). Anal. Cacl for C₁₉H₁₃O₂F: C, 78.08; H, 4.45; F, 6.16. found: C, 78.19; H, 4.56; F, 6.28.

1-(1-Hydroxy-4-iodo-naphthalen-2-yl)-3-(3,4,5-trimethoxy-phenyl)-propenone (3j): IR (ν max) cm^{-1} : 3260 (OH), 1626 (C=O), 1557, 1490 (ring C=C), $^1\text{H NMR}$ (300 MHz, DMSO): δ 3.74 (s, 9H, three - OCH₃), δ 6.86 (d, J= 16 Hz 1H, H_a), δ 7.56 (d, J= 16 Hz 1H, H_b), δ 7.36-8.13 (m, 6H, Ar-H), δ 7.78 (s, 1H, Ar-H), δ 13.2 (s, 1H, OH). MS m/z: 490 (M⁺). Anal. Cacl for C₂₂H₁₉O₅I: C, 53.87; H, 3.87; I, 25.91. found: C, 54.02; H, 4.01; I, 26.04.

3. Results and Discussion

In continuation of our earlier research works on solvent-free organic synthesis [28, 29] herein, we would like to report a simple, efficient and solvent-free synthesis of some new chalcones by grinding method (Scheme-I). Claisen-Schmidt condensation of 2-acetyl-1-naphthol / substituted 2-acetyl-1-naphthol (**1a-j**) with various substituted benzaldehydes (**2a-j**) in the presence of solid KOH in combination with grinding at room temperature in the absence of solvent afforded chalcones (**3a-j**). The yields of the products are excellent and purity is high as compared to conventional method. The process is simple, efficient, and economical. It is also consistent with the green chemistry approach because it does not need heating or microwave irradiation. The structures of the newly synthesized chalcones (**3a-j**) were established by their elemental analysis and spectral data (IR, ¹HNMR and MS).

Scheme



3a: R, R₁= H, R₂, R₃= OCH₃

3b: R, R₁, R₃= H, R₂= OCH₃

3c: R=I, R₁, R₃=H, R₂=OCH₃

3d: R=Br, R₁=H, R₂, R₃= OCH₃

3e: R=Br, R₁, R₂, R₃= OCH₃

3f: R, R₁, R₃= H, R₂= Cl

3g: R= I, R₁, R₃= H, R₂= Cl

3h: R= Br, R₁, R₃= H, R₂= F

3i: R, R₁, R₃= H, R₂= F

3j: R=H, R₁, R₂, R₃= OCH₃

Table-1: Physical data of Synthesized Chalcones.

Entry	M.P. °C	Yield (%)	Time (min.)
3a	154	88	4.5
3b	188	92	6.5
3c	259	85	7.0
3d	128	90	5.5
3e	142	87	5.0
3f	178	91	4.0
3g	227	94	5.0
3h	247	87	6.5
3i	196	93	8.0
3j	278	94	5.0

4. Conclusion

In summary, we have developed a simple, efficient and more eco-friendly method for synthesis of chalcones by grinding technique. The notable advantages of present method are no organic solvent required (except for the product recrystallisation), waste minimization, simple operation, clean reaction profile, easy work-up, shorter reaction time (4-8 min.), high yields (84-95 %) and eco-friendly as compared to conventional method.

Competing Interests

The authors declare that they have no competing interests.

Authors' Contributions

Acknowledgement

Authors are thankful to Principal, Yeshwant Mahavidyalaya, Nanded for providing laboratory facilities and to Director of IICT, Hyderabad for providing spectra of samples. Authors are also thankful to University Grants Commission, New Delhi for sanctioning major research grant.

References

[1] Hijova E, 2006. Bioavailability of chalcones. Bratisl Lek Listy, 107: 80-84.

- [2] Liu M, Wilairat P, Mei-Lin G, 2001. Antimalarial aloylated and hydroxylated chalcones: structure activity relationship analysis. *Journal of Medicinal Chemistry*, 44: 4443-4452.
- [3] Opletalova V, 2000. Chalcones and their heterocyclic analogues as potential therapeutic agents of bacterial diseases. *Ceska Slov Farm*, 49: 278-284.
- [4] Lee SH, Nan JX, Zhao YZ, Woo SW, Park EJ, Kang TH, Seo GS, Kim YC, Sohn DH, 2003. The Chalcones Butein from *Rhus verniciflua* shows antifibrogenic activity. *Planta*, 69: 990-994.
- [5] Konieczny MT, Konieczny W, Sabisz M, Skladanowski A, Wakiec R, Augusttynowiczkopec E, Zwolska Z, 2007. Synthesis of isomeric, oxathiolone fused chalcones and comparison, their activity towards various microorganisms and human cancer cell lines. *Chemical and Pharmaceutical Bulletin*, 55: 817-820.
- [6] Oyedapo AO, Mankanju VO, Adewunmi CO, Iwalewa EO, Adenowo TK, 2004. Antitrichomonal activity of 1,3-diaryl-2-propen-1-one on *Trichomona gallinae*. *African Journal of Traditional, Complementary and Alternative Medicines*, 1: 55-62.
- [7] Jin F, Jin YL, Sohn DW, Kim S-A, Sohn DH, Kim YC, Kim HS, 2007. Structural requirements of 2',4',6'-Tris (methoxymethoxy) chalcones derivatives for anti-inflammatory activity. *Archives of Pharmacal Research*, 30: 1359-1367.
- [8] Narender T, Khaliq T, Shewata, Nishi, Goyal N, Gupta S, 2005. Synthesis of chromenochalcones and evaluation of their in vitro antileishmanial activity. *Bioorganic and Medicinal Chemistry*, 13: 6543-6550.
- [9] Aponte JC, Verastegui M, Malaga E, Zimic M, Quiliano M, Vaisberg AJ, Gilman RH, Hammond GB, 2008. Synthesis, cytotoxicity and anti-*Trypanosoma cruzi* activity of new chalcones. *Journal of Medicinal Chemistry*, 51:6230-6234.
- [10] Prasad YR, Rao AL, Rambabu R, Ravikumar P, 2007. Synthesis and biological evaluation of some novel chalcone derivatives. *Oriental Journal of Chemistry*, 23: 927-937.
- [11] Rao MS, Kotes J, Narukulla R, Duddeck H, 2004. Synthesis and spectroscopic characterization of some chromanochalcones and their dihydro derivatives. *Arkivoc*, XIV: 96-102.
- [12] Calvino V, Picallo M, Lopez-peinado AJ, Martin-Aranda RM, Duran-valle CJ, 2006. Ultrasound accelerated Claisen-Schmidt condensation: A green route to chalcones. *Applied Surface Science*, 252: 6071-6074.
- [13] Eddarir S, Catelle N, Bakkour Y, Ranlando C, 2003. An efficient synthesis of chalcones based on the Suzuki reaction. *Tetrahedron Letters*, 44: 5359-5363.
- [14] Petrov O, Ivanova Y, Gerova M, 2008. $\text{SOCl}_2/\text{EtOH}$: Catalytic system for synthesis of chalcones. *Catalysis Communication*, 9: 315-317.
- [15] Sebti S, Solhy A, Somabh A, Kossir A, Oamimoun H, 2002. Dramatic activity enhancement of natural phosphate catalyst by lithium nitrate: An efficient synthesis of chalcones. *Catalysis Communication*, 3: 335-339.
- [16] Macquarrie DJ, Nazih R, Sebti S, 2002. KF/natural phosphate as an efficient catalyst for synthesis of 2'-hydroxychalcones and flavanones. *Green Chemistry*, 4: 56-59.
- [17] Dong F, Jian C, Zhenghao F, Kai G, Zuliang L, 2008. Synthesis of chalcones via Claisen-Schmidt condensation reaction catalyzed by acyclic acidic ionic liquids. *Catalysis Communication*, 9: 1924-1927.
- [18] Perozo-Rondon E, Martin Aranda RM, Casal B, Duran-Valle CJ, Lau WN, Zhang XF, Yeung KL, 2006. Sonocatalysis in solvent free conditions: An efficient eco-friendly methodology to prepare chalcones using a new type of amino grafted zeolites. *Catalysis Today*, 114: 183-187.

- [19] Craig M, Comisor P, Sarage E, 2004. Kinetics of crossed aldol condensations in high-temperature water. *Green Chemistry*, 6: 227-231.
- [20] Zhang Z, Dong YW, Guan WU, Wang GW, 2003. Efficient and clean aldol condensation catalyzed by sodium carbonate in water. *Chemistry Letters*, 32: 966-967.
- [21] Tanemura K, Suzuki T, Nishida Y, Horaguchi T, 2005. Aldol condensation in water using polyethylene glycol 400. *Chemistry Letters*, 34: 576-577.
- [22] Thirunarayanan G, Vanangamudi G, 2006. Synthesis of some 4-bromo-1-naphthyl chalcones using silica-sulfuric acid reagent under solvent free conditions. *Arkivoc*, XII: 58-64.
- [23] Sarda SR, Jadhav WN, Wasmakar SK, Dake SA, Pawar RP, 2009. Solvent-free NaOH-Al₂O₃ supported synthesis of 1,3-diaryl-2-propene-1-ones. *International Journal of Chem Tech Research*, 2: 265-269.
- [24] Hazarkhani H, Kumar P, Kondiram KS, Shafi Gadwal IM, 2010. Highly selective Claisen-Schmidt condensation catalyzed by silica chloride under solvent-free reaction conditions. *Synthetic Communications*, 40: 2887-2896.
- [25] Rateb, Nora M, Zohdi, Hussein F, 2009. Atom-efficient, solvent-free, green synthesis of chalcones by grinding. *Synthetic Communications*, 39: 2789-2794.
- [26] Toda F, 1995. Solid state organic chemistry: efficient reactions, remarkable yields, and stereoselectivity. *Account of Chemical Research*, 28: 480-486.
- [27] Rothenberg G, Dowine AP, Raston CL, Scott JL, 2001. Understanding solid/solid organic reactions. *Journal of American Chemical Society*, 123: 8701-8708.
- [28] Vibhute AY, Mokle SS, Nalwar YS, Vibhute YB, Gurav VM, 2009. An efficient and operationally simple synthesis of some new Schiff Bases using grinding technique. *Bulletin of Catalysis Society of India*, 8: 164-168.
- [29] Vibhute A, Mokle S, Karamunge K, Gurav V, Vibhute Y, 2010. A simple and efficient method for solvent-free iodination of hydroxylated aromatic aldehydes and ketones using iodine and iodic acid by grinding method. *Chinese Chemical Letters*, 21: 914-918.