

Anthelmintic Resistance to Commonly Used Drugs in Naturally Infected Sheep in and Around Bekoji

Diriba Lemma*, Birhanu Abera and Eyob Eticha

Asella Regional Veterinary Laboratory, Asella, Ethiopia

Abstract

The study aimed at assessing parasite control practices in use and to determine existence of gastro intestinal nematodes resistance for Albendazole, Tetraclozash and Ivermectin, in naturally infected sheep under field conditions. In Lemu Bilbilo district, sheep of both sexes randomly select from small holder farmers from different Pa's were used for the study. Before the actual experiment, screening was done to identify sheep naturally infected with Gastro intestinal nematodes and Eggs per Gram (EPG) greater than 200 were eligible for inclusion in the field experiment following guidelines by Coles, et al. Accordingly a total of 48 animals grouped into four treatment groups Albendazole, Tetraclozash, Ivermectin and control (left untreated). Albendazole, Tetramisole and Ivermectin were found 90.53%, 98.1% and 96.59% respectively.

The percentage reduction of faecal egg count (95% confidence intervals) for Albendazole, Tetraclozash and Ivermectin were, 95.83%, (95.1-84.4) 98.1% (99-90.99) and 96.59% (97.01-91.3) respectively. Resistance is suspected on albendazole based on the findings, it was concluded that development of anthelmintic resistance could be prevented by avoid frequent and unnecessary treatments anthelmintics, opting instead for strategic deworming.

Keywords

Anthelmintic • Resistance • Nematodes • Helment

Introduction

The control of parasitic helment in domestic animals relies largely on the use of few antihelminthic clases of drugs as a consequence antihelminthic resistance becoming a world-wide problem for both small and large scale farmers, sheep and goats are mainly found in arid and semi-arid areas of sub Saharan Africa. They play a vital role in rural economies through the provision of meat, milk, household income, manure and skin. In sub Saharan Small ruminants have also been reported to survive better under drought conditions than cattle due to their low body mass and low metabolic requirements [1]. Ethiopia lies within the tropical latitudes of Africa, and has an extremely diverse topography, a wide range of climatic features and a multitude of agro-ecological zones, which makes the country suitable for different agricultural production systems. This in turn has contributed to the existence of a large diversity of farm animal genetic resources in the country. Ethiopian livestock production systems are broadly characterized as low input, mixed crop-livestock, agro-pastoral and pastoral systems; as well as medium input, peri-urban and urban enterprises. In Ethiopia, parasitological investigations of small ruminants in the humid central highland regions of the country have demonstrated that nematodes of the genera *Haemonchus*, *Trichostrongylus*, *Oesophagostomum*, *Bunostomum*, *Strongyloides*, *Cooperia*, *Nematodirus* and *Trichuris* are the most common [2,3]. The liver flukes of the genus *Fasciola* (*F. hepatica* and *F. gigantica*) are also of particular importance in ruminant livestock found in these regions [4,5]. Another study by Debela also reported that *H. contortus*, *Strongyloides papillosus* and *Trichostrongylus* spp. were the most prevalent Gastro Intestinal (GI) nematode species infecting goats in the Rift Valley of southern Ethiopia [6]. Small-holder farmers and pastoralists of Ethiopia practice varying degrees of parasite control in their livestock. These practices range from the use of anthelmintic drugs of varying quality, to the use of traditional

medicines. Livestock diseases are one of the major production constraints frequently observed in the oromia region among which helminth parasites are the biggest causes of production losses [7]. However, the indiscriminate use of these drugs without appropriate association with other methods to fight parasite infection has favored the development of helminth resistance to various drugs in the gastrointestinal tract of small ruminants.

Effective management of GI parasites in grazing livestock relies on the strategic use of highly efficacious chemotherapy [8]. However, decades of indiscriminate use has resulted in widespread development of Anthelmintic Resistance (AR) including multi-drug resistance in various GI nematode populations [9]. AR is the situation where normal dosage of a drug does not produce a consistent reduction in worms or excreted eggs failure to identify AR and manage its development will incur severe production penalties due to the impact of parasitic gastroenteritis [8,10]. Considering the narrow range of available drugs and slow rate of new drug development, AR presents an alarming global threat demanding vigilant monitoring and management.

In Ethiopia, various anthelmintic have been used in different parts of the country for the treatment of sheep and goats helminth parasites [11,12]. The use of anthelmintic has been practiced for a long time, and constitutes a considerable share of the costs spent by the country in the control of helminthosis [11]. Also, smuggling and misuse of veterinary drugs involving anthelmintics is a wide spread practice in the country [13]. Some of these drugs, particularly albendazole and tetramisole, have been continuously imported and distributed to every corner of the country under different trade names and by different manufacturers [14-18].

Anthelmintic resistance in nematodes of small ruminants has been in all continents and is now recognized as one of the greatest threats to grazing livestock production. Originally the situation was confusing in small ruminants because the poor efficacy of anthelmintics could be due to true resistance or to inadequate dosage of drug. Therefore, the objective of this study was to investigate the existence of GIN resistance for albendazole, tetra-misole and ivermectin, in naturally infected sheep under field conditions in Lemu Bilbilo district [19-25].

Materials and Methods

Study area

The study was conducted from September, 2017 to June 2018 At Lemu Bilbilo district of Arsi zone, south eastern part of Ethiopia which is located at 231 kms. From Addis Ababa and latitudes of about 039016.340N and longitudes of about 07°33.131E as well as elevation ranging from 2755-2878 meters above sea levels. The area receives the annual mean minimum and maximum rain fall of about 800 and 1400 mms respectively,

*Address for Correspondence: Diriba Lemma, Asella Regional Veterinary Laboratory, Asella, Ethiopia; Tel: 0910172216; Email: diribalema@gmail.com

Copyright: © 2021 Lemma D, et al. This is an open-access article distributed under the terms of the creative commons attribution license which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Received: July 08, 2021; Accepted: July 22, 2021; Published: July 29, 2021

which comes from the long and short rainy seasons. The minimum and maximum temperature is 18°C and 28°C respectively. The small ruminant populations in this area were 291,285 sheep and 28, 864 goats (Figure 1).



Figure 1. Map of study area.

Study animals and study design

Sheep of both sexes randomly select from small holder farmers from different Pa's were used for the study. The suckling lambs, pregnant females, very old sheep (>5 years) and animals treated with anthelmintic or traditional medicine were excluded from study. In addition, each sheep selected were individually identified with a numbered ear tagged. Before the actual experiment, screening was done to identify sheep naturally infected with Gastro intestinal nematodes and Eggs per Gram (EPG) greater than 200 were eligible for inclusion in the field experiment following guidelines by Coles, et al. accordingly a total of 48 animals grouped into four treatment groups: albendazole, tetramisole, ivermectin and control (i.e. left untreated). On day 0, fecal samples were collected from each animal enrolled in the study, and then the animals were either treated with an anthelmintic or left untreated. Dose rate and route of administration were based on manufactures' recommendation and anthelmintics were chosen based on the frequency of utilization in the study area described in Table 1. Fecal samples were collected again 10 to 14 days post-treatment from all animals included in the study, and the changes in the EPG were determined (Table 1).

Trade name	Generic name	Manufacture	Dose	Route
Albenda-QK 300 MG	Albendazole	Chengdu qiankun vet. pharm. co. ltd., China	7.5 mg per kg	oral
Ivermectin 1%	Ivermectin	Chengdu qiankun vet. pharm. co. ltd., China	0.2 mg per kg	Subcutaneous
Tetraclozash-900	Tetramisole+ Oxyclozanide	Ashish life science pvt, India	7.5 mg per kg	oral

Table 1. Drugs used in the trial.

Data analysis

The effectiveness of different anthelmintics was evaluated by calculating FECRT% by applying the formula $E = [1 - (T/C)] * 100$, where E=percentage efficacy; C=mean number of eggs in the control group; T=mean number of eggs in the treated group; where the value of FECRT % less than 95% was suspected as developed resistance or poor efficacy described by Coles, et al.

Results

In the present study out of 162 examined 72 (44.4%) were found to be

positive for the gastrointestinal nematode eggs. The major GI nematodes observed in ruminants at the study area were Strongyle type, Strongyloides and Trichuris with the prevalence of 22%, 9% and 1%, respectively. In this mixed nematode eggs were also examined with prevalence of 12.4%. The mean pre and post treatment faecal egg counts (EPG) and the percentage of Faecal Egg Count Reduction (FECR) and The percentage reduction of ivermectin were found 90.53%, 98.1% and 96.59 % respectively (Tables 2 and 3).

Nematode egg type	No examined	No of positive (prevalence)
Strongyle	162	36(22.2%)
Strongyloides	162	15(9.3%)
Trichuris	162	2(1.2%)
Mixed type	162	20(12.4%)
Total	162	73(45%)

Table 2. Proportion of GI nematode species in ruminants at the study area.

Drug	No of animals used in each group	Pre-treatment mean ± SEM	Post-treatment mean ± SEM	Reduction (%)	95%CL (UCL-LCL)
Albendazole	n=12	550.00 ± 112.983	104.17 ± 39.147	90.53	95.1-84.4
Ivermectin	n=12	1212.50 ± 207.814	37.50 ± 23.131	96.59%	97.01-91.3
Tetraclozash	n=12	1395.83 ± 339.700	20.83 ± 8.65	98.10%	99-90.99
Control	n=12	975.00 ± 297.114	1100 ± 238.127	NA	NA

Table 3. The faecal egg count reduction percentage following treatment with albendazole, tetraclozash and ivermectin, in sheep.

Discussion

The presence of anthelmintic resistance on the investigated area was initially declared by considering the percentage reduction in faecal egg counts was less than 95% and the lower limit of the 95% confidence interval was less than 90%. If only one of these criteria is met, anthelmintic resistance is suspected. Based on this criterion, the FECR percentage and the lower confidence limit obtained from lemu Bilbilo district smallholder sheep production system revealed the absence of a significant level of GIN resistance to tetramisole and ivermectin. This finding is in line with other studies conducted in various parts of Ethiopia [12,14-16]. These studies also reported the absence of anthelmintic resistance to albendazole, tetramisole and ivermectin in small ruminants in Ogaden. This finding revealed that albendazole was suspected for resistance in the area.

Conclusions and Recommendations

The current findings indicated that albendazole was suspected for development of resistance, while ivermectin and tetraclozash were found to be not resistance.

To prevent further development of anthelmintic resistance in this area, the following practices are recommended:

- Avoid frequent and unnecessary treatments anthelmintics, opting instead for strategic deworming.

- Avoid under dosing of animals.
- Further studies, are needed to determine the anthelmintic resistance status of the different species of GINs.
- A targeted selective treatment traditionally practiced by farmers should be promoted and supported by coprological examination to identify animals in need of treatments.
- Anthelmintic efficacy tests should also be part of the worm management programs to monitor treatment failures.

References

1. Wesongah, LJO, Chemulitti FD, Wesonga L and Munga P, et al. "Trypanosomosis and Other Parasitic Diseases Affecting Sheep and Goats Production in Two Group Ranches, Narok District, Kenya." (2003).
2. Bekele, T, Woldeab T, Kassi AL and Sherington J. "Factors Affecting Morbidity and Mortality On-farm and On-station in the Ethiopian Highland Sheep." *Acta Trop* 52(1992): 99-109.
3. Tembely, S, Kassi LA, Rege JE and Sovani S, et al. "The Epidemiology of Nematode Infections in Sheep in a Cool Tropical Environment." *Vet Parasitol* 70(1997): 129-141.
4. Ngategize, PK, Bekele T and Tilahun G. "Financial Losses Caused by Ovine Fascioliasis in the Ethiopian Highlands." *Trop Anim Health Prod* 25(1993): 155-161.
5. Mezgebu, M. "A Survey on Ovine Fascioliasis and Lungworm Infection in Addis Ababa and the Surrounding Highland Areas." DVM Thesis, Debre Zeit, Ethiopia, (1995).
6. Debela, E. "Epidemiology of Gastro-intestinal Helminthiasis of Rift Valley Goats Under Traditional Husbandry System in Adami Tulu District, Ethiopia." *Ethiop J Sci* 25(2002): 35-44.
7. Miller, JE and Horohov DW. "Immunological Aspects of Nematode Parasite Control in Sheep." *J Anim Sci* 84(2006): 124-132.
8. Good, B, Hanrahan JP, Waal DT and Patten T, et al. "Anthelmintic-resistant Nematodes in Irish Commercial Sheep Flocks- The State of Play." *Ir Vet J* 65(2012): 21.
9. Kaplan, RM. "Drug Resistance in Nematodes of Veterinary Importance: A Status Report." *Trends Parasitol* 20(2004): 477-481.
10. Silvestre, A, Leignel V, Berrag B and Gasnier N, et al. "Sheep and Goat Nematode Resistance to Anthelmintics: Pro and Cons among Breeding Management Factors." *Vet Res* 33(2002): 465-480.
11. Biffa, D, Jobre Y and Chaka H (2006). "Ovine Helminthosis, A Major Health Constraint to Productivity of Sheep in Ethiopia." *Anim Health Res Rev* 7(2006): 10-118.
12. Asmare, K, Gelaye E and Ayelet G. "Anthelmintic Resistance Test in Gastrointestinal of Small Ruminants in Southern Ethiopia." *Bullet Anim Health Prod Afr* 53(2005): 89-95.
13. Maingi, N, Bjorn H, Thamsborg SM and Bogh HO. "A Survey of Anthelmintic Resistance in Nematode Parasites of Goats in Denmark." *Vet Parasitol* 66(1996): 53-66.
14. Kumsa, B, and Wosene A. "Efficacy of Albendazole and Tetramisole against *Haemonchus contortus* in Experimentally Infected Lambs." *Intern J Appl Res Vet Med* 4(2006): 94-98.
15. Sheferaw, D, and Asha A. "Efficacy of Selected Anthelmintics Against Gastrointestinal Nematodes of Sheep Owned by Smallholder Farmers in Wolaita, Southern Ethiopia." *Ethiop Vet J* 14(2010): 31-38.
16. Tadesse, A, Asmare K, Bekele J and Abebe G. "Study on Anthelmintic Resistance Nematodes at Goat Farm of Hawassa College Agriculture Southern Ethiopia." *Ethiop Vet J* 13(2009): 49-57.
17. Waller, PJ, Baldock C, Rolfe P and Riffkin G, et al. "Anthelmintic Resistance: Report of Working Party for the Animal Health Committee of the Standing Committee on Agriculture (SCA)." SCA Technical Report Series. Canberra, Australia. (1989): 26.
18. Waller, PJ. "Anthelmintic Resistance." *Vet Parasitol* 72(1997): 391-405.
19. Demeke, N and Zewdu S. "A Review on Anthelmintic Resistance and Potential Risk Factors in Domestic Ruminants College of Veterinary Medicine, University of Gondar, Gondar, Ethiopia." *Acta Parasitologica Globalis* 8(2007): 58-67.
20. Fikru, R, Lelisa K, Gezhagn M and Bersisa K, et al. "Efficacy of Commonly Used Anthelmintic Drugs in Naturally Infected Sheep and Goats in Central Oromia, Ethiopia." *Res J Pharmacol* 7(2013): 48-53.
21. Jemal, A, Ararsa D, Dareje R and Dinaol B, et al. "Gastrointestinal Nematode Parasites of Small Ruminants and Anthelmintics Efficacy Test in Sheep of Haramaya District, Eastern Ethiopia." *Anim Vet Sci* 5(2017): 39-44.
22. Klauk, V, Pazinato R, Lopes LS and Cucco DC, et al. "*Trichostrongylus* and *Haemonchus* Anthelmintic Resistance in Naturally Infected Sheep from Southern Brazil." *An Acad Bras Cienc* 86(2014): 777-784.
23. Sissay, MM. "Helminth Parasites of Sheep and Goats in Eastern Ethiopia Epidemiology, and Anthelmintic Resistance and its Management." (2007).
24. Tembely, S. "Development and Survival of Infective Larvae of Nematode Parasites of Sheep on Pasture in a Cool Tropical Environment." *Vet Parasitol* 79(1998): 81-87.
25. Yitayew D, Zewdu S, Kalkidan G and Dagmawi Y. "Anthelmintics Resistance against Gastrointestinal Nematodes of Sheep: A Review." *World J Agricul Sci* 12(2016): 245-253.

How to cite this article: Lemma, Diriba, Birhanu Abera and Eyob Eticha. "Anthelmintic Resistance to Commonly Used Drugs in Naturally Infected Sheep in and Around Bekoji." *J Vet Sci Techno* 12 (2021): 99.