

Association Between Hyperuricemia and Metabolic Syndrome in a Senegalese Population

Fatou Cisse*

Department of Odontostomatology, Cheikh Anta Diop University of Dakar, Senegal

Abstract

Background: Metabolic Syndrome (MS) is a real public health problem in our regions. Its severity requires early detection strategies based on clinical, anthropometric and biological criteria to ensure adequate management of this condition. Hyperuricemia, although not part of these criteria, is frequently associated with MS components. Thus, we have set ourselves the goal of finding a relationship between uric acid and MS.

Methodology: We conducted a prospective study on 441 subjects received as part of an annual medical visit. The subjects included in the study benefited from a complete clinical examination and blood tests provided by this annual medical check (including blood sugar, lipid status, uric acid). The biochemical parameters were determined using enzymatic methods adapted to the A15 automaton. The metabolic syndrome has been defined according to the criteria of NCEP-ATP III

Results: The prevalences of metabolic syndrome and hyperuricemia were 6.57% and 6.34%, respectively. Hyperuricemia was strongly associated with MS (OR = 3.87, $p = 0.007$) and some of these components including hypoHDLemia (OR = 5.09), hypertriglyceridemia (OR = 3.05), and abdominal obesity (OR = 2.52)

Conclusion: The positive association between hyperuricemia and MS demonstrates the interest of dosing uric acid in subjects at risk

Keywords: Hyperuricemia • Metabolic syndrome • Senegal

Introduction

Uric acid is the end product of purine metabolism in humans. Serum uric acid concentrations will therefore depend on the balance between intakes (endogenous and exogenous) and excretions. However, in a large number of cases, hyperuricemia results from a lack of renal elimination of uric acid associated or not with excessive food intake. Many studies have shown a relationship between high blood levels of uric acid and hypertension, obesity and hypertriglyceridemia, which are components of the MS. This syndrome associated with a risk of diabetes and cardiovascular disease, is one of the most important public health problems in the world with a growing prevalence. The severity of this syndrome justifies early detection strategies based on the search for simple clinical, anthropometric and biological arguments in order to guarantee early management of this condition. Hyperuricemia, although not part of the criteria for defining the metabolic syndrome, is most often associated with its components; this is why, in this

work, we set ourselves the objective of seeking a relationship between uric acid and SM.

Materials and Methods

This is a prospective study on 441 subjects received as part of a systematic annual visit to a Senegalese company involving administrative, technical and service staff; Included in the study were all subjects who had undergone a full clinical examination and blood tests provided for by this annual medical check-up (including blood sugar, lipid balance, uric acid).

Anthropometric data were collected on survey cards designed for this purpose. The biochemical parameters were measured using enzymatic methods adapted on the A15 analyser (Biosystems).

The SM was defined according to the criteria of (NCEP-ATP III): abdominal obesity (waist > 102 cm (men) and > 88 cm (women), high blood pressure (BP $\geq 130/85$ mm Hg), hypertriglyceridemia (≥ 1.50

*Address to correspondence: Dr Fatou Cisse, Department of Odontostomatology, Cheikh Anta Diop University of Dakar, Senegal, Tel: 221776123465; E-mail: kinciss@yahoo.fr

Copyright: © 2021 Cisse F. This is an open-access article distributed under the terms of the creative commons attribution license which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Received: 26 August, 2021; **Accepted:** 09 September, 2021; **Published:** 16 September, 2021.

g/l), lower cholesterol HDL (<0.40 g/l (men) or <0.50 g/l (women)) and hyperglycemia (≥ 1.10 g/l); Subjects with at least three of these criteria were considered to have MS. Hyperuricemia has been defined as serum uric acid concentrations ≥ 72 mg/l in men and ≥ 60 mg/l in women.

All observations were entered and coded on Windows Excel 2010 (Microsoft, USA) and then analyzed using SPSS Statistics 24 (IBM, Chicago, IL, USA). Excel software was also used to present the tables, as well as to group the modalities of certain variables before their analysis.

Descriptive statistics presented the data in the form of tables. It then summarized the qualitative variables in the form of proportions (percentages) and the quantitative variables in the form of means +/-Standard Deviation.

For the qualitative variables, the comparison of the percentages required the Pearson chi-square test with Yates correction if necessary for small samples.

The statistical inference necessary for the analytical study was used to study the associations using statistical tests for the comparison of variables (Logistic regression) and an approximation of the relative risk using the Odds Ratio (OR) with a confidence interval 95% (95% CI). A risk value of the first kind $\alpha = 0.05$ was considered as a significant threshold.

Results and Discussion

The average age of our study population was 38.82 ± 10.35 with a sex-ratio M/F of 1.12. The prevalence of MS and hyperuricemia were 6.57% and 6.34% respectively (Table 1). The prevalence of hyperuricemia was 41.38% in subjects with MS versus 23.35% in subjects without MS.

Variables	Results
Number of individuals	441
Age (years) Mean \pm SD	38,82 \pm 10,35
Gender-ratio M/F	1,12
Men N (%)	233 (52,83%)
Women N (%)	208 (47,18%)
Metabolic Syndrom N (%)	29 (6,57%)

Table 1. Characteristics of the study population.

Hyperuricemia and MS were strongly associated (OR = 3.87, $p = 0.007$). The study of the relationship between hyperuricemia and the other components of MS has shown a strong association with hypoHDLemia (OR = 5.09), hypertriglyceridemia (OR = 3.05) and abdominal obesity (OR = 2.52). We also found a higher prevalence of hyperuricemia in subjects over 50 years with a male predominance.

Our study found a positive association between hyperuricemia and SM (OR = 2.29, $p = 0.05$). These results corroborate those of numerous epidemiological data and experimental. The relationship between uric acid levels and the risk of developing MS has been established by a meta-analysis of prospective studies. Indeed, hyperuricemia is a risk factor for MS. This could be explained by its

constant association with the different components of the SM. In fact, in our study, significant associations were found between hyperuricemia and hypertension, abdominal obesity and dyslipidemia (Table 2). This is in agreement with the data of the literature. However, we have not found an association between hyperuricemia and hyperglycemia.

Components	Hyperuricemia		OR	P
	Yes	No		
Abdominal obesity				
Yes	10	92	2,52	0,04
No	14	325		
HTA				
Yes	15	171	2,14	0,09
No	10	245		
Hypertiglyceridemia				
Yes	6	41	3,05	0,04
No	18	376		
hypoHDLemia				
Yes	7	28	5,09	< 0,001
No	19	387		
Hyperglycemia				
Yes	4	29	2,54	0,2
No	21	387		

Table 2. Association between hyperuricemia and the various components of MS.

These results could be linked to the inhibitory action of glucose on the reabsorption of uric acid which thus promotes its excretion. This role of glucose cannot explain the high prevalence of hyperuricemia in type 2 diabetics found by some authors. According to the latter, hyperuricemia in diabetics was favored by the existence of an MS or one of its components.

The study of the distribution of hyperuricemia according to age and sex shows on the one hand an increase in the prevalence in subjects over 50 years and on the other hand higher rates in men compared women. The opposite being observed for the SM. These opposite trends could be linked to the biases introduced by the differences in population size in the subgroups. Increased uric acid with age has been reported by Woyesa. However, Fu and colleagues found no significant association between uric acid and age. As for the male predominance, it has been widely described in the literature.

Conclusion

A positive association has been found between hyperuricemia and SM in the Senegalese population. This demonstrates the value of measuring uric acid in subjects at risk. Further work will be required

to demonstrate whether hyperuricemia is a cause or a consequence of MS.

Conflicts of Interest

The authors declare that they have no conflict of interest

Author's contribution

Fatou Cisse developed the study protocol and the data collection. Fatou Cisse, Tarik Lamkinsi and Souleymane Thiam contributed to the analysis of the results and the drafting of the final document. The methodology and the statistical analyzes were performed by Tarik Lamkinsi. All authors have read and approved the final version of the manuscript.

References

1. Avula, Naveen Reddy, Shenoy D. "Evaluation of Association of Hyperuricaemia with Metabolic Syndrome and Insulin Resistance." *J Clinical Diag Res* 10 (2016): 32-34.
2. Cameron, Adrian J, Magliano DJ, Zimmet PZ, and Welborn TA, et al. "The Metabolic Syndrome as a Tool for Predicting Future Diabetes: The AusDiab Study." *J Internal Med* 264 (2008): 177-186.
3. Carbone, Federico, Montecucco F, Mach F, And Pontremoli R, Et Al. "The Liver and the Kidney: Two Critical Organs Influencing the Atherothrombotic Risk in Metabolic Syndrome." *Thromb Haemost* 110 (2013): 940-958.
4. Cibickova, L, Langova K, Vaverkova H, and Kubickova V, et al. "Correlation of Uric Acid Levels and Parameters of Metabolic Syndrome." *Physiol Res* 66 (2017): 481.
5. Conen, David, Wietlisbach V, Bovet P, and Shamlaye C, et al. "Prevalence of Hyperuricemia and Relation of Serum Uric Acid with Cardiovascular Risk Factors in a Developing Country." *BMC Public Health* 4 (2004): 1-9.

How to cite this article: Cisse Fatou. "Association Between Hyperuricemia and Metabolic Syndrome in a Senegalese Population." *J Cosmo Trichol* 7 (2021) : 26876