

## Bilateral Complex Traumatic Cleft of the Ear Lobes

Richard E Nnabuko\*

Department of Plastic Surgery, National Orthopaedic Hospital, Enugu Nigeria

Bilateral complex traumatic cleft of the earlobes in a 54 year old lady. Aetiology was from wearing of heavy ear rings. Duration was for 10 years and had progressively worsened till the situation seen in the pre-operative photographs occurred.

Had reconstruction with modified L-plasty and complete closure of the earring hole. Fresh piercings done on non-scarred parts when previous scars settled.



\*Corresponding author: Richard E Nnabuko, Department of Plastic Surgery, National Orthopaedic Hospital, Enugu Nigeria, E-mail: [richnnabuko@yahoo.com](mailto:richnnabuko@yahoo.com)

Received November 13, 2014; Accepted November 17, 2014; Published November 19, 2014

Citation: Nnabuko RE (2014) Bilateral Complex Traumatic Cleft of the Ear Lobes. J Trauma Treat 4: 1102. doi:[10.4172/2167-1222.10001102](https://doi.org/10.4172/2167-1222.10001102)

Copyright: © 2014 Nnabuko RE. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.