

Body Mass Index in Children and Their Parents: A Cross-Sectional Study in a Study Population of Children from Southern Italy

Nicola Fuiano¹, Antonella Lonero², Giuliana Diddi², Vincenza Luce², Fabrizia De Palma², Maria Felicia Faienza² and Maurizio Delvecchio^{2*}

¹Pediatric Allergy Service, ASL FG, Torremaggiore (FG), Italy

²"B Trambusti" Unit, Department of Biomedical Sciences and Human Oncology, AOU Policlinico di Bari, Bari, Italy;

*Corresponding author: Maurizio Delvecchio, MD, PhD, "B Trambusti" Paediatrics Unit, A.O.U. Consorziale Policlinico, 70100 Bari, Italy. Tel: +393393379630; Fax: +390805592287; E-mail: mdelvecchio75@gmail.com

Received date: November 22, 2014; Accepted date: December 29, 2014 Published date: December 31, 2014

Copyright: © 2014 Fuiano N, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

The prevalence of overweight and obesity has increased over the last decades. Parental obesity plays an important role in determining childhood obesity. We aimed to evaluate the relationship between parental and offspring weight status in a population of children from South of Italy, as no data have ever been published from this area. We recruited 636 children (5.7 ± 1.5 years old) and their parents. Seventy-three (11.5%) and sixteen (2.5%) children were overweight and obese, respectively. Offspring weight status was significantly associated with parents' weight status. The linear regression analysis showed that offspring BMI was more affected by paternal than by maternal BMI. Our data confirmed that parents' weight status plays an important role on children's BMI. Interestingly, in our study, parents' height and weight were measured and not reported as in most of the previous papers, strengthening our conclusions. We suggest that intensive nutritional education and preventive programs should be performed in children with overweight / obese parents rather than in children with normal weight parent. Furthermore, nutritional education should be performed also for overweight parents to modify preventable risk factor for pediatric obesity.

Keywords: Obesity; Childhood; Parents; BMI

Introduction

The worldwide prevalence of overweight and obesity has dramatically increased among young people during the last decades. Furthermore, the relative raise over the last twenty years was higher in developing Countries (+65%) than in developed Countries (+48%) [1]. In Italy, the prevalence of overweight and obesity in children and adolescents aged from 6 to 17 years ranges from 13.7% to 23.6% and from 3.7% to 12.3% [2-5], respectively.

Obesity is associated with many health problems in childhood and represents a risk factor for obesity in adulthood [6]. It is associated with a proinflammatory and prothrombotic state [7], leading to cardiovascular dysfunction, type 2 diabetes mellitus, metabolic syndrome, snoring, asthma, early puberty, menstrual irregularities, steatohepatitis, pulmonary, renal, musculoskeletal, gastrointestinal, and psycho-social problems [6,8].

Once established, obesity is difficult to treat. Thus it is important to identify modifiable risk factors, contributing to the development of overweight and obesity, which have to be targeted by preventive programs [9]. Among the risk factors, parental obesity was identified as a predominant risk factor for childhood obesity, likely due to a combination of genetic, epigenetic, social, and environmental factors [10]. Consequently, the risk of childhood obesity is higher in obese parents' children [11-15]. Children with 2 obese parents are 10 to 12 times more likely to be obese [15-17]. The same-sex correlation between parents and child seems to be stronger than the opposite-sex correlations [18].

Cross-sectional studies have demonstrated that parental body mass index (BMI) is predictive of offspring BMI, in keeping with the evidence that parental lifestyle affects the children's behavior. Other studies reported that infant BMI is more strongly associated with maternal than paternal obesity in different age brackets [19,20]. Inverse association between socio-economic status and adiposity in children is well established [11,15], and the influence of the father seems to be stronger than that of the mother [14].

A limit of these studies is that parental height and weight were often reported and not measured, affecting the reliability of the results. Furthermore, no studies on population recruited in South of Italy have ever been performed. Prompted by this consideration, we focused our attention on the relationship between parental and offspring weight status in a population of preschool and school children recruited in a single Centre from South of Italy.

Subjects and Methods

The study population included 636 children (5.7 ± 1.5 years old, age range 2.9 – 9.0 years, 348 males - 54.7%;) and their biological parents. Children were recruited at school during a routine medical examination. Anthropometric measurements were performed by a single trained physician using standard procedures. Standing height by a wall-mounted Harpenden Stadiometer and weight in underwear by an electronic scale with digital readings accurate to 0.1 kg were measured. BMI was calculated dividing the weight in kilograms by the square of height in meters and expressed as Standard Deviation Score (SDS) [21]. Overweight and obesity were defined in according to the IOTF growth charts [22]. Adulthood obesity was defined as BMI ≥ 30 kg/m², overweight as BMI between 25 and 30 kg/m². The study was submitted and approved by the local ethic committee. Statistical

analysis was run by SPSS Software v.17. Statistical difference was considered significant when $p < 0.05$.

Results

Males and females were not different in terms of age at recruitment, BMI SDS, and rate of normal / underweight, overweight, and obesity

	Males	Females	Whole samples	Fathers	Mothers
Age (years)	5.7 ± 1.5	5.7 ± 1.6	5.7 ± 1.5	37.3 ± 6.2	33.8 ± 5.8
BMI (SDS)	0.15 ± 1.12	0.04 ± 1.06	0.1 ± 1.1	---	---
BMI (kg/m ²)	17.0 ± 2.8	16.8 ± 2.6	16.9 ± 2.7	25.7 ± 2.9	23.9 ± 3.8
Under / normalweight (n. patients, %)	293 (84.2%)	254 (88.2%)	547 (86%)	299 (47.0%)	437 (68.7%)
Overweight (n. patients, %)	43 (12.3%)	30 (10.4%)	73 (11.5%)	291 (45.8%)	145 (22.8%)
Obese (n. patients, %)	12 (3.7%)	4 (1.4%)	16 (2.5%)	46 (7.2%)	54 (8.5%)

Table 1: Features of offspring and their parents at recruitment. The values as expressed as mean ± standard deviation.

The chi-square analysis showed that offspring weight status was significantly associated with fathers ($p = 0.01$) and mothers ($p = 0.001$) weight status (Table 2). In males, there was no statistically significant

association with paternal or maternal weight status, while in females the rate of obesity was significantly higher in obese mothers' daughters ($p = 0.003$).

Offspring classification	Father classification			Mother classification		
	Normal / Underweight	Overweight	Obesity	Normal / Underweight	Overweight	Obesity
Normal / underweight (n. patients, %)	269 (49.2%)	242 (44.2%)	36 (6.6%)	391 (71.5%)	114 (20.8%)	42 (7.7%)
Overweight (n. patients, %)	22 (30.1%)	44 (60.3%)	7 (9.6%)	41 (56.2%)	22 (30.1%)	10 (13.7%)
Obesity (n. patients, %)	8 (50.0%)	5 (31.3%)	3 (18.8%)	5 (31.3%)	9 (56.3%)	2 (12.5%)

Table 2: Distribution of normal / underweight, overweight, and obesity in the offspring on the basis of mother's and father's weight status.

Offspring BMI was significantly correlated with both mothers ($r^2 = 0.157$ and $p < 0.001$) and fathers BMI ($r^2 = 0.169$ and $p < 0.001$). In males, BMI was correlated with mothers BMI ($r^2 = 0.153$ and $p = 0.004$) and fathers BMI ($r^2 = 0.144$ and $p = 0.007$). In females, BMI was correlated with mothers BMI ($r^2 = 0.176$ and $p = 0.003$) and fathers BMI ($r^2 = 0.191$ and $p = 0.001$). Both mothers and fathers BMIs were significantly different among normal / underweight, overweight and obese children (Figures 1a and 1b).

The linear regression analysis showed that offspring BMI was affected by both paternal and maternal BMI (offspring BMI = $-2.242 + 0.149 * \text{fathers BMI} + 0.135 * \text{mothers BMI}$, $R = 0.215$, $p = 0.001$). The same result was obtained both in males (offs= 0.001) than and in females (offspring BMI = $-2.726 + 0.171 * \text{fathers BMI} + 0.145 * \text{mothers BMI}$, $R = 0.245$, $p < 0.001$).

Discussion

Our data obtained in a quite large population of children from South of Italy confirmed that the parents weight status strongly affected children BMI [13,14]. Maternal obesity plays a significant role

mostly in determining daughters weight status. In the present study, the parents height and weight were measured and not reported as in the majority of the previous papers. This is an important point, because women typically under-report their weight while men typically over-report their height [18].

Even if a recent twin-study [23] concluded that genetic factors play the most important role in determining children weight status, available evidence suggests that obesity results from multiple interactions between genes, environmental and behavioral factors [24]. A lot of papers analyzed the relationship between parents and offspring weight status, showing that parents can indirectly influence unhealthy dietary (fatty foods, snacking, sweetened beverages), physical activity or sedentary behaviors (TV and computer addiction) of their children [10].

Besides these evidences, different data were reported about parents and offspring weight status at different age. Some data showed that early childhood growth was affected by both paternal and maternal BMI [25], with an independent and additive effect [26]. On the other hand, it was suggested that the maternal weight affected the severity of

childhood obesity at age 7, but both maternal and paternal weight affected offspring BMI at 15 years [27]. The mothers-offsprings correlations in BMI seemed to reach the maximum at 11 years, while the father-offsprings correlation at 16 years [28]. Similarly, Safer et al. showed that both of parents BMI affect offspring BMI, but the effect of mothers BMI appeared sooner [29].

Perez-Pastor et al. reported a close association between fathers and sons BMI, as well as between mothers and daughters BMI, indicating a social or behavioral influence and not a genetic inheritance of obesity. By the age of 8, there was a significant gender interaction with a risk of obesity 10-fold greater in girls and six-fold greater in boys [18]. In contrast, Linabery et al. and Heude et al. reported that infant BMI was more strongly associated with maternal than paternal obesity overall, with significant differences at birth and from 2-3 years [19] and from birth to puberty [20].

Our study confirmed the association between parental and offspring BMI, showing that both of parents BMIs plays a role by itself on offspring BMI. The regression analysis suggested that paternal BMI plays a major role than maternal BMI in determining offspring BMI, in contrast with some papers showing a larger effects exerted by maternal BMI [19,20,29] and other ones reporting a similar effects exerted by both of the parents [25,26].

Our data largely confirm the correlation between parental and offspring BMI described in literature, however some findings are brand new. Firstly, we show that father BMI affects offspring BMI more than the maternal BMI. Interestingly, beside the effect of parental BMI on offspring BMI, the maternal obesity is significantly associated with daughter BMI in our study population, a finding never reported before, helping us to identify a group of patients prone to develop obesity. Finally, children of our study population are young as compared to other papers, suggesting that the effect of parental obesity is exerted in young children.

The differences in age at recruitment of children and parents, social environment, and population size among all the studies likely accounted for the differences in the results. In conclusion, our paper confirms that offspring BMI is largely affected by parents BMI. Fathers BMI seems to play a major role on BMI, even if overweight / obese daughters were more frequently among overweight / obese mothers offspring. Obesity is a difficult condition to treat, involving behavior and eating attitude and nutritional education plays a key role in the prevention of obesity.

Overweight is a worldwide increasing social and medical issue, requiring great consideration not only from physicians. The identification of modifiable risk factors plays a key role in the prevention of pediatric obesity and our study shows that parental obesity is a risk factor for offspring obesity. Nutritional education should be an actual activity, provided by general practitioners and hopefully also at school setting. All children should be educated, but education nutritional programs should involve also their parents as parental BMI plays an important role in offspring weight status. In those families with overweight / obese parents, the prevention programs are recommended to be intensive, aiming to avoid an early onset of excessive weight gain. We recommend that in such families, also the parental obesity has to be targeted to facilitate weight loss in the children.




Figure 1a: Fathers BMI (kg/m²) on the basis of offspring weight status.




Figure 1b: Mothers BMI (kg/m²) on the basis of offspring weight status.

References

1. De Onis M, Blossner M, Borghi E (2010) Global prevalence and trends of overweight and obesity among preschool children. *Am J Clin Nutr* 92: 1257-1264.
2. Marras V, Macchis R, Foschini ML, Pilia S, Sortino M et al. (2006) Prevalence of overweight and obesity in primary school children in southern Sardinia, Italy. *Ital J Pediatr* 32: 251-55.
3. Celi F, Bini V, De Giorgi G, Molinari D, Faraoni F et al. (2003) Epidemiology of overweight and obesity among school children and adolescents in three provinces of central Italy, 1993-2001: study of potential influencing variables. *Eur J Clin Nutr* 57: 1045-51.
4. Maffei C, Consolaro A, Cavarzere P, Chini L, Banzato C et al. (2006) Prevalence of overweight and obesity in 2- to 6-year-old Italian children. *Obesity* 14 (5): 765-69.
5. Albertini A, Tripodi A, Fabbri A, Mattioli M, Cavrini G et al. (2008) Prevalence, determinants and indicators of risk of obesity in six-year-old children living in Central-North Italy. *Obes Rev* 9:4-10.
6. Reilly JJ, Methven E, McDowell ZC, Hacking B, Alexander D et al. (2003) Health consequences of obesity. *Arch Dis Child* 88: 748-752.
7. Giordano P, Del Vecchio GC, Cecinati V, Delvecchio M, Altomare M et al. (2011) Metabolic, inflammatory, endothelial and haemostatic markers in a group of Italian obese children and adolescents. *Eur J Pediatr* 170: 845-850.

8. Brunetti L, Tesse R, Miniello VL, Colella I, Delvecchio M et al. (2010) Sleep-Disordered breathing in obese children: the Southern Italy experience. *Chest* 137(5): 1085-90.
9. Danielzik S, Langnäse K, Mast M, Spethmann C, Müller MJ (2002) Impact of parental BMI on the manifestation of overweight 5-7 year old children. *Eur J Nutr* 41(3): 132-138.
10. Maffei C (2000) Aetiology of overweight and obesity in children and adolescent. *Eur J Pediatr* 159 (suppl 1): 35-44.
11. Van Stralen MM, v et al. (2012) Weight status of European preschool children and associations with family demographics and energy balance-related behaviours: a pooled analysis of six European studies. *Obes Rev* Mar 13 Suppl 1: 29-41.
12. Maffei C, Micciolo R, Must A, Zaffanello M, Pinelli L (1994) Parental and perinatal factors associated with childhood obesity in north-east Italy. *Int J Obes Relat Metab Disord* 18: 301-305.
13. Maffei C, Talamini G, Tato L (1998) Influence of diet, physical activity and parents' obesity on children's adiposity: a four-year longitudinal study. *Int J Obes Relat Metab Disord* 22: 758-764.
14. Lazzeri G, Pammolli A, Pilato V, Giacchi MV (2011) Relationship between 8/9-yr-old school children BMI, parents' BMI and educational level: a cross sectional survey. *Nutr J* 19: 10-76
15. Kleiser C, Schaffrath Rosario A, Mensink GB, Prinz-Langenohl R, Kurth BM (2009) Potential determinants of obesity among children and adolescents in Germany: results from the cross-sectional KiGGS Study. *BMC Public Health* Feb 2;9:46.
16. Whitaker KL, Jarvis MJ, Beeken RJ, Boniface D, Wardle J (2010) Comparing maternal and paternal intergenerational transmission of obesity risk in a large population-based sample. *Am J Clin Nutr* 91(6): 1560-1567.
17. Feummeler BF, Lovelady CA, Zucker NL, Østbye T (2013) Parental obesity moderates the relationship between childhood appetitive traits and weight. *Obesity (Silver Spring)* April 21(4): 815-823.
18. Perez-Pastor EM, Metcalf BS, Hosking J, Jeffery AN, Voss LD et al. (2009) Assortative weight gain in mother-daughter and father-son pairs: an emerging source of childhood obesity. Longitudinal study of trios (EarlyBird43). *Int J Obes* 33 (7): 727-735.
19. Linabery AM, Nahhas RW, Johnson W, Choh AC, Towne B et al. (2013) Stronger influence of maternal than paternal obesity on infant and early childhood body mass index: the Fels Longitudinal Study. *Pediatr Obes* Jun 8(3): 159-169.
20. Heude B, Kettaneh A, Rakotovo R, Bresson JL, Borys JM et al. (2005) Anthropometric relationships between parents and children throughout childhood: the Fleurbaix-Laventie Ville Santé Study. *Int J Obes* 29(10): 1222-1229.
21. Milani S, Balsamo A, Spada E, Bona G, et al. (2006) Italian cross-sectional growth charts for height, weight and BMI (2 to 20 yr). *J Endocrinol Invest* 29 (7): 581-593.
22. Cole TJ and Lobstein T (2012) Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatr Obes* 7(4): 284-294.
23. Wardle J, Carnell S, Haworth CMA, Plomin R (2008) Evidence for a strong genetic influence on childhood adiposity despite the force of the obesogenic environment. *Am J Clin Nutr* 87: 398-404.
24. Reilly JJ, Armstrong J, Dorosty AR, Emmett PM, Ness A et al. (2005) Early life risk factors for obesity in childhood: cohort study. *Br Med J* 330: 1357-1364.
25. Botton J, Heude B, Maccario J, Borys JM, Lommez A et al. (2010) Parental body size and early weight and height growth velocities in their offspring. *Early Hum Dev* 86 (7): 445-450.
26. Knight B, Shields BM, Hill A, Powerell RJ, Wright D et al. (2007) The impact of maternal glycemia and obesity on early postnatal growth in a non diabetic Caucasian population. *Diabetes Care* 30: 777-783.
27. Svensson V, Jacobsson JA, Fredriksson R, Danielsson P, Sobko T et al. (2011) Associations between severity of obesity in childhood and adolescence, obesity onset and parental BMI: a longitudinal cohort study. *Int J Obes (Lond)* 35(1): 46-52.
28. Lake JK, Power C, Cole TJ (1997) Child to adult body mass index in the 1958 British birth cohort: associations with parental obesity. *Arch Dis Child* 77: 376-381.
29. Safer DL, Agras WS, Bryson S, Hammer LD (2001) Early body mass index and other anthropometric relationships between parents and children. *Int J Obes Relat Metab Disord* 25: 1532-1536.