

Clinical Correlations between Endoscopic Findings and Histopathology

Narayan Murthy*

Department of Gastroenterology, University of Delhi, New Delhi, India

Introduction

Choosing appropriate study endpoints is a challenge faced in both animal models and patient-based research, particularly in the context of Inflammatory Bowel Disease (IBD). In animal models of IBD, the severity of the disease is often measured using semi-quantitative evaluation of intestinal histopathology, which is considered a reliable method (considered a "gold standard"). This evaluation involves using ordinal categorical scales, ranging from normal to severe, to assess various aspects such as immune cell infiltration and crypt loss. To obtain an overall severity score, different subscores measuring these histopathological features are combined. However, it is important to note that the relevance of specific histopathological features may vary between different animal models of IBD.

Epithelial destruction is highly relevant in the acute dextran sodium sulphate (DSS) model, whereas epithelial hyperplasia is more relevant in the T-cell transfer model. Concerns exist regarding the validity of disease severity indices currently used for the Interleukin-10 knock-out [IL10 KO] mouse model of chronic intestinal inflammation [1]. Several histopathological scoring systems have been described for this model, resembling those used for the T-cell transfer model. However, the final disease severity scores are simple combinations of item scores, lacking a methodologically rigorous scheme of index development. Consequently, there is uncertainty about the inclusion of relevant and reliable items in the final index.

Description

Only a minority of such patients will suffer clinically significant bleeding. Although mechanisms such as hypoperfusion leading to reperfusion injuries have been suggested the precise cause remains unclear. Mechanical ventilation for over 48 hours and coagulopathy are supposed to be the most grounded risk factors. Prior liver sickness, intense kidney injury requiring renal substitution treatment, furthermore, other organ brokenness may likewise increment risk. Prophylactic treatment of these lesions may not be without harm. Treatment may be a risk factor for ventilator-associated pneumonia. This can have a negative impact upon recovery resulting in prolonged critical care and hospital stays. Might be lost prompting incidental pneumonic taking care of. Patients might suction suddenly. This might be especially logical in patients who have created gastric balance as a result of basic disease. Parenteral sustenance requires focal venous access and line-related inconveniences like contamination, dying and pneumothorax is conceivable. At the cell level, glycogenolysis and gluconeogenesis drain liver and muscle glycogen stores quickly [2].

Intra-Endoscopy and histopathology assessments in mouse models for IBD are essential for both fundamental and translational research. Several research groups have developed ordinal classification indices, which to some extent measure overlapping features and generally have face validity [appear to be valid]. However, formal assessment of the operating properties of these indices had not been performed. In the current study, we evaluated histopathology and endoscopy scoring systems for reliability, validity and

***Address for Correspondence:** Narayan Murthy, Department of Gastroenterology, University of Delhi, New Delhi, India, E-mail: prof.murthy@edu.in

Copyright: © 2025 Murthy N. This is an open-access article distributed under the terms of the creative commons attribution license which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Received: 01 February, 2025, Manuscript No. cgi-25-161573; **Editor assigned:** 03 February, 2025, Pre QC No. P-161573; **Reviewed:** 15 February, 2025, QC No. Q-161573; **Revised:** 20 February, 2025, Manuscript No. R-161573; **Published:** 27 February, 2025, DOI: [10.37421/2952-8518.2025.10.296](https://doi.org/10.37421/2952-8518.2025.10.296)

ability to discriminate differences between groups known to vary in severity in mouse models for IBD, and developed simplified prediction models from these systems. We included two widely used mouse models for chronic intestinal inflammation: the T-cell transfer and the IL10 KO models. Items within the scoring systems were chosen based on the existing indices that are currently used for these mouse models. Usually an inter-rater ICC of >0.4 is used as criterion for including an item in the development of a new index. We showed that all histology items had a higher reliability than 0.4 [the lowest inter-rater ICC was 0.48, for 'ulceration']. Our observers agreed that this component was the most difficult item to score, especially on poorer quality slides. Indeed, removing the poor-quality slides from the dataset increased the inter-rater ICC of 'ulceration' to 0.53. In addition to 'ulceration', the model-building process excluded 'muscle thickness' and 'abscess', items for which the inter-rater ICCs were low. To enable a better distribution of disease severity scores, which enables better model building, the healthy control animals were excluded from the process of developing a new histological index. This process was started by investigating the bivariate relationships between the VAS score and the individual items. 'Inflammation' and 'crypt loss' exhibited non-linear relationships with the VAS and were re-categorized for the model-development process [3,4].

The MCHI was shown to be able to discriminate between control animals, untreated animals, and animals treated with a treatment of known efficacy. The new histological index has the potential to reduce the sample size required for detecting treatment effects, and as the readers do not have to score all initial features, to reduce the time required to read slides. The reliability coefficients of the histological components were considerably lower for the IL10 KO model than for the T-cell transfer model. As there was a lower number of slides available from the IL10 KO model for the model-development process, and the IL10 KO mice exhibited less severe disease, the MCHI may be a more efficient instrument in the T-cell transfer model. For the five items from the endoscopy scoring system introduced by Becker et al., the inter-observer reliability was also assessed as 'moderate' to 'substantial' [5].

Conclusion

There is an absence of evidence to support platelet transfusion in a patient who is not actively bleeding and the BSG and NICE specifically recommend against this practice. Although not covered by guidelines from either NICE, the BSG, or the ACG, platelet transfusions may be encountered in two other situations: platelets are often given to patients with thrombocytopenia prior to invasive procedures such as lines or drains; secondly, patients taking antiplatelet agents such as aspirin may have normal platelet counts but have a functional platelet defect and platelet transfusion may be considered.

Acknowledgement

None.

Conflict of Interest

None.

References

1. Akeel, Mohammed, Erwa Elmakki, Atef Shehata and Ahmed Elhafey, et al." Prevalence and factors associated with *H. pylori* infection in Saudi patients with dyspepsia." *Electron Physician* 10 (2018): 7279.

2. Geleto, Ayele, Catherine Chojenta, Abdulbasit Musa and Deborah Loxton. "Barriers to access and utilization of emergency obstetric care at health facilities in sub-Saharan Africa: A systematic review of literature." *Syst Rev* 7 (2018): 1-14.
3. Black, Christopher J., Lesley A. Houghton, and Alexander C. Ford. "Insights into the evaluation and management of dyspepsia: recent developments and new guidelines." *TAG* 11 (2018): 1756284818805597.
4. Hassan, Taha MM, Samia I. Al-Najjar, Ibrahim H. Al-Zahrani and Fadi IB Alanazi, et al. "*Helicobacter pylori* chronic gastritis updated Sydney grading in relation to endoscopic findings and *H. pylori* IgG antibody: Diagnostic methods." *J Microsc Ultrastruct* 4 (2016): 167-174.
5. Banke-Thomas, Oluwasola, Aduragbemi Banke-Thomas and Charles Anawo Ameh. "Utilisation of maternal health services by adolescent mothers in Kenya: Analysis of the demographic health survey 2008–2009." *Int J Adolesc Med Health* 30 (2016): 20160042.

How to cite this article: Murthy, Narayan. "Clinical Correlations between Endoscopic Findings and Histopathology." *Clin Gastroenterol J* 10 (2025): 296.