

Coffee Consumption Prevents Liver Cancer Development

Abhishek Bhurwal¹ and Vinod Rustgi^{1,2*}

¹Division of Gastroenterology and Hepatology, Rutgers Robert Wood Johnson School of Medicine, New Brunswick, NJ, USA

²Center for Liver Diseases and Liver Masses, New Brunswick, NJ, USA

Abstract

Coffee consumption has its benefits on human health. In the specific area of hepatology, previous studies have suggested that coffee decreases liver enzymes and inhibits progression of liver fibrosis. Additionally, in a recent systematic review and meta-analysis, coffee consumption was shown to have an inverse association with the development of liver cancer. That well-designed study suggested that coffee drinking would be beneficial in terms of preventing liver cancer. Coffee drinkers at risk of developing hepatocellular cancer should consider the potential benefits of drinking coffee

Keywords

Coffee • Liver fibrosis • Inverse association • Hepatocellular cancer

Commentary

The origins of the coffee bean far outdate reliable written records but historical references place its discovery in the Ethiopian highlands over 1000 years ago [1]. The popularity of coffee has continued to grow with an estimated 255 kilograms consumed every second around the world [2]. Naturally, with such wide use, the effect of coffee consumption on human health has become an important area of research. Previous studies have shown that coffee consumption decreases the risk of developing diabetes mellitus, heart disease and stroke [3-5]. In a recent meta-analysis, coffee drinking was determined to have an inverse association with the development of liver cancer [6]. Coffee consumption has been shown to influence the levels of liver enzymes such as aspartate aminotransferase, alanine aminotransferase and gamma-glutamyltransferase [7]. Additionally, there has been evidence to suggest a decrease in the rate of hepatic fibrosis progression in coffee drinkers [8].

In the September issue of Journal of Gastrointestinal and Liver diseases, Bhurwal et al. performed a systematic review and meta-analysis evaluating twenty studies to determine the impact of coffee consumption in preventing liver cancer [6]. Overall, coffee drinkers had a significantly decreased risk of development of liver cancer compared to non-drinkers (Risk Ratio (RR) 0.69; 95% Confidence Interval (CI) 0.56-0.85; $p < 0.01$). Subgroup analysis revealed that drinking more than 2 cups of coffee had significant impact in preventing liver cancer (RR 0.51; 95% CI 0.38-0.69) [6]. Even though the analysis was well designed with multiple studies across several continents, it had its limitations. There is lack of standardization of coffee preparation and serving size which prevents standardization. The duration of the coffee intake was also unknown. Therefore, further clinical trials eliminating the confounders and determining a possible dose-response would be valuable. Despite the limitations, the results from the systematic review and meta-analysis suggest a beneficial impact of coffee in preventing hepatocellular cancer.

***Address for Correspondence:** Vinod Rustgi, Center for Liver Diseases and Liver Masses, Division of Gastroenterology and Hepatology, Rutgers Robert Wood Johnson School of Medicine, New Brunswick, NJ, USA, Tel: +301-801-5814; E-mail: vinod.rustgi@rutgers.edu

Copyright: © 2020 Bhurwal A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Received October 06, 2020; **Accepted** October 20, 2020; **Published** October 28, 2020

However, the application of the coffee consumption to prevent liver cancer in daily practice is still uncertain. Even though the main compounds associated with protective influence are caffeine and chlorogenic acid, there are more than a thousand components of coffee which could be responsible [9]. Significant coffee consumption has its own adverse effects such as hypertension, tachycardia and physical dependence [10]. Further studies are needed to determine the specific ingredient(s) of coffee exerting a protective role in liver disease. Nonetheless, the data from Bhurwal et al. indicate that coffee consumption is associated with a significantly decreased risk of liver cancer development [6]. Coffee drinkers at risk of developing liver cancer may possibly benefit from coffee consumption. As every good meta-analysis, the study by Bhurwal et al. points out the need for more data from well-designed studies.

Role of Authors

A. Bhurwal and V. Rustgi wrote and edited the commentary.

Conflict of Interest

The authors declared no conflicts of interest with respect to the authorship and/or publication of this article. The authors also certify that they have no financial affiliation/interest (eg, employment, stock holdings, consultant arrangements, and honoraria in the subject matter, materials, or products mentioned in this manuscript).

Financial Disclosure

The authors received no financial support for the research and/or authorship of this article.

References

1. Illy E. "The Complexity of Coffee." *Science Am* 286(2002):86-91.
2. International Coffee Organization (ICO) Review (2018).
3. Wu, Jiang-nan, Suzanne C Ho, Chun Zhou and Wen-hua Ling et al. "Coffee Consumption and Risk of Coronary Heart Diseases: A Meta-Analysis of 21 Prospective Cohort Studies." *Int J Cardiol* 137(2009):216-25.
4. Higdon, Jane V and Balz Frei. "Coffee and Health: A Review of Recent Human Research." *Crit Rev Food Sci Nutr* 46(2006): 101-23.
5. Zhang, WL, E Lopez-Garcia, TY Li and FB Hu et al. "Coffee Consumption and Risk of Cardiovascular Events and All-Cause Mortality Among Women With Type 2 Diabetes." *Diabetologia* 52(2009): 810-7.
6. Bhurwal, Abhishek, Puru Ratta, Sho Yoshitake and Lauren Pioppo et al. "Inverse Association of Coffee With Liver Cancer Development: An Updated Systematic Review and Meta-Analysis." *J Gastrointest Liver Dis* 29(2020): 421-428.

7. Nakanishi, N, K Nakamura , K Nakajima and K Suzuki et al. "Coffee Consumption and Decreased Serum Gamma-Glutamyltransferase: A Study of Middle-Aged Japanese Men." *Eur J Epidemiol* 16(2000): 419-23.
8. Modi, Apurva A, Jordan J Feld , Yoon Park and David E Kleiner et al. "Increased Caffeine Consumption Is Associated With Reduced Hepatic Fibrosis." *Hepatology* 51(2010):201-9.
9. Yan, Yuan, Na Liu , Ni Hou and Lei Dong et al. "Chlorogenic Acid Inhibits Hepatocellular Carcinoma In Vitro and In Vivo." *J Nutr Biochem* 46(2017):68-73.
10. Mesas, Arthur Eumann, Luz M Leon-Munoz , Fernando Rodriguez-Artalejo and Esther Lopez-Garcia. "The Effect of Coffee on Blood Pressure and Cardiovascular Disease in Hypertensive Individuals: A Systematic Review and Meta-Analysis." *Am J Clin Nutr* 94(2004): 1113-26.

How to cite this article: Bhurwal, Abhishek and Vinod Rustgi. "Coffee Consumption Prevents Liver Cancer Development." *J Inflamm Bowel Dis Disor* 5 (2020): 136.