

# Comparison of Healthcare Resource Utilization (HcRU) between Age Groups of Women Diagnosed with Gynecological Cancers in the U.S

Godwin Okoye\*, R. Vincent and H. Antwi

<sup>1</sup>Department of Gynecology, University of Maryland, Baltimore, USA

<sup>2</sup>Department of Oncology, Government Medical College, Nagpur, Maharashtra, India

## Abstract

**Objective:** Previous studies have shown that older women are more likely to be diagnosed with gynecological cancers (GC) setting the precedent for more attention to be given to older women than younger women with GC. This study seeks to compare nationally representative healthcare use among younger women and older women diagnosed with gynecological cancer in the United States.

**Methods:** A retrospective cross-sectional repeated measures design was used to make comparisons between baseline characteristics of two age groups of women diagnosed with gynecological cancers. Data were extracted from the National Health Interview Survey (NHIS). A weighted sample of 2633569 responses who reported having at least one kind of GC was included from 2015 to 2018. GC was defined as those who reported having uterine, cervical, or ovarian cancer. Characteristics of patients were compared across two age groups. 1. Women less than 50 considered to be most likely premenopausal. 2. Women more than 50 considered to be most likely postmenopausal. Nine health services use were also quantified between both age groups and a multivariable logistic regression model was used to determine the likelihood of health care use among both age groups with the older women being the reference group.

**Results:** Among the baseline characteristics, alcohol use and smoking were seen to be most prevalent among the younger women with the following proportions 88% and 79.93% for alcohol use and 59.79% and 50.26% for smoking at  $p < 0.01$ . Younger women had higher obstetrics and gynecologic visits OR 2.13 (1.59-2.84), Emergency room visits OR 1.55 (1.17-2.06), Hospitalization OR 1.57 (1.15-2.14), Preventive care OR 2.49 (1.88-3.31). Adjusting for Alcohol use and smoking yielded similar odds ratio. Patterns of hospitalization increased over time for younger women and use of surgery and surgical procedures wasn't statistically significant among both age groups.

**Conclusion:** Based on this study, younger women tend to use comparable healthcare services as older women diagnosed with gynecologic cancer. Therefore, more studies should be conducted with the inclusion of younger women.

**Keywords:** Healthcare Resource • Utilization • Uterine neoplasm • Cervical neoplasm • Ovarian neoplasm • Menopausal

## Introduction

According to Centers for Disease Control and Prevention (CDC), gynecological cancer (GC) refers to when malignant cells originate from the female reproductive organs. There are five major types: cervical, ovarian, uterine, vaginal, and vulvar cancers [1]. All women are at risk of developing one or more types of GC and the risk increases with age. Between 2012-2016, about 94,000 women were diagnosed with gynecological cancer each year. It is interesting to note that Uterine cancer is the fourth most common cancer for women in the U.S with about 66% diagnosed at an early stage due to the abnormal vaginal bleeding that occurs as an early symptom [2]. Incidence rate for cervical cancer has dropped by more than 50% in the past 50 years due to screening exercises that can detect changes in the cervix before they become malignant [3]. There has been an upward trend in the incidence of ovarian cancer owing to the increasing use of oral contraceptives

and hormonal therapy for menopausal women [4]. GC generally accounts for about 11.2% of all female deaths [5].

Over the years, it has been discovered that GC is more prevalent in older, mostly post-menopausal women [6]. This has gathered much attention in the scientific community, encouraging research that could improve health outcomes for older women with GC [7,8]. However, little research attention has focused on the clinical burden of GC among younger, mostly premenopausal women [9]. This research focuses on self-reported use of health care resource of younger women with GC when compared to older women to improve their health outcomes and general survivorship.

The main objective of this study was to compare healthcare use among two distinct age groups of women with GC. i.e., women most likely to be premenopausal and women most likely to be post-menopausal.

## Materials and Methods

### Study design and data source

A retrospective cross-sectional repeated measures study was conducted to compare all patient-reported demographics and HcRU related to GC in women most likely to be pre-menopausal and women most likely to be post-menopausal using the sample adult file of NHIS (National Health Interview Survey) data from 2015 to 2018. NHIS is the principal source of information on the health of the civilian noninstitutionalized population of the United States. It is one of the major data collection programs of the National Center for Health Statistics (NCHS) which is part of the Centers for Disease Control and

\*Address for Correspondence: Godwin Okoye, Department of Gynecology, University of Maryland, Baltimore, USA; Tel: 4439359693, E-mail: gokoye8@gmail.com

**Copyright:** © 2022 Okoye G, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Date of Submission:** 02 August, 2022, Manuscript No. jomp-22-70991; **Editor Assigned:** 06 August, 2022, PreQC No. P-70991; **Reviewed:** 14 August, 2022, QC No. Q-70991; **Revised:** 16 August, 2022, Manuscript No. R-70991; **Published:** 25 August, 2022, DOI: 10.37421/2576-3857.2022.07.166

Prevention (CDC). The main objective of NHIS is to monitor the health of the United States population through the collection and analysis of data on a broad range of health topics [10].

## Study sample

We identified women who reported having a diagnosis of either of the following: Cervical, Ovarian or Uterine cancer. A few women in the cohort had more than one GC. In this study, women most likely to be premenopausal were defined as those who are less than 50 years old while women who are most likely to be post-menopausal were defined as those more than 50 years old.

## Baseline demographics, clinical and behavioral characteristics

According to the systematic review conducted by Phillips, et al., we utilized the behavioural model of health services utilization as a framework [11]. we selected our covariates based on individual and population level factors of patients in the cohort. These include Risk factors (race, marital status, obesity, pregnancy [12-14] Enabling factors (health insurance coverage, region in the U.S [15,16], Behavioral factors (smoking, drinking [17].

## Health care resource utilization

We focused on nine areas of HcRU which are: obstetrics and gynecological visit, specialist visit, general practitioner visit, hospitalization or in-patient admissions, emergency room visit, office-based services, home healthcare services, surgery and surgical procedures, preventive healthcare. Preventive health care focused on women in the cohort who have ever undergone pap smear or who have received HPV vaccination. The number of individuals who reported utilizing any of the above health service was recorded at baseline and proportions were also taken by age group.

## Statistical analysis

Baseline characteristics and outcome measures of interest were obtained for individuals in the cohort. We reported count and column percentage for categorical variables. The covariates were: race, census region, marital status, obesity, smoking, alcohol, health insurance, and pregnancy. Comparison between defined age groups of the women in the cohort was done for baseline characteristics using the Pearson's chi square ( $\chi^2$ ) test.

For HcRU, values for the nine areas mentioned above was computed and stratified amongst age group. Pearson's chi square ( $\chi^2$ ) test was used as a measure of association. Annual trends in health care resource utilization Figure 1 was extrapolated to give an insight on the independent changes happening across each year within the study period.

A univariate logistic regression model was used to compare HcRU between women most likely to be pre-menopausal (defined as women < 50 years) and women most likely to be post-menopausal (defined as women >50 years) across nine areas of health services mentioned above. Only predictors with  $p < 0.05$  were utilized for multivariate logistic regression model.

All statistical analysis were performed using SAS v9.4 ((SAS Institute Inc, Cary, NC), and statistical significance was set at  $\alpha = 0.05$ .

## Classification of evidence

This study provides class IV evidence that women most likely to be premenopausal diagnosed with GC can use comparable health care resources with postmenopausal women diagnosed with GC despite GC being more prevalent in women most likely to be postmenopausal.

## Results

Based on our data, Gynecological cancers were discovered in 1518 responses representative of 2633569 responses who reported having at least one kind of GC. We had 1047 women (1742470 weighted) who are most likely to be post-menopausal and 471 women (891099 weighted) who are most likely to be pre-menopausal. The following cases were found for the individual types of GC: 577 women reported having uterine cancer, 728 women with cervical

cancers, and 323 ovarian cancers. The mean age and standard deviation of the women with GC was: Uterine ( $64 \pm 13$ ), Cervical ( $53 \pm 15$ ) and Ovarian ( $60 \pm 15$ ). Overall, the mean age of women in the cohort was ( $58 \pm 16$ ). Statistical significance at  $p < 0.05$  was achieved for behavioral factors in the study which includes alcohol consumption and smoking.

Based on this study, Table 1A and 1B displays the baseline characteristics of women in the cohort. Alcohol use and smoking was associated with the two age groups being compared at  $p < 0.05$ . In simpler terms, pre-menopausal women were more likely to consume alcohol when compared to post-menopausal women with approximate values of 88% and 79.93% respectively. For smoking, the approximate values are 59.79% and 50.26%.

## Health care resource utilization

Table 2 displays percentage values for HcRU of interest across age groups considered to be most likely premenopausal and most likely postmenopausal. Based on our data, Premenopausal women utilized more health care for Obstetrics and Gynecological visits (51.78% vs. 33.51%), Emergency room visits (41.29% vs. 31.19%), Hospitalization (30.59% vs. 21.9%) and Preventive care (46.33% vs. 25.72%). Surgery/surgical visits was the only HcRU parameter of interest that wasn't statistically significant at  $p < 0.05$ .

Figures 1(A-I) illustrated annual trends in HcRU. A clear distinction was marked for premenopausal women in the four areas mentioned above where they tend to use more healthcare than postmenopausal women. One notable outcome of this research was the upward trend in hospitalization for premenopausal women.


Table 3 displays results from the logistic regression models comparing premenopausal women with GC and postmenopausal women with GC. Out of nine parameters of interest, premenopausal women showed significant HcRU in four of the parameters. With the following results: Obstetrics and Gynecological visits (OR 2.38, (95% CI 1.89-2.97)), Emergency room visits (OR 1.73, (95% CI 1.39-2.17)), In-patient hospitalization (OR 1.92 (95% CI 1.51-2.43)), Preventive care (OR 2.695 (95% CI 2.14-3.39)). Surgery/surgical visits was not consistent with an association between premenopausal and postmenopausal women. Similar results were observed in the sensitivity analysis displayed in Table 4.

## Discussion

This retrospective cross sectional repeated measures study used NHIS data from 2015-2018 to compare HcRU among women most likely to be premenopausal and women most likely to be post-menopausal who have had at least one diagnosis of ovarian cancer, uterine cancer, or cervical cancer.

From our baseline statistics, race wasn't statistically significant when compared with both age groups, but it is known from previous studies that being white increases the risk of gynecologic cancer [18]. It is also interesting to note that both age groups had similarity in being obese which interesting because there is an association between obesity and age [19]. Behavioral factors like drinking and smoking were found to be strongly associated with younger women with gynecological cancer which isn't surprising because of past literatures pointing out drinking and smoking as a coping mechanisms for young adult women [20].

To improve the understanding of longitudinal use patterns, we quantified HcRU changes using a combination of descriptive statistics and regression models. Overall, of nine essential area of health care use in gynecological cancer, premenopausal women were seen to significantly use preventive health care such as pap smear and HPV vaccination which is known to prevent cervical cancer or detect it at an earlier stage. This utilization pattern is consistent with a paper written by Sirovich, et al. [21]. Premenopausal women were also observed to have been more hospitalized. The trend for hospitalization was also observed to have had a continuous increase throughout the study period. Although younger women are expected to have lesser hospitalization due to expected less comorbidities of certain diseases that could be predictors of increased hospitalization [22]. Premenopausal women also had more frequent emergency room visits and visits to Obstetrics and Gynecologist maybe due



**Figure 1.** Annual trends in Health Care Resource Utilization (A-I) across women with gynecological cancer stratified by age group.

**Table 1A.** Sample demographic characteristics of women with gynecologic cancers, 2015-2018.

Baseline		Overall n= 2.63 * 10 <sup>6</sup>	Post-menopausal n= 1.74 * 10 <sup>6</sup>	Pre-menopausal n=0.89 * 10 <sup>6</sup>	P
Race n(%)	White	2.25 (85.3)	1.5 (86.26)	0.74 (83.43)	0.2636
	Not White	0.38 (14.7)	0.23 (13.74)	0.15 (16.57)	-
Marital Status n(%)	Married	1.28 (48.63)	0.86 (49.13)	0.42 (47.67)	0.68
	Not Married <sup>a</sup>	1.35 (51.36)	0.89 (50.87)	0.47 (52.33)	-
Alcohol n(%)	Never	0.46 (17.34)	0.35 (20.07)	0.11 (12)	<0.01*
	Drinker	2.17 (82.66)	1.39 (79.93)	0.78 (88)	-
Census region n(%)	Northeast	0.44 (16.79)	0.33 (18.83)	0.11 (12.81)	0.1076
	Midwest	0.68 (25.94)	0.44 (25.24)	0.24 (27.28)	-
	South	0.96 (36.4)	0.61 (34.97)	0.35 (39.18)	-
	West	0.55 (20.87)	0.36 (20.93)	0.19 (20.72)	-
BMI n(%)	Obesity	1.18 (44.71)	0.79 (45.34)	0.39 (43.47)	0.5893
	Non-Obesed	1.46 (55.29)	0.95 (54.66)	0.5 (56.52)	-
Smoking n(%)	Never	1.23 (46.52)	0.87 (49.74)	0.36 (40.21)	<0.01*
	Smoker	1.4 (53.48)	0.88 (50.26)	0.53 (59.79)	-
Pregnancy n(%)	Never	0.21 (7.83)	0.13 (7.39)	0.08 (8.67)	0.45
	≥1	2.43 (92.17)	1.61 (92.6)	0.81 (91.32)	-
Insurance n(%)	Unchanged <sup>b</sup>	1.48 (56.05)	0.99 (57.36)	0.48 (53.49)	0.2593
	Changed	1.15 (43.95)	0.74 (42.64)	0.41 (46.51)	-

**Table 1B.** The baseline characteristics of women in the cohort.

Baseline		Overall n=1518	Post-menopausal n=1047	Pre-menopausal n=471	P
Race n(%)	White	1306 (86.03)	910 (86.91)	396 (84.08)	0.14
	Not White	212 (13.97)	137 (13.09)	75 (15.92)	-
Marital Status n(%)	Married	543 (35.77)	361 (34.48)	182 (38.64)	0.1176
	Not Married <sup>a</sup>	975 (64.23)	686 (65.52)	289 (61.36)	-
Alcohol n(%)	Never	252 (16.6)	201 (19.20)	51 (10.83)	<0.001*
	Drinker	1266 (83.4)	846 (80.8)	420 (89.17)	-
Census region n(%)	Northeast	258 (17)	189 (18.05)	69 (14.65)	0.4258
	Midwest	379 (24.97)	260 (24.83)	119 (25.27)	-
	South	518 (34.12)	350 (33.43)	168 (35.67)	-
	West	363 (23.91)	248 (23.69)	115 (24.42)	-
BMI n(%)	Obesity	663 (43.68)	455 (43.46)	208 (44.16)	0.7981
	Non-Obesed	855 (56.32)	592 (56.54)	263 (55.84)	-
Smoking n(%)	Never	676 (44.53)	496 (47.37)	180 (38.22)	<0.001*
	Smoker	842 (55.47)	551 (52.63)	291 (61.78)	-
Pregnancy n(%)	Never	132 (8.7)	90 (8.6)	42 (8.92)	0.8372
	≥1	1386 (91.3)	957 (91.4)	429 (91.08)	-
Insurance n(%)	Unchanged <sup>b</sup>	912 (60.08)	412 (39.35)	194 (41.19)	0.4987
	Changed	606 (39.92)	635 (60.65)	277 (58.81)	-

N/B: \* refers to p-values that are <0.05 which entails statistical significance.

Refers to women who have never been married (i.e., excluding widowed, divorced, separated)

Refers to women who had the same insurance as the previous year.

**Table 2.** Health services use among two distinct age group of women with gynecological cancer using NHIS data.

Post-menopausal n= 1.74 * 10 <sup>6</sup>	Pre-menopausal n=0.89 * 10 <sup>6</sup>	P
0.58 (33.51)	0.46 (51.78)	<.0001*
0.86 (49.13)	0.31 (35.28)	<.0001*
1.47 (84.63)	0.64 (71.45)	<.0001*
0.54 (31.19)	0.37 (41.29)	0.0021*
0.16 (9.09)	0.03 (3.69)	0.0014*
1.64 (94.06)	0.79 (88.89)	0.0016*
0.43 (24.76)	0.22 (24.51)	0.9352
0.38 (21.9)	0.27 (30.59)	0.0039*
0.45 (25.72)	0.41 (46.33)	<.0001*

to the fact that most of them are within reproductive age and are most likely to have a partner or be willing to have children as described by Schramm, et al. [23]. Surgery/surgical visits was not associated with the age group. This could be due to hysterectomy performed as a choice treatment for gynecologic cancer. Although in past literatures [24], younger women were seen to visit

general practitioners more than older women, the reverse was the case for our current study.

Our study has several strengths, first we utilized the NHIS (National Health Interview Survey) which is the principal source of information on the health of



**Table 3.** OR comparing health care resource utilization among women most likely to be premenopausal and women most likely to be postmenopausal who have gynecological cancer.

Health Care Resource	OR (95% CI)	p value
ObGyn visits	2.13 (1.59 - 2.84)	<0.001
Specialist visits	0.56 (0.43 - 0.75)	<0.001
General Practitioner visits	0.45 (0.32 - 0.63)	<0.001
Emergency room	1.55 (1.17 - 2.06)	<0.001
Home health care	0.38 (0.21 - 0.71)	0.0022
Office-based visits	0.51 (0.33 - 0.78)	<0.001
Surgery/Surgical visits	0.99 (0.72 - 1.36)	0.9354
Hospitalization	1.57 (1.15 - 2.14)	0.0044
Preventive care	2.49 (1.88 - 3.31)	<0.001

**N/B:** Reference group is the older women (women most likely to be postmenopausal)  
**OR:** Odds Ratio, **CI:** Confidence Interval.

**Table 4.** OR comparing health care resource utilization among women most likely to be premenopausal and women most likely to be postmenopausal who have gynecological cancer (controlling for alcohol use and smoking).

Health Care Resource	Adj OR (95% CI)	p value
ObGyn visits	2.23 (1.67 - 2.96)	<0.001
Specialist visits	0.57 (0.43 - 0.76)	<0.001
General Practitioner visits	0.46 (0.33 - 0.63)	<0.001
Emergency room	1.52 (1.14 - 2.02)	<0.001
Home health care	0.39 (0.21 - 0.73)	0.003
Office-based visits	0.54 (0.35 - 0.83)	0.0052
Surgery	0.97 (0.69 - 1.36)	0.8581
Hospitalization	1.55 (1.13 - 2.11)	0.0065
Preventive care	2.54 (1.91 - 3.38)	<0.001

**N/B:** Reference group is the older women (women most likely to be postmenopausal)  
**OR:** Odds Ratio, **CI:** Confidence Interval.

civilian noninstitutionalized population in the United States. This survey data is able to capture the many demographic and socioeconomic characteristics of participants. It also finds its use in monitoring trends in illness and disability. To the best of our knowledge, this study is the first to make comparison between healthcare resource utilization of women who are most likely to be premenopausal and women most likely to be postmenopausal diagnosed with gynecological cancer in the U.S. As such, this could serve as the basis for other study to be conducted.

There are a few limitations to this study. First, some important variables were missing in the previous years. As a result of missing variables, our study was limited to four years. One notable inconsistency in covariates was use of chemotherapy and absence of specific procedures like hysterectomy. Pap smear and HPV vaccination was recorded in the 2018 data and with all the response missing. Second, given the nature of this study as self-reported questionnaire, it is subject to non-response and reporting bias. However, according to Rosenman, et al. [25], there is some agreement between self-reported questionnaires and medical records. Third, due to the short survival rate and late detection of some gynecological cancers, this population-based survey may have underestimated the prevalence.

## Conclusion

This study presents some evidence that younger women who are most likely to be premenopausal utilize healthcare in a comparable way as older women who are most likely to be postmenopausal does. As such, research relating to gynecological cancer should not be limited to the most prevalent group.

## References

- Centers for Disease Control and Prevention. "Basic information about gynecologic cancers. Centers for Disease Control and Prevention." (2021).
- Tarver, Talicia. "Cancer facts & figures 2012. American cancer society (ACS) (2012): 366-367.
- "Cervical Cancer: Statistics." Cancer.Net. (2022).
- American Society of Clinical Oncology. "Ovarian, fallopian tube and peritoneal cancer: Statistics." Cancer. Net. (2020).
- American Cancer Society. "Cancer facts & figures 2022."
- Sundar, Sudha, Richard D. Neal and Sean Kehoe. "Diagnosis of ovarian cancer." *BMJ* 351 (2015).
- Dumas, Lucy, Alistair Ring, John Butler and Tania Kalsi, et al. "Improving outcomes for older women with gynecological malignancies." *Cancer Treat Rev* 50 (2016): 99-108.
- "Retooling for an Aging America: Building the Health Care Workforce."
- Mattsson, Elisabet, Lisa Ljungman, Kim Einhorn and Inger Sundström Poromaa, et al. "Perceptions of care after end-of-treatment among younger women with different gynecologic cancer diagnoses – a qualitative analysis of written responses submitted via a survey." *BMC Womens Health* 20 (2020): 1-9.
- Bureau, U. S. C. (2021, October 8). National Health Interview Survey (NHIS).
- Phillips, Kathryn A., Kathleen R. Morrison, Ronald Andersen and Lu Ann Aday. "Understanding the context of healthcare utilization: assessing environmental and provider-related variables in the behavioral model of utilization." *Health Serv Res* 33 (1998): 571.
- Kato, Ikuko, Suketami Tominaga, and Chikahiro Terao. "An epidemiological study on marital status and cancer incidence." *JPN J Cancer Res* 80 (1989): 306-311.
- Madison, Terri, David Schottenfeld, Sherman A. James and Ann G. Schwartz, et al. "Endometrial cancer: Socioeconomic status and racial/ethnic differences in stage at diagnosis, treatment, and survival." *Am J Public Health* 94 (2004): 2104-2111.
- Kalliala, Ilkka, Georgios Markozannes, Marc J. Gunter and Evangelos Paraskevidis, et al. "Obesity and gynaecological and obstetric conditions: Umbrella review of the literature." *BMJ* 359 (2017).
- Yabroff, K. Robin, Katherine Reeder-Hayes, Jingxuan Zhao and Michael T. Halpern, et al. "Health Insurance Coverage Disruptions and Cancer Care and Outcomes: Systematic Review of Published Research." *J Natl Cancer Inst* 112 (2020): 671-687.
- Fogleman, Amanda J., Georgia S. Mueller and Wiley D. Jenkins. "Does where you live play an important role in cancer incidence in the U.S.?" *Am J Cancer Res* 5 (2015): 2314.
- Klein, William M.P., Mary E. O'Connell, Michele H. Bloch and Susan M. Czajkowski, et al. "Behavioral research in cancer prevention and control: emerging challenges and opportunities." *J Natl Cancer Inst* 114 (2022): 179-186.
- Collins, Yvonne, Kevin Holcomb, Eloise Chapman-Davis and Dineo Khabele, et al. "Gynecologic cancer disparities: A report from the health disparities taskforce of the society of gynecologic oncology." *Gynecol Oncol* 133 (2014): 353-361.
- Jura, Magdalena and Leslie Kozak. "Obesity and related consequences to ageing." *Age (Dordr)* 38 (2016): 1-18.
- De Angelis, Cristina, Antonio Nardone, Francesco Garifalos and Claudia Pivonello, et al. "Smoke, alcohol and drug addiction and female fertility." *Reprod Biol Endocrinol* 18 (2020): 1-26.
- Sirovich, Brenda E and H. Gilbert Welch. "The frequency of Pap smear screening in the United States." *J Gen Intern Med* 19 (2004): 243-250.
- Diaz-Montes, Teresa P and Robert L. Giuntoli. "Volume-based care among young women diagnosed with uterine cancer." *ISRN Surg* 2011 (2011).
- Schramm, Katharina, Juliane Nees, Janine Hoffmann, Thomas Bruckner, et al. "Emergency consultations in obstetrics: Identification of decisive, contributing and associated factors." *Arch Gynecol Obstet* 302 (2020): 821-828.
- Ezendam, Nicole P.M., Kim A.H. Nicolaije, Dorry Boll, Marnix L.M. Lybeert, et al. "Health care use among endometrial cancer survivors: A study from profiles, a population-based survivorship registry." *Int J Gynecol Cancer* 23 (2013): 1258-1265.
- Rosenman, Robert, Vidhura Tennekoon and Laura G. Hill. "Measuring bias in self-reported data." *Int J Behav Healthc Res* 2 (2011): 320.

**How to cite this article:** Okoye, Godwin, R Vincent and H Antwi. "Comparison of Healthcare Resource Utilization (HcRU) between Age Groups of Women Diagnosed with Gynecological Cancers in the U.S." *J Oncol Med & Pract* 6 (2022):166.