

Determination the Origin of Mineralization in the Coastal Aquifer Northeast Tunisia by Isotopic Method

Mzoughi Aroua^{1*}, Ben Hamouda Mohamed Fethi² and Bouhlel Salah¹

¹Laboratory of Mineral Resources and Environment, Faculty of Sciences of Tunis, University of Tunis El Manar, Tunis, Tunisia

²Hydrology and Isotopic Geochemistry Unit, CNSTN Sidi-Thabet Technology Center, Tunisia

Abstract

A geochemical and isotopic techniques were undertaken to characterize groundwater in Northeast Tunisia. Hydrogeochemical investigations demonstrated that groundwater can be classified into different water facies. The Ras Djebel-RafRaf aquifer showed a (Ca-Cl-SO₄) and (Na-Cl-NO₃) water type. Data inferred from ¹⁸O and deuterium isotopes in groundwater samples indicated recharge with modern rainfall. Water characterized by lower δ¹⁸O and δ²H values is interpreted as recharged by non-evaporated rainfall originating from Mediterranean air masses from Mediterranean air masses at higher altitude. However, water with relatively enriched δ¹⁸O and δ²H contents is thought to reflect the occurrence of an evaporation process related to the long-term practice of flood irrigation.

Keywords: Hydrogeology; Geochemistry; Coastal aquifer; Tunisia

Introduction

The region of Bizerte (Figure 1) is situated in boundary Northern of Tunisia. It forms with the Cap Blanc the most advanced point of Africa. It is exposed to humid winds from the northwest and West, and receives an average rainfall of 570 mm/year [1], This coastal occupies an area of 50 Km². It is limited by Djebels bouchoucha, Touchela and Bab Banzart in NW, Djebels Ennadour and Demna in South, the Mediterranean Sea in the North and NE.

This zone is influenced by a sub-humid Mediterranean climate. The annual temperature is around 18° C. The ETP, about 1197 mm/year. This basin is drained by many wadis. The use of water is essential since the Ras Djebel is an agricultural region by excellence, in Fact the major part of population practise agricultural activities.

In result, this aquifer undergoes increased anthropic pressure through overexploitation and construction of two dams at Wadi Beni Ata and Wadi Shaab Eddoud.

Geology Setting

From a geologic standpoint, the main geologic features of this study area are [2,3].

(a) Upper Cretaceous, represented by the lower senonian and the lower campanian which is divided into two distinct entities: the marls senonian and the calcareous marl campanian.

(b) Lower Paleocene, represented by grey marls.

(c) Eocene, formed by limestone.

(d) Mio-Pliocene, formed by continental materials: Marls, clays, conglomerates, gypsum and sands.

(e) Quaternary, represented by the villafranchian, formed essentially by limestone.

Materials and Methods

For the evaluation of groundwater quality, 50 water samples were collected from the shallow aquifer of Ras Djebel plain during October (Figure 1), the groundwater samples were analysed for chemical and isotopic composition.

Measurement of temperature, electrical conductivity and pH were

measured in the field. Major elements (SO₄²⁻, Cl⁻, NO₃⁻, Ca²⁺, Mg²⁺, Na⁺ and K⁺) were analysed at the laboratory of National Center for Nuclear Sciences and Technologies. These elements are expressed in mg/L⁻¹. Salinity was analyzed by evaporating a specified volume of water sample in a graduated capsule for 24 h in an incubator at 105°C.

The dry residue concentration was determined by subtracting the final mass from the initial mass of the sample that was placed in the incubator. Ca²⁺ and Mg²⁺ were measured by Liquid Ion Chromatography while K⁺ and Na⁺ were measured by flame photometer.

The results of chemical analyses of the investigated groundwater samples are presented in Table 1.

Results and Discussion

Hydrochemistry

Physical-chemical parameters: The overall chemical characteristics of groundwater are presented in Table 1. The pH values range from 6, 88 to 8, 3 with an average value of 7, 3. The temperature values are between 20, 2°C and 25°C. The values measured on the surface water are influenced by the surface air temperatures.

The total dissolved solids and the Electrical Conductivity (EC) values range from 1, 03 g/L to 12, 96 and 0, 2 to 0, 7 μS/cm. The spatial distribution maps (Figure 2) shows that salinity increase in the direction of groundwater flow. The Electrical Conductivity (EC) is well correlated with the total dissolved solids (TDS).

The factor of correlation is in the order of 1.

Major ion chemistry:

***Corresponding author:** Mzoughi Aroua, Unit of Geology and Applied Geochemistry (UR11ES16), Department of Geology, Faculty of Sciences of Tunis, University of Tunis El Manar, 2092 Tunis, Tunisia, Tel: +21621482388; E-mail: Arwa_mzoughi@yahoo.com

Received April 05, 2017; Accepted April 27, 2017; Published May 04, 2017

Citation: Aroua M, Fethi BHM, Salah B (2017) Determination the Origin of Mineralization in the Coastal Aquifer Northeast Tunisia by Isotopic Method. Hydrol Current Res 8: 273. doi: [10.4172/2157-7587.1000273](https://doi.org/10.4172/2157-7587.1000273)

Copyright: © 2017 Aroua M, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

N°èch	pH	Cond	sec	Ca	Mg	Na	K	SO ₄	Cl	NO ₃	HCO ₃
1	7.2	2.41	1567	220	40.32	328.9	8.19	324.52	479.25	23.3	351.54
2	6.9	5.37	3491	368	86.64	749.8	9.75	845.24	955.7	70.59	423.46
3	7.1	2.4	1560	180	32.88	280.6	28.47	333.26	323.1	97.4	572.26
4	7.2	2.15	1398	200	32.4	230	8.59	349.32	323.15	152.1	353.4
5	7.3	2.46	1599	142	62.16	331.2	8.97	467.31	370.23	86	523.9
6	7.7	2.4	1560	106	65.52	354.2	15.6	402.26	388.57	144.4	416.64
7	7.1	2.49	1619	156	48.48	299	30.81	354.42	455.79	90.87	407.34
8	8	2.92	1898	201	66.48	386.4	9.75	573.34	556.17	19.11	391.84
9	7.1	3.82	2483	137	104.64	616.4	7.02	492.12	753	40.17	701.84
10	7.5	4.04	2626	196	47.76	464.6	31.98	534.34	657.9	29.15	502.2
11	7.2	1.66	1079	89	28.32	149.5	72.15	202.16	238.48	49.92	221.34
12	7.2	3.56	2314	114	59.76	595.7	24.57	464.89	648.78	21.35	513.98
13	7	6.7	4355	319	67.44	1299.5	85.8	986.48	1567.7	41.24	544.36
14	7	5.37	3491	401	104.64	832.6	93.6	854.89	1327.5	12.68	564.82
15	7.2	2.93	1905	222	26.64	503.7	3.51	452.14	670.24	45.24	550.56
16	7.2	2.81	1827	260	30.96	411.7	2.34	335.22	660.9	47.58	412.3
17	7	3.36	2184	252	30.96	496.8	6.24	521.76	608.74	15.11	626.2
18	7.1	3.22	2093	178	66.72	397.9	31.98	254.72	632.97	48.26	574.12
19	7	3.25	2113	270	138.72	328.9	38.22	443.93	733.48	57.82	515.84
20	7.1	3.3	2145	250	104.88	368	23.79	459.82	806.07	54.21	493.52
21	6.9	2.97	1931	205	86.88	393.3	17.16	304.88	744.33	34.42	512.74
22	7	3.31	2152	263	119.04	342.7	7.41	350.4	843.33	53.24	550.56
23	7.3	1.44	936	66	39.36	138	18.33	160.01	147.61	73.22	288.3
24	7.2	3.4	2210	240	47.52	446.2	13.26	526.93	560.96	60.16	600.78
25	6.9	7.09	4609	542	207.84	984.4	8.19	987.69	1742.3	99.94	485.46
26	7	4.03	2620	277	124.08	517.5	3.12	499.09	971.78	99.84	512.12
27	7.1	3.66	2379	315	0.48	471.5	3.9	466.06	561.69	82.49	396.18
28	7.1	3.6	2340	322	14.16	427.8	15.99	380.74	691.25	22.72	458.18
29	7.1	5	3250	357	58.32	611.8	27.3	550.1	1046	77.03	522.66
30	7.1	3.55	2308	338	24	577.3	19.11	282.92	1019.3	83.07	556.14
31	7.1	4.17	2711	318	62.64	738.3	16.38	599.52	1118.6	47.97	515.22
32	6.9	3.94	2561	386	14.64	489.9	8.19	629.57	821.75	46.12	434.62
33	7.2	2.94	1911	200	15.6	411.7	30.42	386.99	550.79	17.36	408.58
34	7	3.75	2438	480	10.8	397.9	6.24	275.25	1145.7	45.14	521.42
35	7.2	3.56	2314	354	17.28	473.8	7.8	393.06	724.18	43.58	535.06
36	7.6	3.22	2098	282	19.92	473.8	5.46	408.41	803.41	16.58	480.5
37	7.1	2.78	1807	251	73.44	202.4	9.75	297.5	398.4	59.48	442.68
38	7.1	3.6	2340	205	81.6	395.6	13.65	339.69	579.64	61.23	555.52
39	7.3	2.31	1502	136	45.6	273.7	13.65	169.04	368.54	17.55	391.84
40	7.8	0.27	174	50	7.68	6.9	8.19	53.45	42.5	11.41	65.1
41	7.4	2.98	1937	244	98.64	404.8	7.02	429.47	671.63	7.51	559.24
42	8.3	0.99	607	83	20.4	92	24.96	130.58	147.25	46.51	132.06
43	7.3	2.28	1482	113	15.36	292.1	9.36	235.8	252.96	45.05	499.72
44	7.1	3.46	2249	282	119.04	342.7	16.38	623.95	540.14	11.31	532.58
45	7.4	4.43	2880	208	139.44	719.9	17.16	744.19	1025.3	16.58	516.46
46	7.5	3.42	2223	222	13.44	427.8	24.18	514.97	668.82	32.08	301.32
47	7.3	4.2	2730	300	42.72	625.6	31.59	570.49	933.48	92.63	420.36
48	7.1	6.3	4095	253	74.88	1016.6	56.94	328.8	1674.2	40.07	603.26
49	7.2	4.79	3114	228	28.08	901.6	64.35	274.27	1041.3	58.79	539.4
50	7.4	1.56	1014	111	15.12	142.6	107.64	152.75	163.99	126.95	272.18

Table 1: Geochemical data for the studied aquifer.

**Chloride and sodium*

Concentrations of sodium and chlorides vary respectively between 6, 9 to 1299, 5 mg/L and 42, 5 to 1742 mg/L. For both Na⁺ and Cl⁻, the highest concentration levels characterize the wells situated on the border of the sea (P13, P25 and P48). The spatial distribution of chloride and sodium concentrations illustrate a similar evolution to that salinity: they both increase in the direction of groundwater flow.

**Calcium*

Ca²⁺ concentrations ranged between 50 and 542 mg/L. The most elevated value was measured in well P25, while lowest level was recorded in well P40. The water of this coastal aquifer is moderately rich in calcium. The mostly analyzed waters are pure, which is in conformity with pH values of these waters. Calcium and pH have an influence on water aggressiveness [4].

Ech	RDJ1	RDJ2	RDJ3	RDJ5	RDJ6	RDJ8	RDJ9	RDJ10	RDJ11	RDJ12	RDJ13	RDJ14	RDJ15	RDJ16	RDJ17	RDJ18	RDJ21	RDJ23	RDJ24	RDJ26
$\delta^2\text{H}$	-27,7	-26,7	-27,6	-30,8	-30,1	-32,8	-31,1	-25,1	-31,1	-31,6	-26,7	-24,3	-27,2	-27,0	-25,7	-28,2	-23,4	-31,8	-28,5	-26,6
$\delta^{18}\text{O}$	-4,4	-3,6	-4,5	-5,0	-4,6	-5,9	-5,0	-4,0	-4,9	-5,0	-4,4	-4,0	-4,5	-4,1	-4,1	-4,2	-4,2	-5,3	-4,8	-4,1

Table 2: Isotopic data for the studied aquifer.




Figure 1: Geological map of RafRaf-Ras Djebel plain.




Figure 2: Map of salinity.

*Magnesium

Mg²⁺ concentrations in Ras Djebel-RafRaf aquifer fluctuated between 0, 48 and 207 mg/L. The highest values were recorded for well P25, whereas the lower one was recorded for well P40.

*Potassium

K⁺ concentrations are relatively low compared to the concentrations of other cations which values vary between 2, 34 and 107, 64.

*Sulfates

The sulphates concentrations of the water in the Ras Djebel-RafRaf aquifer varied between 53, 45 and 987, 69 mg/L. The highest concentrations were measured for well P25 and P13 while the lowest value is characterized by well P40.

*Nitrates

NO₃⁻ concentrations varied between 7, 51 and 144, 4 mg/L. The highest concentration was measured for well P6 while the lowest value was characterized by well P41.

Forty two percent of groundwater samples taken during this study, show nitrate concentrations exceeding the maximum European admissible nitrate concentration limit in drinking water (50 mg/L).

The examination of nitrate distribution map (Figure 3) reveals that high nitrate concentrations appear to be related to the fertilizer application and long term flood irrigation practises, highlighting the significant contribution of the return flow of irrigation waters to the degradation of groundwater [5-8].

Saturation Index

Water saturation states for minerals as anhydrite (SI anh), gypsum (SI gyp), aragonite (arg) and calcite (cal) were computed. If the SI is <0, dissolution is considered as the dominant process for the related mineral and when is >0, precipitation of mineral is likely to be occurring in the system [9].

The majority of samples are supersaturated towards aragonite from which the carbonates are precipitated. This supersaturation of water towards these carbonates minerals such as aragonite and calcite shows that the possibility of dissolution of these minerals by the waters is not possible and that cannot contribute to the acquisition of mineralization [10]. In addition, the majority of samples of this groundwater are under saturated towards gypsum.

Piper diagram

The piper diagram [3] (Figure 4) was determined to precisely recognize the different groundwater facies. Nitrate concentration was taken into account when plotting this diagram because of its relative abundance in the groundwater.

The data represented in this groundwater samples are classified into two major groups: (Ca-Cl-SO₄) and (Na-Cl-NO₃) water type.

Relationships between TDS and major elements

Plots of various major elements as a function of TDS value were drawn to determine the contributing elements to groundwater mineralization and the major hydrogeochemical process. Shows that Cl⁻, SO₄²⁻ and Na⁺ are well correlated with TDS presenting very close values of 1 (respectively: 0, 88; 0, 67 and 0,87). In result chlorides, sulphates and sodium are the main contributors to groundwater mineralization.

However, the contents of calcium and bicarbonate participate significantly is not determining role in the acquisition of mineralization. In addition to TDS, the correlation matrix established between sulphates and Cl⁻ ion (Figure 5A) shows that the majority of points are situated above the line (1:1). The increase of Cl⁻ in comparison to sulphates proves that the Cl⁻ has probably two origins: It is linked to a marine intrusion or dissolution of the halite. This hypothesis is confirmed by the relation between the Cl⁻ and Na⁺ ions (Figure 5B).

The relationship between Ca²⁺ and Cl⁻ ions (Figure 5C) shows excess of Cl⁻ against Ca²⁺, which demonstrates a process of precipitation of gypsum (CaSO₄·2H₂O) or anhydrite (CaSO₄). Besides, the effect of precipitation is felt in the relationship between the SO₄²⁻ and Ca²⁺ ions (Figure 5D) where the sulphate deficiency is related to the ion exchange.

Isotopic composition of groundwater

Isotope geochemistry techniques are valuable tools in investigating many problems in hydrology and evaluating hydrogeological and hydro chemical controlling mechanisms in any groundwater system [11,12].

They provide significant insights into water origin, recharge circumstances (time and location), water flow directions and residence times [11].

*Stable isotope of water (δ¹⁸O, δ²H):

Stable isotopes (δ¹⁸O, δ²H) have conservative proprieties and provide information on the groundwater recharge process (Table 2) [13]. They can offer an evaluation of physical process that affects water masses, such as evaporation and mixing [14].

The isotopic compositions of the groundwater samples from the Grombalia shallow aquifer range from -5, 3‰ to -3, 6‰ for δ¹⁸O and from -31‰ to -24, 3‰ for δ²H.

The correlation diagram of δ¹⁸O/δ²H (Figure 6) shows the position of all samples relative to the Global Meteoric Water Line (GMWL: δ²H=8δ¹⁸O+10) [15] and the Local Meteoric Water Line of the Tunis-Carthage (LMWL: δ²H= 8δ¹⁸O+12, 4) [16,17].

Nevertheless, these groundwater samples (Figure 6A) can be further divided into two groups:

- The first group, which is placed below the GMWL is probably the consequence of the infiltration of an evaporated component likely deriving from the return flow of irrigation water.
- The second group, placed between the GMWL and the LMWL, shows that the precipitation ensuring the recharge of aquifer from a mixture of oceanic and Mediterranean vapour masses.

The plot of chloride versus δ¹⁸O (Figure 6B) indicates the different processes responsible for the variation of groundwater salinity.

Some water samples show a correlation between these two elements. It is probable that the dissolution mechanism takes precedence over evaporation in the acquisition of the mineralization.

Conclusion

Based on hydrogeological characteristics the aquifer of Ras Djebel Northeast Tunisia can be classified into two groups: Ca-Cl-SO₄ and Na-Cl-NO₃ water type. The acquisition of mineralization mainly through natural mineralization mechanisms such as the ion exchange and the geology of groundwater. Nevertheless, anthropogenic process related to agricultural practices also plays an important role in the salinization of groundwater.




Figure 3: Map of salinity.






Figure 4: Piper diagram.



4. Gilbert JR (1990) L'agressivité des eaux de surface en région tropicale, ses conséquences sur le réseau de distribution. TSM, l'eau. 85^{ème} année, N°3, pp: 47-86.
5. Charfi S, Zouari K, Feki S, Mami E (2012) Study of variation of groundwater quality in a coastal aquifer in North eastern Tunisia using multivariate factor analyses. Quaternary International 302: 199-209.
6. Hadas A, Sagiv B, Haruvy N (1999) Agricultural practices, soil fertility management modes and resultant nitrogen leaching rates under semi-arid conditions. Agricultural Water Management 42: 81-95.
7. Oenema O, Boers PCM, Van Erort MM, Fraters B, Vander Meer HG, et al. (1998) Leaching of nitrate from agriculture to groundwater: the effect of policies and measures in the Netherlands. Environmental Pollution 102: 471-478.
8. Rouvier H (1977) Géologie de l'extrême Nord Tunisien: Tectonique et paléogéographique superposés à l'extrémité de la chaîne Maghrébine. Thèse de 3^{ème} cycle, Univ Paris VI, 2: 898.
9. Freeze RA, Cherry JA (1979) Groundwater. Prentice Hall Inc., Englewood Cliffs, NJ, USA, p: 604.
10. Ben Hamouda MF (2008) Approche hydrogéochimique et isotopique des systèmes aquifères de CapBon, Cas des nappes de la côte orientale et d'El Haouaria, Tunisie. Thèse en sciences agronomiques de l'INAT.
11. Clark I, Fritz P (1997) Environmental isotopes in Hydrogeology. Lewis Publishers, New York, USA.
12. Tijani MN, Loehnert EP, Uma KO (1996) Origin of saline groundwater's in the Ogoja area, lower Benue trough, Nigeria. Journal of African Earth Sciences 23: 237-252.
13. Gat JR (1971) Comments on the stable isotope method in the regional groundwater investigation. Water Resources Research 7: 980-993.
14. Geyh MA (2000) An Overview of ¹⁴C Analysis in the Study of Groundwater. Radiocarbon 42: 99-114.
15. Craig H (1961) Isotopic variation in meteoric water. Science 133: 1702-1703.
16. Singh B, Singh Y, Sekhon GS (1995) Fertilizer-N uses efficiency and nitrate pollution of groundwater in developing countries. Journal of Contaminant Hydrology 20: 167-184.
17. Zouari K, Aranyosy JF, Mamou A, Fontes JC (1985) Étude isotopique et géochimique des mouvements et de l'évolution des solutions de la zone aérée des sols sous climat semi-aride (sud tunisien).

References

1. Balti M (1986) Notes sur l'exploitation de la nappe phréatique du Ras Djebel Rapport interne DGRE. réf 3/27, p: 19.
2. Burrollet PF (1951) Étude géologique des bassins Mio-Pliocène du Nord Est de la Tunisie. Ann Mines et Géol, n°7, 82p+ annexes.
3. Piper AM (1944) A graphic procedure in the geochemical interpretation of water analyses. Transactions, American Geophysical Union 25: 914-923.