

Development and Validation of Spectrophotometric Methods for the Determination of Mesalazine in Pharmaceutical Formulation

Fatima Altayib Alasha Abdalla and Abdalla Ahmed Elbashir*

Chemistry Department, Faculty of Science, University of Khartoum, Sudan

Abstract

Two simple, accurate and precise spectrophotometric methods for the quantitative analysis of mesalazine (MSZ) in pharmaceutical formulation have been described. The first method (A) is based on the charge transfer reaction with alizarin red sulphonate (ARS) in the solution of pH 8.0 to form a violet product showing maximum absorbance at 600 nm. The second method (B) is a derivatisation method involving reaction of MSZ with 1,2-naphthoquinone-4-sulphonate (NQS) in alkaline medium at pH 12.0 to form an orange product exhibiting maximum absorbance at 470 nm. All variables affecting the reactions were studied and optimized. Beer's law was obeyed in the concentration ranges of 15-97.5 and 2-22 µg/mL for methods A and B, respectively. The molar absorptivity, Sandell sensitivity, detection and quantification limits were calculated. The two methods were successfully applied to the determination of MSZ in pharmaceutical formulation.

Keyword: Spectrophotometric; Mesalazine; Pharmaceutical formulation; 1,2-naphthoquinone-4-sulphonate (NQS)

Introduction

Mesalazine (MSZ), chemically known as 5-amino-2-hydroxybenzoic acid, is an anti-inflammatory drug used to treat and also maintain the remission of mild to moderate ulcerative colitis or Crohn's disease [1,2]. MSZ has been shown to be a potent scavenger of reactive oxygen species that play a significant role in the pathogenesis of inflammatory bowel disease, inhibition of natural killer cell activity, inhibition of antibody synthesis, inhibition of cyclooxygenase and lipoxygenase pathways and impairment of neutrophil function [3,4]. A number of analytical methods have been reported for the determination of MSZ in pharmaceutical dosage forms and biological fluids including spectrofluorometric [5], micellar electrokinetic chromatography [6], differential pulse voltammetry [7], HPLC [4,8-12], LC/MS/MS [13] and spectrophotometric [14-20]. Chromatographic methods have been extensively used and recommended. However these methods generally require complex and expensive equipment, provision for use and disposal of solvents, labor-intensive sample preparation procedures and personal skills in chromatographic techniques. Spectrophotometric methods are the most convenient technique because of their inherent simplicity, high sensitivity, low cost, and wide availability in quality control laboratories. However, the spectrophotometric methods [14-20] that have been reported for the determination of MSZ in their pharmaceutical formulations were associated with some major disadvantages, such as lack of sensitivity, tedious extraction procedures, heating, cooling and time consumption. In this paper, development and validation of two new spectrophotometric methods for the determination of MSZ in pharmaceutical tablets that overcome these drawbacks will be described.

Materials and Methods

Apparatus

All of the spectrophotometric measurements were made with a Double beam UV 1800 ultraviolet visible spectrophotometer provided with matched 1-cm quartz cells (SHIMATZU- Japan) with temperature maintained at 25°C. pH was determined using a model pH211 pH meter (Hanna, Italy). Thermostatically controlled water-bath (LAUDA, ecoline RE 220).

Materials

All chemicals were of analytical or HPLC grade. Distilled water was used in all experiments. Chemicals (suppliers) were as follows: The standard of MSZ was obtained from Sigma-Aldrich, tablet formulation Pentasa (Pharbil Pharma Bielefeld, Germany (Marketing Authorization Holder : Ferring GmbH, Kiel, Germany) was purchased from a local pharmacy with labeled amount 500 mg MSZ, Alizarin Red Sulphonate (ARS) [Hopkin & Williams LTD (England)], sodium 1,2-naphthoquinone-4-sulphonate (NQS) was purchased from [Aldrich Chemical Co., St. Louis, USA].

Preparation of reagents and solutions

Stock standard solution of Mesalazine (MSZ): An accurately weighed amount 0.100 g of MSZ was dissolved in 30 mL water then transferred into a 100 mL standard flask, completed to the mark with ethanol for method A (ratio of water to ethanol was 30:70) and distilled water for method B to obtain a stock solution of (1000 µg/mL). The stock solutions were further diluted with the same solvent to obtain working solutions of (150 and 200 µg/mL), for method A and B respectively.

Standard solution of Alizarin Red Sulphonate (ARS): An accurately weighed amount (0.5134 g) of ARS was dissolved in 30:70 water and ethanol, transferred into a 100 mL standard flask, completed to the mark with the same solvent to obtain a solution of 15×10^{-3} mol/L. The solution was freshly prepared and protected from light during use.

Sodium 1, 2-Naphthoquinone-4-Sulfonic Solution (NQS): An accurately weighed 0.1041 g of NQS was dissolved in distilled water, transferred into a 100 mL standard flask and diluted to the mark with

*Corresponding author: Abdalla A Elbashir, Chemistry Department, Faculty of Science, University of Khartoum, PO Box 321, Khartoum, 11115, Sudan, E-mail: hajaae@yahoo.com

Received January 24, 2014; Accepted March 21, 2014; Published March 24, 2014

Citation: Alasha Abdalla FA, Elbashir AA (2014) Development and Validation of Spectrophotometric Methods for the Determination of Mesalazine in Pharmaceutical Formulation. Med chem 4: 361-366. doi:10.4172/2161-0444.1000166

Copyright: © 2014 Alasha Abdalla FA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

distilled water and mixed well to prepare (4×10^{-3} mol/L). The solution was freshly prepared and protected from light during use.

Buffer solutions: Buffer solution of pH 8.0 was prepared by adding 4.49 mL of 0.1 mol/L HCl and 95.51 mL of 0.1 mol/L Na_2HPO_4 and adjusting to pH 8.0, pH 12.0 was prepared by adding 100 mL 0.05 M Na_2HPO_4 and 53.8 mL of 0.1 M NaOH, and made up to 200 mL with distilled water. Other buffer solutions were also prepared according to literature method.

Sample solution: A sample of finely powdered tablet nominally equivalent to 0.100 g MSZ was dissolved in about 30 mL distilled water in a 100 mL volumetric flask. Then swirled, sonicated for 15 min, then filled to the volume with ethanol for method A (ratio of water to ethanol was 30:70) and distilled water for method B, The contents were mixed well and filtered. This prepared solution was diluted quantitatively to obtain a suitable concentration for the analysis.

Assay procedures

Method using ARS (Method A): Aliquots of MSZ solution were added to 10 mL volumetric flasks to give final concentrations of (15-97.5 $\mu\text{g/mL}$). 2.0 mL Buffer solution of pH 8.0 was added followed by 1.0 mL ARS solution (15×10^{-3} mol/L). The reaction was allowed to proceed at temperature 40°C for 10 min after which the reaction mixture was made up to the mark with water and the absorbance was measured at 600 nm against a water blank similarly prepared.

Method using NQS (Method B): Aliquots of MSZ solution was added to 10 mL volumetric flasks to give final concentrations of (2-22 $\mu\text{g/mL}$). 1.0 mL Buffer solution of pH 12.0 was added followed by 1.0 mL NQS solution (4×10^{-3} mol/L). The reaction was allowed to proceed at room temperature for 10 minutes after which the reaction mixture was made up to the mark with water and the absorbance was measured at 470 nm against blank similarly prepared.

Stoichiometry of the reaction (Job's method)

Stoichiometry of the reaction of MSZ with ARS and NQS was established by Job's method of continuous variation [21]. Equimolar aqueous solutions of MSZ with ARS and NQS (3×10^{-3} , 2.5×10^{-3} mol/L respectively) were prepared in 10 mL volumetric flasks containing complementary proportions: (0:10, 1:9, 2:8, 3:7, 4:6, 5:5, 6:4, 7:3, 8:2, 9:1, 10:0). The solutions of method A and B were further treated as described under the general recommended procedure.

Results and Discussion

Method A

Alizarin Red S (ARS) has been used as a color-developing reagent in the spectrophotometric determination of pharmaceutical amines [22-26]. The reaction of MSZ with ARS to produce a violet colored charge transfer product of the n- π type, this compound is considered to be an intermediate molecular association complex which dissociates in the corresponding radical anion in the solvent.

At optimum conditions the radical anion (absorbing species) in the medium after mixing of the reagent and showed maximum absorption at 600 (Figure 1). The absorbance was found to increase linearly with increasing concentration of MSZ.

The effect of pH was studied by forming the colored product in the presence of various pH the absorbance of the proton transfer product was measured. Figure 2 shows that at pH from 1.0 to 4.0, no MSZ-ARS product was formed whereas at pH greater than pH 4.0 the absorbance

increased rapidly with increasing pH up to pH 8.0 and then decrease, after pH greater than 8.0, the decrease in absorbance of the product was observed. The maximum absorbance was attained at pH value of 8.0.

The reaction was found to be dependent on ARS concentration with the absorbance of the reaction solution increasing as the ARS concentration increased. Maximum absorbance was attained on using 1.0 mL of 15×10^{-3} mole/L ARS above which it decreased. Therefore, this concentration was used in all subsequent work Figure 3.

The reaction time was determined by following the color development at room temperature and in thermostatically controlled water-bath adjusted at temperature in range from ($30-90^\circ\text{C}$). The highest absorbance is obtained 40°C for 10 minutes.

Method B

MSZ exhibits maximum absorbance (λ_{max}) at 298 nm. Being in the ultraviolet, absorbance at this wavelength is susceptible to interference from co-extracted excipients in the tablet formulation. Accordingly,




Figure 1: Absorption spectra of MSZ (50 $\mu\text{g/mL}$) against blank (a), ARS against blank (b), and the reaction product of MSZ (50 $\mu\text{g/mL}$) with ARS against reagent blank (c).





Figure 2: Effect of pH on the reaction of MSZ with ARS, (30 $\mu\text{g/mL}$ of MSZ, 1.0 mL buffer solution, 1.0 mL of ARS 13×10^{-3} mol/L) at room temperature, and reaction time 10 min.

derivatization of MSZ to produce a chromophore absorbing more in the visible region was appropriate. MSZ contains a primary aliphatic amino group, which is suitable for derivatization by NQS, an analytical chromogenic reagent for the determination of primary and secondary amines [27-35]. MSZ was found to react with NQS under the experimental conditions to form an orange colored product exhibiting λ_{\max} at 470 nm (Figure 4). Under the optimum reaction conditions, the absorbance was found to obey the Beer-Lambert law.

The effects of pH on the reaction of MSZ with (NQS) were examined by varying the pH from 4.0 to 13.0, the results revealed that the absorbance increased with increasing up to pH 12. The maximum readings were attained at pH value of 12.0. At pH value more than 12.0, a decrease in the reading occurred. This was attributed probably to the increase in the amount of hydroxide ion that holds back the reaction of MSZ with NQS [33] (Figure 5).

The effect of temperature on the reaction was also studied by varying the temperature from 25°C to 90°C. The highest absorbance is obtained at room temperature for 10 min.



The effect of different NQS concentrations on the absorbance of product was studied; it is evident that the absorbance increases with increasing NQS concentration and reached maximum on using 1 mL of 4×10^{-3} mol/L NQS. Therefore, this concentration was used in all subsequent work (Figure 6).

Stoichiometry of the reaction (Job's method)

Under the optimum conditions shown in Table 1, the stoichiometric ratio between MSZ and each of investigated reagents (ARS, NQS) was found to be 1:1 (Figures 7 and 8). Based on this ratio, the reaction pathways were postulated to be proceeded as shown in scheme 1 and scheme 2.

Validation of the proposed method

Linearity and sensitivity: Calibration curves for Methods A and B in the ranges 15-97.5 µg/mL and 2-22 µg/mL were linear with correlation coefficients (r^2) of 0.9986 and 0.9997 for methods A and B, respectively. The molar absorptivities (ϵ) at 600 nm and 470 nm for Methods A and B were 1.184×10^3 and 7.733×10^3 L/mole/cm, respectively. The sandell's sensitivity values were 0.129 and 0.020 for methods A and B respectively.

The limit of detection (LOD) is defined as the minimum level at which the analyte can be reliably detected for the two methods was calculated using the following equation

$$\text{LOD} = 3.3s/k$$



Where s is the standard deviation of replicate determination values under the same conditions as for the sample analysis in the absence of the analyte and k is the sensitivity, namely the slope of the calibration graph. In accordance with the formula, the detection limits were found to be 4.92 and 0.56 µg/mL for method A and B, respectively.

The limit of quantification (LOQ) is defined as the lowest concentration that can be measured with acceptable accuracy and precision

$$\text{LOQ} = 10s/k$$

According to this equation, the limit of quantification was found to be 14.89 and 1.69 µg/mL for Method A and B respectively; these parameters for the two methods are summarized in Table 2.

Accuracy and precision: The accuracy and precision of the






Figure 6: Effect of NQS concentration on the reaction with MSZ (20 µg/mL of MSZ, 1.0 mL buffer solution (pH 12.0), 1.0 mL of NQS, at room temperature (25°C) and reaction time 10 min.

Condition	Method A	Method B
pH	8	12
Buffer volume (ml)	2	1
Temperature (°C)	40	25
Reaction time (min)	10	10
Reagent concentration mole/L	15x10 ⁻³	4x10 ⁻³

Table 1: Optimum conditions for the reactions of two methods.




Figure 7: Job's method for the reaction of MSZ with ARS, 2.0 mL buffer solution (pH 8), at 40°C, and reaction time 10 min.






Figure 8: Job's method for the reaction of MSZ with NQS, 1.0 mL buffer solution (pH 12), at room temperature (25°C) and reaction time 10 min.



Parameters	Method A	Method B
Linear range(µg/mL)	15-97.5	2-22
limit of detection, LOD(µg/mL)	4.92	0.56
Limit of quantification, LOQ (µg/mL)	14.89	1.69
Slope	0.00705	0.0426
Intercept	0.01055	0.01944
Correlation coefficient(r)	0.99862	0.99971
Molar absorptivity ε (L mol ⁻¹ cm ⁻¹)	1.184x10 ³	7.733x10 ³
Sandell sensitivity	0.129	0.020

Table 2: Summary of quantitative parameters and statistical data using the proposed procedure.

Method	Concentration(µg/mL)	Recovery (% ± RSD)*	Relative error %
Method A	30	101.24 ± 0.890	-1.24
	60	97.27 ± 0.237	-2.73
	90	100.36 ± 0.309	-0.37
Method B	6	96.85 ± 1.87	-0.031
	12	102.81 ± 0.37	0.028
	18	99.23 ± 2.31	-0.037

*three determinations were used for the proposed methods

Table 3: Precision results for the proposed methods.

proposed method were determined at three concentration levels of MSZ (within the linear range) by analyzing three replicate analyses on pure drug of each concentration. The percentage relative error as accuracy and percentagerelative standard deviations (RSD) as precision for the results did not exceed 2.5% for the two methods as shown in Table 3, indicating the good reproducibility and repeatability of the two methods. This good level of precision and accuracy was suitable for quality control analysis of MSZ in their pharmaceutical formulation.

Robustness: Robustness was examined by evaluating the influence of small variation in the method variables on its analytical performance. In these experiments, one parameter was changed whereas the others were kept unchanged, and the recovery percentage was calculated each

Method A Condition	Recovery \pm RSD*	Method B Condition	Recovery \pm RSD*
pH 7.8 8.2	100.61 \pm 1.791 99.82 \pm 0.902	pH 11.8 12.2	101.62 \pm 1.326 102.01 \pm 1.762
Temperature ($^{\circ}$ C) 35 45	100.92 \pm 0.893 103.44 \pm 1.308	Temperature ($^{\circ}$ C) 20 30	104.83 \pm 2.790 102.28 \pm 1.968
Time (min) 8 12	98.72 \pm 0.456 99.19 \pm 0.454	Time (min) 8 12	103.34 \pm 1.305 103.37 \pm 1.735
ARS concentration (mole/L) 14.8x10 ⁻³ 15.2x10 ⁻³	102.66 \pm 1.318 98.09 \pm 0.459	ARS concentration (mole/L) 3.8x10 ⁻³ 4.2x10 ⁻³	100.84 \pm 1.336 98.49 \pm 2.278

*value are mean of three determinations

Table 4: Robustness of the proposed methods.

Method	Brand name and dosage form	Labeled claim 500 mg	Amount found	(% found \pm RSD)*
Method A	Pentasa	500	488.85	97.77 \pm 0.46
Method B	—	—	490.85	98.17 \pm 1.70

*Three determinations were used for the proposed method

Table 5: Determination of MSZ in tablets by the proposed methods.

time. It was found that small variation in the methods variables did not significantly affect the procedures. This provided an indication of the reliability of the proposed methods during routine work; recovery values were shown in Table 4.

Applications of the methods: The proposed methods were applied to the pharmaceutical formulation containing MSZ. The results are shown in Table 5. Indicate the high accuracy of the proposed methods for the determination of the studied drug. The proposed methods have the advantage of being virtually free from interferences by excipients. The percentages were 97.77 \pm 0.46 and 98.17 \pm 1.67 for method A and B, respectively (Table 5).

Conclusions

The development spectrophotometric methods for the determination of MSZ in pharmaceutical formulation are simple, sensitive, rapid and accurate. The methods are superior to the previously reported spectrophotometric methods [14-20] in terms of the simplicity and sensitivity. The methods are practical and valuable for routine application in quality control laboratories for analysis of MSZ.

References

- Actis GC, Pellicano R, Rizzetto M, Ayoubi M, Leone N, et al. (2009) Individually administered or co-prescribed thiopurines and mesalamines for inflammatory bowel disease. *World J Gastroenterol* 15: 1420-1426.
- Gonçalves E, Almeida LM, Dinis TC (1998) Antioxidant activity of 5-aminosalicylic acid against peroxidation of phosphatidylcholine liposomes in the presence of alpha-tocopherol: a synergistic interaction? *Free Radic Res* 29: 53-66.
- Sandborn WJ, Feagan BG, Lichtenstein GR (2007) Medical management of mild to moderate Crohn's disease: evidence-based treatment algorithms for induction and maintenance of remission. *Aliment Pharmacol Ther* 26: 987-1003.
- Palumbo G, Carlucci G, Mazzeo P (1995) Simultaneous determination of 5-aminosalicylic acid, acetyl-5-aminosalicylic acid and 2,5-dihydroxybenzoic acid in endoscopy intestinal biopsy samples in humans by high-performance liquid chromatography with electrochemical detection. *J Pharm Biomed Anal* 14: 175-180.
- Zadeh HA, Kohansal S (2012) Determination of mesalazine by spectrofluorometry in human serum after solid-phase extraction with Ni-Al layered double hydroxide as a nanosorbent. *J Braz Chem Soc* 23: 473-481.
- Gotti R, Pomponio R, Bertucci C, Cavrini V (2001) Determination of 5-aminosalicylic acid related impurities by micellar electrokinetic chromatography with an ion-pair reagent. *J Chromatogr A* 916: 175-183.
- Nigovic B, Simunic B (2003) Determination of 5-aminosalicylic acid in pharmaceutical formulation by differential pulse voltammetry. *J Pharm Biomed Anal* 31: 169-174.
- Haney PW, Dash AK (1997) Simple liquid chromatographic method for the analysis of 5-aminosalicylic acid and its degradation product. *J Chromatogr A* 765: 233-239.
- Palumbo G, Bacchi S, Primavera L, Palumbo P, Carlucci G (2005) A validated HPLC method with electrochemical detection for simultaneous assay of 5-aminosalicylic acid and its metabolite in human plasma. *Biomed Chromatogr* 19: 350-354.
- Hussain FN, Ajjan RA, Moustafa M, Anderson JC, Riley SA (1998) Simple method for the determination of 5-aminosalicylic and N-acetyl-5-aminosalicylic acid in rectal tissue biopsies. *J Chromatogr B Biomed Sci Appl* 716: 257-266.
- Bystrowska B, Nowak J, Brandys J (2000) Validation of a LC method for the determination of 5-aminosalicylic acid and its metabolite in plasma and urine. *J Pharm Biomed Anal* 22: 341-347.
- Nobilis M, Vybiralova Z, Sladkova K, Lisa M, Holcapek M, et al. (2006) High-performance liquid-chromatographic determination of 5-aminosalicylic acid and its metabolites in blood plasma. *J Chromatogr A* 1119: 299-308.
- Pastorini E, Locatelli M, Simoni P, Roda G, Roda E, et al. (2008) Development and validation of a HPLC-ESI-MS/MS method for the determination of 5-aminosalicylic acid and its major metabolite N-acetyl-5-aminosalicylic acid in human plasma. *J Chromatogr B Analyt Technol Biomed Life Sci* 872: 99-106.
- Achariya SK, Sahu A, Das S, Sagar P, Annapurma MM (2010) Development and Validation of Spectrophotometric methods for the Estimation of Mesalazine in Pharmaceutical Preparations. *J Pharm Educ Res* 11: 63-67.
- Darak V, Karadi AB, Arshad MD, Appalraju S (2011) Derivative Spectroscopic Determination of Mesalazine in Tablets dosage forms. *Pharma science monitor* 2: 31-35.
- Madhavi V, Panchakshari V, Prathyusha TN, Sekaran CB (2011) Spectrophotometric Determination of Mesalazine in Bulk and Tablet dosage forms based on Diaz-coupling reaction with Resorcinol. *International journal of pharmaceutical sciences review and research* 105-109.
- Moharana AK, Banerjee M, Panda S, Muduli JN (2011) Development and validation of UV spectrophotometric method for the determination of Mesalazine in bulk and tablet formulation. *International Journal of Pharmacy and Pharmaceutical Sciences* 3:19-21.
- Rohitas M, Agrawal A, Jain AK, Lariya NK, Kharya AK, et al. (2010) Development of Simultaneous Spectrophotometric Method of Mesalazine and Prednisolone in Same Dosage Form. *Int J Appl Pharm* 2: 8-11.
- Srikanth K, Emmanuel KA, Raju Rasayan RK (2010) Spectrophotometric determination of oxybutynin chloride through Ion association complex formation. *Rasayan J Chem* 3: 179-187.
- Basavaiah K, SriLatha, Swamy JM (1999) Spectrophotometric determination of ceterizine hydrochloride with Alizarin Red S. *Talanta* 50: 887-892.
- Job P (1964) *Advanced Physicochemical Experiments*, 2nd Ed., Edinburgh: Oliver and Boyd 54
- Kishore M, Rao YH, Janardhan M (2010) Rapid determination of Ziprasidone and Buclizine Hydrochloride in pharmaceutical formulations (Tablets) by simple spectrophotometric method. *International Journal of Pharma Sciences and Research* 1: 438-444.
- Farhadi K, Savojbolaghi AK, Maleki R (2003) Study of Reactions of Triiodide and Alizarin Red S with Some Important Phenothiazines. Development of an Indirect Titrimetric and a Spectrophotometric Method for the Assay of Phenothiazine Derivatives. *Journal of the Chinese Chemical Society* 50: 153-159.
- Gouda AA, E Sheikh R, El-Azzazy RM (2012) Charge Transfer Spectrophotometric Determination of Zolmitriptan in Pure and Dosage Forms. *Analytical and Bioanalytical Techniques* 3: 6.
- Hassan WS, E-Henawee MM, Gouda AA (2008) Spectrophotometric determination of some histamine H1-antagonists drugs in their pharmaceutical preparations. *Spectrochim Acta A Mol Biomol Spectrosc* 69: 245-255.

-
26. Elbashir AA, Abdalla FAA (2013) Application of alizarin red S as an ion-pair reagent for the spectrophotometric determination of Olopatadine Hydrochloride in pharmaceutical formulation. *American Academic & Scholarly Research Journal* 5: 22-30.
 27. Wang H.Y, Xu LX, Xiao Y (2004) Spectrophotometric determination of ampicillin sodium in pharmaceutical products using sodium 1,2-naphthoquinone-4-sulfonic as the chromogenic reagent. *Spectrochim Acta Part A* 60: 2933–2939.
 28. Darwish IA (2005) Kinetic spectrophotometric methods for determination of trimetazidine dihydrochloride. *Anal Chim Acta* 551: 222–231.
 29. Ebraheem SAM, Elbashir AA, Aboul-Enein HY (2011) Spectrophotometric methods for the determination of gemifloxacin in pharmaceutical formulations. *Acta Pharmaceutica Sinica B* 4: 248–253.
 30. Elbashir AA, Elwagee HEA (2012) Spectrophotometric determination of pyrimethamine (PYM) in pharmaceutical formulation using 1,2-naphthoquinone-4-sulfonate (NQS). *JAAUBAS* 11: 32–36.
 31. Elbashir AA, Ahmed SM, Aboul-Enein HY (2012) New spectrofluorimetric method for determination of cephalosporins in pharmaceutical formulations. *J Fluoresc* 22: 857-864.
 32. Ali Ahmed SM, Elbashir AA, Suliman FE, Aboul-Enein HY (2013) New spectrofluorimetric method for determination of cephalosporins in pharmaceutical formulations. *Luminescence* 28: 734-741.
 33. Ali Ahmed SM, Elbashir AA, Aboul-Enein HY (2011) New spectrophotometric method for determination of cephalosporins in pharmaceutical formulations. *Arab. J Chem* 10.1016/j.arabjc.2011.08.012.
 34. Elbashir AA, Ahmed AA, Ali Ahmed SM, Aboul-Enein HY (2012) 1,2-Naphthoquinone-4-Sulphonic Acid Sodium Salt (NQS) as an Analytical Reagent for the Determination of Pharmaceutical Amine by Spectrophotometry. *Appl. Spectrosc. Reviews* 47: 219–232.
 35. Elbashir AA, Awad SF (2013) A New Spectrophotometric Method for Determination of Penicillamine in Pharmaceutical Formulation Using 1, 2-naphthoquinone-4-sulfonate (NQS). *J Pharmacovigilance* 1:2.