**Open Access** 

# Directional Solar Asymmetries in Sound Pyrenean Catalan Horses

#### Parés Casanova Pere M<sup>\*</sup> and López-Navarro

Department of Animal Sciences, University of Lleida, Alcalde Rovira Roure, Lleida, Catalonia, Spain

#### Abstract

Most published attempts to quantify equine hoof form are based on lineal and angular measurements. Here we applied geometric morphometric methods to study shape variation of hoof outlines in a sample of 25 distal forelimbs (13 right and 12 left). Limbs belonged to sound "Cavall Pirinenc Català" horse, an equine meat local breed from Pyrenees, slaughtered in an abattoir. The outline of each hoof was represented by a set of two landmarks and 86 semi-landmarks. Results reflected directional asymmetric, e.g. consistent differences between medial and lateral contours for all four limbs. Right limbs tended to supinate (rotation of medial wall towards out), while left limbs tended to pronate (rotation of lateral wall towards in). Such morphological adjustments may be an important consideration for hoof practitioners and may imply an important reconsideration of "normal" feet evaluation.

Keywords: Equine • Hoof conformation • Forelimb • Laterality • Sidedness

### Introduction

Bilateral symmetry among animals is rarely perfect, i.e., when measurements of left and right structural parts do not always are perfectly equal [1,2]. In a population, bilateral asymmetry can be observed to occur in three general patterns, *viz*. fluctuating asymmetry (FA), directional asymmetry (DA) and anti-symmetry (AS). FA represents a random variation on the right and left side; DA involves left and right sides differences but *in the same direction*; in AS can be considered a macroscopic form of FA [3-6].

Geometric morphometrics (GM) is based on the Cartesian coordinates of landmarks (measurement points) that are homologous across all measured individuals [7]. The set of all landmarks preserves the geometry of the studied configurations, so GM is of superior statistical power than most traditional morphometric approaches and is particularly effective for exploratory studies [7].

Landmark configurations need to be registered (superimposed) prior to any statistical analysis because the coordinates not only contain information on the shape of the measured objects, but also on their position, scale, and orientation [8]. The most common superimposition technique in GM is Generalized Procrustes Analysis, consisting of three steps: (i) translation to have the same centroid (average landmark position), (ii) scaling to have the same size, and (iii) iteratively rotation to minimize the summed squared distances between the landmarks and the corresponding sample average [7]. Overall size is measured as centroid size, the square root of the summed squared distances between the landmarks and their centroid [8]. The coordinates of the superimposed landmark configurations are called Procrustes shape coordinates as they contain information about the shape of the landmark configurations only [8].

Curvatures express the rate of change of the angle between the tangent to the curve and the x-axis [9]. Curvatures cannot be well described using visual or traditional metric methods [9]. Semi-landmarks are points along curvatures. They are initially placed at approximately corresponding positions and their exact locations are ulteriorly estimated statistically in order to create geometrically homologous points that can be used in the subsequent analysis as if they were anatomical landmarks [9].

Many biological structures, such as hoof outlines, consist of relatively smooth curves and lack homologous landmark points that can be identified in all individuals. Object symmetry for hooves can be considered if the axis of symmetry is included within the structure itself as the axial plan and separates lateral and medial sides as "halves" [10,11]. The convenient statistical properties of GM together with its effective visualization allow for a potentially very powerful exploratory of the hoof solar structure [12]. Separation of DA and FA is of great importance to the shape analysis of symmetric structures, as it is a key step that allows for distinction between systematic asymmetry, which is genetically fixed and heritable for DA, and a sign of developmental instability for DA [6].

Handedness (or laterality, as it is referred to when applied to horses) means they prefer either the right or left side. Recent researchers have found that horses have a preferred side of the body, just like humans [13,14]. In the present paper we apply GM methods to study object shape asymmetry of the entire equine hoof outline. We extracted the hoof shape from surface scans of the sole (see the methods section below) using landmarks and semilandmarks. To the best of our knowledge, symmetries of equine hoof outline shapes using GM have scarcely been studied [15].

## **Materials and Methods**

#### **Studied population**

The Catalan Pyrenean horse ("Cavall Pirinenc Català") is a compact, broad-built horse with a small population (<4,600) which is located in the NE part of the Pyrenees, along the Catalan-French border [16]. Mainly bred for

\*Address for Correspondence: Parés Casanova Pere M, Department of Animal Sciences, University of Lleida, Alcalde Rovira Roure, Lleida, Catalonia, Spain, Tel: 34973706460; E-mail: peremiquelp@ca.udl.cat

Copyright: © 2020 Parés Casanova Pere M, et al. This is an open-access article distributed under the terms of the creative commons attribution license which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Received: 13 March 2020; Accepted: 10 July 2020; Published: 17 July 2020

meat production, it is reared outdoors throughout the year, normally not receiving additional food beside some low-quality straw in winter except when foals are selected for slaughter [17]. Then, they are gathered in paddocks and receive additional feeding with hay and concentrates during the last months before slaughter, at about 10-12 months of age ("poltres", body mass about 350 kg) [17]. Animals of this breed do not receive any hoof care, trimming, or shoeing; therefore, their hooves must be considered naturally shaped. Anyway, hoof problems are rather rarely encountered, being the forelimb "splay foot" -the hoof wall flaring outwards- the most frequently found non-functional abnormality (pers. obs.).

#### **Sampled limbs**

At a commercial abattoir, 25 distal forelimbs (13 right and 12 left) were obtained from Catalan Pyrenean yearlings (<12 months) after slaughter. Sex and exact age were not registered. Animals were apparently healthy and sound. At the abattoir, limbs were disarticulated at the level of the carpus and were rinsed with water to clarify hoof outline. Ground surface was taken outlining the round edges with a pen on a paper. The original drawings were subsequently photocopied.

#### **Extraction of shape**

The outline of the hoof was digitized by 86 semi-landmarks and two true landmarks at the two axial-most positions (Figure 1). We used the sliding landmark algorithm to estimate the position of the semi-landmarks in all individuals, enabling the joint analysis of anatomical landmarks and curves (represented by semi-landmarks) [18]. All landmark configurations were superimposed by a Generalized Procrustes Analysis, standardizing for position, size, and orientation of the configurations. The resulting Procrustes shape coordinates were used for further statistical analysis [7]. The most common algorithm for this purpose is the sliding landmark algorithm, which iteratively slides the semi-landmarks along their curves in order to minimize local shape differences (the bending energy of the thin-plate spline





The software TpsUtil v. 1.50 [19] was used to prepare and organize the images. Landmarks were digitized twice, using TpsDig v. 1.40 [19], by one of the authors (Noelia) in two different sessions. In order to compare Procrustes to tangent space distances between individuals, the procedure performed using TpsSmall v. 1.33 [19] reflected a high degree of approximation of

shapes in the sample (i.e., shape space) in relation to the reference shape (i.e., tangent space) (r=0.999), which allows accurate capture of the nature and extent of skull shape deformations in subsequent statistical analyses.

A regression of centroid size versus shape (regression scores) was done to verify if allometry existed. Size was computed as centroid size (or, the square root of the sum of squared distances from the landmarks to their centroid) [7]. In the absence of allometry, it is the only size measure uncorrelated with all the shape variables [7].

For shape, a Procrustes ANOVA indicated the degrees of freedom, means of squares, F and P values for the effects from individuals, sides, Individual × sides effect and measurement error. The individual's effect denoted the individual variations of shape and size of hooves individual animals. The main effect of sides indicated the variation between sides and considered as the measure of DA. The Individual × sides is the mixed effect, which indicates there is failure of hoof shape differences to be on the same side in every case and therefore indicates there FA in the data. Lastly, measurement error is a gauge of the possible effect of measurement error on the estimates of FA [20].

#### Data analysis

All morphometric analysis were analysed in MorphoJ v. 1.06c [21] and PAST v. 2.17c softwares [22]. Confidence level was stablished at 95%.

### Results

#### Allometry

No significate regression of centroid size versus shape appeared, with only a 1.32% of shape variation explained by size variation (p=0.200).

#### Hoof shape symmetries

For hoof shape, Procrustes ANOVA highly significant variations in symmetry within individuals and sides (DA), but not for side\*individual interaction (FA) (Table 1). FA presented mean squares about 1.6 to 7.2 times lower than for DA, which accounted for around a 25% of total asymmetric variation. FA was significative for left forelimb. Right limbs tended to supinate (rotation of medial wall towards out), while left limbs tended to pronate (rotation of lateral wall towards in).

Table 1: Procrustes ANOVA of hooves shape in object symmetry for 25 distal forelimbs (13 right and 12 left) belonging to sound "Cavall Pirinenc Català" horse. Mean squares (MS) are the amount of variation from the one higher level in the hierarchy. The F value represents the comparison of each MS to the one lower level of MS which could be the source of error. Shape differences among individual hooves ("Individual" effect) were strongly significant (p<.0001) and were approximately about 0.4 to 1.2 times larger than the asymmetry between both sides of the hooves. DA ("Sides" effect) proved to be highly significant and considerably larger than the FA ("Individual × sides" interaction effect), which was very subtle for left forelimb. Error is a gauge of the possible effect of measurement error on the estimates of FA.

1/ Right forelimb (n=13)							
Effect	SS	MS	df	F	Ρ		
Individual	0.09501	0.0000920673	1032	3.41	<.0001		
Sides	0.00428	0.0000497111	86	1.84	<.0001		
Individual × sides	0.02787	0.0000270028	1032	1.19	0.0006		
Error	0.05093	0.0000227758	2236				

2/ Left forelimb (n=12)							
Effect	SS	MS	Df	F	Ρ		
Individual	0.08183	0.0000865028	946	3.19	<.0001		
Sides	0.00387	0.0000449869	86	1.66	0.0003		
Individual × sides	0.02563	0.0000270950	946	1.07	0.0998		
Error	0.05212	0.0000252497	2064				

# Discussion

This study aim was to assess hoof asymmetries on solar surface in a local equine breed maintained under extensive management and consequently, which has no care of feet, so conclusions may express horse natural wearing. We applied geometric morphometric methods to study shape asymmetries for each forelimb separately with an outline represented by a set of landmarks and semi-landmarks.

Hooves are not pieces of rigid tissue that forever hold the shape. They are malleable enough to change shape when put under pressure, and such forces would not act uniformly [23]. The medial side of the hoof bears most of the descending weight of the horse [24]. In response to this, the 'plastic' hoof develops a medial side that is steeper and a lateral side that is more oblique, so it changes shape. Hoof DA in the studied sample is largely present, and attributable to differential mechanical loading, e.g. laterality. Laterality represents a dominance. Limb dominance is a common occurrence in many species [25], including horse [14,26]. Like humans [27] most horses are right forelimb dominant [14,28,29] and this would explain the detected subtle FA only in left forelimb, which would represents a very low disturbing effect. As asymmetric shape variation of hoof surface was not determined by allometric changes relating to size, we suppose that FA is due to a function process rather than to a developmental (e.g. wall growth) effect.

### Conclusion

"Normal hooves" in horses are therefore directionally asymmetrical but, notably, with a clear right supination and a left pronation. These results would be considered by both farriers and veterinarians as an important feature of hoof conformation. Now it would be interesting to study if this unevenness coincides with both kinetic and kinematic asymmetrical pressure differences.

## **Conflicts of Interest**

The authors declare no conflicts of interest. The post-mortem use of this kind of non-edible parts did not require an approval from the Ethics Committee.

# Acknowledgements

We thank MAFRISEU SA for providing access to the legs and the individual information for each animal. We are also grateful to anonymous reviewers for their insightful recommendations.

# **Authors' Contributions**

PMPC designed the study and directed implementation and data collection. NLN collected data. Both authors analyzed the data and drafted the manuscript.

# **Supporting Information**

The contents of all supporting data are the sole responsibility of the authors.

### References

- 1. Valen, Van. "A study of fluctuating asymmetry." *Evolution* (1962): 125-142.
- Klingenberg, Christian Peter, Marta Barluenga, and Axel Meyer. "Shape analysis of symmetric structures: quantifying variation among individuals and asymmetry." *Evolution* 56 (2002): 1909-1920.
- Auffray, Jean-Christophe, Vincent Debat, and P. Alibert. "Shape asymmetry and developmental stability." On growth and form: spatio-temporal pattern formation in biology. *John Wiley and Sons Ltd, NY, USA* 1 (1999): 309-324
- Tomkins, Joseph L, and Kotiaho, JS. "Fluctuating Asymmetry." In Encyclopedia of Life Sciences (2001): 1-5.
- Mancini, Sandra, Sharon L. Sally, and Rick Gurnsey. "Detection of symmetry and anti-symmetry." *Vision Research*. 45 (2005): 2145-2160.
- De Coster, Greet, Stefan Van Dongen, Phillista Malaki, and Muchane, M, et al. "Fluctuating Asymmetry and Environmental Stress: Understanding the Role of Trait History." *PLoS One* 8 (2013): 1-9.
- 7. Adams, Dean C., James Rohlf, and Dennis E Slice. "A field comes of age: geometric morphometrics in the 21st century." Hystrix, 24 (2013): 7-14.
- Rohlf, F James. "Geometric morphometrics simplified." *Trends Ecol Evol* 20 (2005): 13-14.
- Gunz, Philipp, and Philipp Mitteroecker. "Semilandmarks: A method for quantifying curves and surfaces." *Hystrix* 24 (2013): 103-109.
- Willmore, Katherine Elizabeth, Christian Peter Klingenberg, and Benedikt Hallgrímsson. "The Relationship Between Fluctuating Asymmetry and Environmental Variance in Rhesus Macaque Skulls." *Evolution* 59 (2005): 898-909.
- Ducos, Monaliza B, and Sharon RM Tabugo. "Fluctuating asymmetry as an indicator of ecological stress and developmental instability of Gafrarium tumidum (ribbed venus clam) from Maak and Lagoon Camiguin Island, Philippines." *Int J Bioflux Society* 7 (2014): 516-523.
- Richtsmeier, Joan T, Valerie Burke Deleon, and Subhash R Lele. "The promise of geometric morphometrics." *Yearbook of Physical Anthropology* 45 (2002): 63-91.
- Hobbs, Sarah Jane, Sandra Nauwelaerts, Jonathan Sinclair, and Clayton, HM, et al. "Sagittal plane fore hoof unevenness is associated with fore and hindlimb asymmetrical force vectors in the sagittal and frontal planes." *PLoS ONE*, 13 (2018), 1-17.
- A, Byström, Clayton HM, Hernlund E, Rhodin M, and Egenvall A. "Equestrian and biomechanical perspectives on laterality in the horse." *Comparat Exercise Physiol* 16 (2020): 35-45.
- Roland, Elizabeth, Susan M Stover, Maury L Hull, and Katie Dorsch. "Geometric symmetry of the solar surface of hooves of thoroughbred racehorses." *American J Vet Res* 64 (2003): 1030-1039.
- 16. Infante, Gil. "Caracterización y gestión de los recursos genéticos de la población equina de carne del Pirineo Catalán (Cavall Pirinenc Català): interrelacion con otras razas cárnicas españolas." Universitat Autònoma de Barcelona. (2011).
- 17. Pere-Miquel, Parés. A Nonlinear Model for Estimating Hoof Surface Area in Unshod Meat-type Horses. *J Equine Vet Sci* 31 (2011): 379-382.
- MA, Webster, and David Sheets. "A practical introduction to landmark-based geometric morphometrics." *Paleontol Soc Papers* 16 (2010): 163-188.
- 19. Rohlf, F James. "The tps series of software." Hystrix 26 (2015): 9-12.

Page 3 of 4

- Fruciano, Carmelo. "Measurement error in geometric morphometrics." Develop Genes Evolut 226 (2016): 139-158.
- Klingenberg, Christian Peter. "MorphoJ: an integrated software package for geometric morphometrics." *Molecul Ecol Resource* 11 (2011): 353-357.
- 22. Hammer, Øyvind, Harper DT, and Ryan PD. "PAST-palaeontological statistics, ver. 1.89." *Palaeontol Electron* 4 (2001): 1-9.
- Dyson SJ, Tranquille CA, Collins SN and Parkin TD. "An investigation of the relationships between angles and shapes of the hoof capsule and the distal phalanx." *Equine Vet J* 43 (2011): 295-301.
- 24. P. Healey. Evaluating for equilibrium of the equine foot.
- Hackert, Rémi, Maes Marc Herbin, Libourel PA and Abourachid A "Limb preference in the gallop of dogs and the half-bound of pikas on flat ground." *Laterality* 13 (2008): 310-319.
- Leśniak, Kirsty, Lisa Whittington, Stephanie Mapletoft, and Jennifer Mitchell, et al. "The influence of body mass and height on equine hoof conformation and symmetry." *J Equine Vet Sci* 77 (2019): 43-49.

- Jandová, Soňa. "Laterality of lower limb and plantar pressure symmetry while walking in young adults." J Human Sport Exercise 14 (2019): 834-840.
- Larose, Claire, Marie-Annick Richard-Yris, Martine Hausberger, and Lesley J Rogers. "Laterality of horses associated with emotionality in novel situations." *Laterality* 11 (2006): 355-367.
- NP, Austin, and Rogers LJ. "Limb preferences and lateralization of aggression, reactivity and vigilance in feral horses, Equus caballus." *Anim Behav* 83 (2012): 239-247.

How to cite this article: Parés Casanova, Pere M, and López-Navarro. "Directional Solar Asymmetries in Sound Pyrenean Catalan Horses". J Vet Sci Technol 11 (2020) doi: 10.37421/jvst.2020.11.596