

Disorganized Speech in Autism Spectrum Disorder: Part of the Problems in Social Pragmatic Communication and Verbal Intelligence

Sakinah Idris^{1,2*}, Louwrese A^{1,3}, Van der Ende J¹, Hillegers M¹, Van Haren N¹ and Greaves Lord K^{1,3,4,5}

¹Department of Child and Adolescent Psychiatry/Psychology, Erasmus Medical Center-Sophia Children's Hospital, Rotterdam, The Netherlands

²Department of Psychiatry, Clinical Psychology Unit, Universiti Teknologi MARA, Malaysia

³Department of Medicine, Yulius Academie and Yulius Autism, Dordrecht, The Netherlands

⁴Department of (Youth) Mental Health and Autism of Lentis Psychiatric Institute, Autism Team Northern-Netherlands of Jonx, Groningen, The Netherlands

⁵Department of Psychology, Clinical Psychology and Experimental Psychopathology Unit, University of Groningen, Groningen, The Netherlands

Abstract

Background and aims: Disorganized speech (DS) refers to the failure to communicate coherently. Children with autism spectrum disorder (ASD) show disorganized speech, such as abrupt topic changes. Disorganized speech can be explained by 1 social pragmatic communication difficulties (SPCD) or 2 social cognition problems (SCP).

Methods: We tested whether children with ASD and severe disorganized speech differed from those with mild or no disorganized speech, using parent's report on SPCD and tested SCP (i.e., Theory of Mind; ToM). A total of 62 children with ASD (aged 7-12 years) were included, 12 with severe disorganized speech, 32 with mild, and 18 with no disorganized speech. The disorganized speech was assessed using the Kiddie Formal Thought Disorder Rating Scale (KFTDS). SPCD was measured using the parent reported Children's Communication Checklist (CCC). SCP was measured using a first order false belief task.

Results: Children with severely disorganized speech showed significantly more inappropriate initiations and less coherence. No significant group differences were found in the false belief task, even after included verbal IQ as a covariate in multivariate analysis.

Conclusion: These findings may contribute to the conceptualization of autistic communication difficulties which may guide intervention optimizations. Implications: Furthermore, we may consider offering children with ASD and disorganized speech interventions that stimulate and train broader social pragmatic communication skills, social cognition, and verbal ability (i.e., conversational skills).

Keywords: Autism spectrum disorder • Disorganized speech • Social pragmatic communication • Social cognition • Verbal intelligence

Introduction

Disorganized speech refers to the degree to which transmission of meaning fails because of incoherent communication, such as abrupt topic changes. Disorganized speech entails verbal signs of illogical thinking and loose associations [1,2]. Studies reported elevated levels of disorganized speech in children with autism spectrum disorder (ASD) as compared to typically developing children [3-5]. Disorganized speech in ASD has formerly been explained as a manifestation of social cognitive problems and as a potential precursor of psychosis [2,6,7]. However, a previous study suggested that disorganized speech may rather be a part of the broader developmental difficulties in social pragmatic communication (SPC) [8]. In other words, learning how to communicate logically and coherently to meet the listener's needs (how to attune one's communication to the social context) is a

developmental process that might be delayed or altered in autistic children with ASD.

Disorganized speech as a part of SPCD

Coherent communication in typically developing (TD) children can be explained in terms of developmental aspects of children's disorganized speech as described by [9]. Before the age of 7, typically developing children may still have difficulty maintaining a topic of discourse. For instance, instead of providing only relevant topic information, they often still make abrupt topic changes. This behavior can be regarded as displaying signs of disorganized speech which is referred to as loose associations (LA) and illogical thinking (ILL). Around the age of 7, these large unannounced topic shifts (LA's) tend to diminish, but smaller unannounced topic shifts (ILL's) may still appear. By the age of 9, most typically developing children have learned to organize their speech and barely display any signs of disorganized speech.

In contrast, at the age of 9, children with ASD have difficulty maintaining an ongoing topic of conversation, which is a signal of developmental delay in social pragmatic communication. As they get older, their development of social pragmatic communication is improved [3,10]. Based on the work by [11], we suggest that disorganized speech reflects the difficulties that children with ASD display in maintaining coherent communication with others. These difficulties are a signal of developmental delay in the development of social pragmatic communication skills, i.e., engaging in reciprocal social communication in such a way that the listener's needs are met.

***Address for Correspondence:** Sakinah Idris, Department of Child and Adolescent Psychiatry/Psychology, Erasmus Medical Center-Sophia Children's Hospital, Rotterdam, The Netherlands, Email: sakinahpsyc@gmail.com

Copyright: © 2022 Idris Sakinah et al. This is an open-access article distributed under the terms of the creative commons attribution license which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Received: 02-March-2022, Manuscript No. jnd-22-56660; **Editor assigned:** 04-March-2022, PreQC No. P-56660 (PQ); **Reviewed:** 18-March-2022; QC No. Q-56660; **Revised:** 23-March-2022, Manuscript No. R-56660; **Published:** 30-March-2022, **DOI:** 10.4172/2329-6895.10.3.484

Are social cognition problems associated with disorganized speech?

In a study by [12], typically developing children and children with ASD were tested in experiments on false beliefs which is an index for Theory of Mind (ToM). The result revealed a strong developmental age trend: 57% of the children aged 4 to 5 years old passed the (first order) ToM task, while almost all 6 to 9 years old children (86%) passed this test. In contrast, children with ASD aged from 10:11 to 15:10 years passed the first order false belief task. Clearly, for children with ASD, the first order mental state attribution is acquired much later [13,14]. Thus, this study is important for taking the developmental perspectives on the theory of mind in autism.

This has led researchers to believe that individuals with ASD may have a specific developmental delay in ToM that can also cause problems with the acquisition of social pragmatic language [15-17]. We hypothesize that this delay in ToM may be related to the significant problems in children's ability to maintain coherent communication, as taking on the perspective of others is needed to understand their communication needs. We based this hypothesis on the assumption of Hale and Tager Flusberg (2005) that SPCD is the consequence of deficits in ToM [11] found that children with ASD (aged 4-13 years) made significant gains over 1 year in their ability to maintain a topic of conversation that was related to ToM. A significant association was found between understanding the first order false belief tasks and the ability to maintain a conversation in a meaningful way among children with ASD [18,19].

Our study aims to explore whether disorganized speech (1) is part of the broader social pragmatic communication difficulties (SPCD) that appear in ASD, and (2) can be explained by underlying social cognition problems (SCP). To test this, we examined (a) whether children in different disorganized speech groups differed regarding their social pragmatic communication skills as reported by their parents and (b) whether the groups differed on ToM performance. We hypothesize that children who have problems in organizing their speech have difficulties to maintain a topic of conversation and have poor social cognitive abilities, i.e., ToM [8].

Methods

Participants and procedure

One hundred 42 children with ASD participated in this study [8,20] of which 62 were selected for analysis in this article (Figure 1). They visited the outpatient clinic at the Department of Child and Adolescent Psychiatry/ Psychology of Erasmus Medical Center-Sophia's Children Hospital Rotterdam, Netherlands between July 2002 to September 2004. Inclusion criteria were: (1) meeting the clinical diagnosis of Diagnostic and Statistical Manual of Mental Disorders (DSM-IV-TR; American Psychiatric Association, 2000) classification of ASD and (2) parents being able to communicate in the Dutch language. The exclusion criteria in this study were (1) children aged below 7 years old because cut off points of ILL and LA were not available for children younger than 7 years, (2) the presence of severe neurological or physical problems, such as epilepsy or muteness. The total group (n=62) children had a mean age of 9.41 years (SD=1.65 years). 84% of the participants were boys (n=52). All children with complete data on Kiddie Formal Thought Disorder Scale (KFTDS), the Children's Communication Checklist (CCC), and the first order ToM task were included in the analyses.

The clinical diagnostic assessment was obtained by a multi-disciplinary team that consisted of a parental interview on the child's early developmental history, medical history, and current functioning. Parents of the participating children additionally filled out a questionnaire, e.g., the CCC. The children also underwent psychological assessments e.g., the KFTDS and the ToM task [21]. Parents of the participants had signed informed consent forms.

The Medical Ethics Committee of the Erasmus Medical Centre approved this study.




Figure 1. Sample description.

Materials

Disorganized speech

Kiddie Formal Thought Disorder Rating Scale (KFTDS) was used to assess disorganized speech, quantified as the presence of loose associations (LA) and illogical thinking (ILL) [6,9]. The validity of these KFTDS scales has been established in children with ASD [4,22]. Children were asked to answer standard questions after listening to two audio taped stories (i.e., "What did you like about this story?" or "Do you think this a true story?"). Children were then asked to make up his/her own story about one of four given topics (the incredible hulk, a witch, a disobedient child, or an unhappy child). This test took 20 to 30 minutes, and the child's speech was audiotaped. All stories were scored according to the KFTDS guidelines [6] by summing frequency counts for illogical thinking and loose associations. Total raw scores were corrected for the variability of speech by converting the number of utterances per minute. These continuous KFTDS scores were divided into different ordinal categories as falling above or below the cut off points, in which a score above the cut-off point indicates a higher likelihood of pathology. Using cut-off points, continuous KFTDS scores become dichotomous as falling above or below the cut-off point. Cut-off points were not calculated for children younger than 7 years old because KFTDS is a reliable measure of disorganized speech in children ages 7-18 years old [6,9].

The inter rater reliability for the total KFTDS score was good, with a kappa of 0.77 [6]. The kappa for ILL was 0.78 and for LA 0.71 [6]. The validity of the KFTDS has been established in children with schizophrenia spectrum disorders and ASD [9].

In our sample of children aged 7-12 (n=62), 71% of children scored above the cut-off point for ILL (n=44), while 19.4% of children scored above the cut-off point for LA (n=50). Based on these results, in the next step, we combine these into one integrated/combined grouping variable. A score above the cut-off point indicated a higher likelihood of disorganized speech. By using the cut-off point value, continuous KFTDS scores were dichotomized as falling above or below the cut-off point. Children with a score above the cut-off point for ILL and LA were assigned to the group 'severe disorganized speech', children with a score above the cut-off point for ILL and below the cut-off point for LA were assigned to the group 'mild disorganized speech', and children below the cut-off point of ILL and LA were assigned to the group 'no disorganized speech'.

The rationale for dividing the children into three groups is based on the study by [8], who found that ILL predicted the severity of autistic symptoms in adolescence, while LA did not. Those findings indicated that children with ASD show ILL from early childhood onwards, while LA deteriorates when they reach the age of 7 years old. In our study, we included children aged between 7 and 12 years and found that 70% of the children scored above the cut-off point for ILL, while 76% of the children scored below the cut-off point for LA. Based on these results, we combined ILL and LA into one integrated/combined grouping variable, as having both LA and ILL is

reflective of the presence of more types of disorganized speech than only having one or the other.

Social pragmatic communication difficulties

The Children's Communication Checklist-CCC [23] was administered to the parents of the included children. The CCC assessed pragmatic aspects of a child's social pragmatic communication difficulties. The questionnaire consists of 70 multiple choice questions on a 3-point scale (0=definitely applies, 1=applies somewhat, 2=does not apply). The lower the score on the CCC, the more impaired the child is. The questions are divided into nine subscales; five subscales for social pragmatic communication skills (coherence, inappropriate initiation, stereotyped language, use of context, and conversational rapport), two subscales to measure formal language (speech production and syntax), and the remaining two subscales assess characteristic behaviors of ASD (social relationships and interests). In this study, we used five social pragmatic communication skills subscales. There are 38 social pragmatic communication items in total, with the sum of the five subscales ranging between 88 to 162. The CCC was translated into Dutch using a two way translation procedure [24]. The internal consistency for the Pragmatic Composite Score was 0.76 [25]. Meanwhile, the internal consistency in our dataset is .80.

Social cognition problems

Theory of mind (ToM) was measured using a Dutch ToM task which was based on the classic Sally-Anne false belief task [13]. A child was told the following story: Ben places a bone in a barrel and goes out to play; while he is gone, Max (Ben's brother) takes the bone from the barrel and hides it in another barrel. After telling this story, children were asked a control question (Where is the bone really?). A wrong answer for this question represents an insufficient understanding of the story and understanding is necessary to perform well on the first order false belief tasks. One child who answered this question wrong was excluded from the analyses. Then, children were asked a false belief question (Which barrel will Ben choose?). The child was given a score of 1 for a correct answer and a score of 0 for the wrong answer.

Covariates

Individual factors were taken into consideration as possible confounding variables. First, sex may have influenced the result because females were usually more verbally oriented [26]. Despite both sexes have similar social understanding difficulties, males showed less restricted, repetitive patterns of behaviors [27]. Intelligence Quotient (IQ) was taken into account as a possible covariate based on [22], who found a significant negative correlation between Total IQ with illogical thinking and loose association on the KFTDS. A lower IQ is also associated with more difficulties in social pragmatic communication [28] and false-belief task [15]. In this study, IQ was measured using the Wechsler's Intelligence Scale for Children-Revised version (WISC-R). Finally, age was taken into account as a covariate because developmental aspects play a vital role in the manifestation of DS and therefore scores could be correlated with age. Age also affects social pragmatic communication [3,11] and ToM [12]. The older the child is, the better his/her social pragmatic communication skills are, both for TD children and children with ASD [10].

Statistical analyses

Of the 142 children recruited, 89 children had complete KFTDS data, while 53 children had incomplete KFTDS data. Six children were younger than 7 years and three children with high LA and low ILL scores were excluded from the analyses. The group of children with scores above the cut-off point of LA and below the cut-off point of ILL was too small to be included separately in the analyses. n=18 children who did not have complete data on the outcome measures, i.e., SPCD and SCP, were also excluded from the analyses (Figure 1).

The effects of potential attrition (sex, age, and IQ) were checked for the group with complete data (N=89) versus the group with incomplete data

(N=53). Then, the descriptive characteristics (sex, age, and IQ) of the children were provided for the total group (n=62) and compared between three subgroups i.e., children with severe (n=12), mild (n=32) and no DS (n=18).

Then, assumptions of normality, independence, and homogeneity of variance were checked for the CCC scores. Also, the assumptions of Fisher-Freeman-Halton Exact test were checked for the ToM data. Pearson correlations were used to estimate the influence of covariates (age and IQ) concerning the main variables (KFTDS groups, CCC and ToM). A Spearman correlation was performed to investigate the influence of sex to KFTDS groups, CCC and ToM. In case one of the covariates showed a significant correlation with both the predictor and the outcome variable, we included this covariate in the subsequent analysis of covariance (ANCOVA) and logistic regression analysis for ToM.

A one-way analysis of variance (ANOVA) was used to explore between-group differences for CCC. We examined whether the groups differed significantly on the subscales, i.e., inappropriate initiation, coherence, stereotyped language, use of content, and rapport (Aim 1). Post hoc pairwise tests were carried out to compare between two groups. In this study, we chose Games-Howell procedure as it remains accurate when sample sizes are unequal [29]. To examine whether disorganized speech influenced their first-order false-belief task (Aim 2), the study used a Fisher-Freeman-Halton Exact Test. We looked at the cross-tabulation and observed the number of children with disorganized speech pass the first-order false-belief task in each group.

In this study, a p<.05 (confidence interval of 95%) is considered to be significant. The statistical analyses have been performed using IBM SPSS Statistics 21.

Results

Descriptive

Sixty-two children with a mean age of 9.41 years, SD=1.65 years; Boys 84%; mean Total IQ 94.35, SD=19.47 were included in the analyses. An attrition analysis was performed. An independent sample t-test showed no statistically significant differences between children with complete data (n=62) and children with missing data (n=80) on age, sex, and intelligence (total, verbal, and performance).

Descriptive statistics were provided for the total group and the subgroups (i.e., children with severe, mild and no disorganized speech) as shown in Table 1. No differences were found for the distribution of sex between the three groups, F (2,59)=.006, p=.98. Age was also not significantly different between the groups, F (2,59)=2.15, p=.13. Significant differences on Total, Verbal, and Performance IQ were found amongst the groups (respectively F (2,59)=6.90, p<.01; F (2,59)=6.60, p<.01; F (2,59)=4.46, p=.02). Post hoc analyses were performed to check the pair-wise differences between groups (Table 1).

Table 1. Descriptive information of the total sample of (n=62) on covariates.

Pair-wise comparison							
	Severe DS (n=12)	Mild DS (n=32)	No DS (n=18)	Anova	No vs mild DS	No vs severe DS	Mild vs severe DS
Sex (% boys)	83	84	83	1.0			
	M (SD)	M (SD)	M (SD)		p	p	p
Age	8.59 (1.27)	9.50 (1.77)	9.81 (1.53)	.13			
IQ:							

- TIQ	77.92 (15.11)	96.28 (18.96)	101.89 (17.30)	<.01**	<.01	<.01	<.01
- VIQ	79.42 (14.04)	96.34 (19.03)	102.78 (16.72)	<.01**	.44	<.01	<.01
- PIQ	80.75 (19.43)	97.44 (20.73)	101.50 (17.00)	.02*	.74	.02	.05

Note: *p<.05; **p<.01

Normality plots of age, IQ, and CCC indicated a normal distribution of scores for these variables. The assumption of homogeneity of variance ($p > .05$) was satisfied for CCC-scores. The first assumption of the Chi-square test, the independence of participants, was confirmed. However, more than 20% of the group cells contained less than 5 participants. Therefore, the Fisher-Freeman-Halton Exact Test outcomes were interpreted [29].

Significant correlations between the dependent variables (CCC and ToM) and covariates are shown in Table 2. Therefore, in the main analyses (i.e., groups comparison regarding CCC subscales i.e., coherence and inappropriate initiation and ToM), we included IQ as a covariate in the models.

Table 2. Correlations between outcome variables and covariates.

	Coherence	Inapprop.	Stereotype	Context	Rapport	CCC	ToM
Sexb	.28*	.16	.09	.19	.12	.23	-.14

Table 3. Differences on CCC pragmatic subscales among disorganized speech groups.

	Severe DS (n = 12)	Mild DS (n = 32)	No DS (n = 18)				No vs mild DS	No vs severe DS	Mild vs severe DS
	M (SD)	M (SD)	M (SD)	F	p	Partial η^2	p	p	p
Pragmatic subscales:									
Coherence	25.83 (1.59)	28.69 (3.04)	29.44 (3.70)	5.46	< .01	.19	.74	<.01	<.01
Coherence corrected for TIQ				2.95	.06	.09	.88	.07	.11
Coherence corrected for VIQ				3.41	.04	.11	.87	.04	.08
Coherence corrected for PIQ				3.37	.04	.10	.86	.04	.09
Inappropriate initiation	22.83 (2.95)	26.13 (4.18)	26.5 (3.29)	4.14	.02	.14	.94	.01	.02
Inappropriate initiation corrected for TIQ				3.42	.04	.11	.98	.06	.05
Inappropriate initiation corrected for VIQ				4.61	.01	.14	.95	.02	.02
Inappropriate initiation corrected for PIQ				2.8	.07	.09	.99	.10	.09

Note: DS: disorganized speech

Due to positive significant differences found in CCC subscales i.e., coherence and inappropriate initiation, analysis of covariance (ANCOVA) were performed with the covariate i.e., the Total, Verbal, and Performance IQ. Only the subscale of 'Inappropriate initiation' shows significant differences between the groups after controlling for Total IQ. After correction for Verbal IQ, both subscales remained significant. However, after controlling for Performance IQ only 'Coherence' remained significant.

Agea	.46**	.28*	.19	.27*	.15	.39**	.19
TIQa	.33**	.10	.05	.10	.12	.19	.33**
VIQa	.28*	.01	-.06	.00	.15	.10	.37**
PIQa	.31*	.19	.17	.19	.07	.27*	.23

Note: *: $p < .05$; **: $p < .01$

Subscales of CCC; Inapprop: Inappropriate initiation; Stereotype: Stereotype language; Context: Use of context; CCC: Total Children's Communication Checklist; ToM: Theory of mind; TIQ: Total IQ; VIQ: Verbal IQ; PIQ: Performance IQ; a: Pearson correlations; b: Spearman correlations

Aim 1: Disorganized speech as a part of SPCD

Table 3 Analysis of variance showed an association between disorganized speech and social pragmatic communication difficulties as reported by the parents in CCC subscales i.e., coherence $F(2,59)=5.46, p < .01, \eta^2=.19$ and inappropriate initiation $F(2, 59)=4.14, p=.02, \eta^2=.14$ with both showing large effect sizes. Post hoc analyses showed that children with severe disorganized speech showed significantly lower scores on these subscales than children with no or mild disorganized speech. There was no significant effect of disorganized speech in other CCC subscales, i.e., stereotyped language, use of context, and rapport.

Aim 2: Are SCP associated with disorganized speech?

All children who took part in the study correctly answered the control question, which allowed us to conclude that they all knew (and implicitly believed) that the bone was put somewhere else after Ben had left. The cross-tabulation in Figure 2 shows the percentage of children passing/failing the first-order false-belief task for each disorganized speech group. 3.1% (1 out of 32) of the children in the mild disorganized speech group

and 5.6% (1 out of 18) of the children in the no disorganized speech group failed the first-order false belief task. By contrast, 25% (3 out of 12) of the children with severe disorganized speech failed the first-order false-belief task. The differences between the three disorganized speech groups were trend significant ($p=0.08$) (Figure 2).




Figure 2. Percentages of children who failed the first-order false-belief task

Table 2 shows that disorganized speech and Verbal IQ are correlated to scores on the first-order false belief task. A logistic regression analysis was conducted with disorganized speech groups as the predictor, passing/failing the ToM task as the outcome, and VIQ as a covariate. In a model with only disorganized speech as a predictor alone it was not significant ($X^2(1)=4.64$, $p=0.10$), and when VIQ was included as a covariate, disorganized speech was also not significantly predicted passing/failing the ToM task ($X^2=8.98$, $df=1$, $p=0.89$) (Table 4).

Table 4. Results of binary logistic regression of children who pass/fail ToM task.

	b (SE)	OR (95% CI)	p
Step 1 DS groups	-1.08 (.69)	0.34(0.09-1.31)	0.12
Step 2 DS groups	-0.12 (0.82)	0.89(0.18-4.43)	0.89
VIQ	0.11 (0.04)	1.11(1.02-1.21)	0.02*

Note: OR: Odd Ratio; CI: Confidence Interval; DS: Disorganized speech group; VIQ: Verbal IQ
* $p<.05$; ** $p<.01$

Discussion

The study aimed to explore whether disorganized speech is part of the broader social pragmatic communication difficulties (SPCD) in children with ASD and whether it might be explained by underlying difficulties in ToM. We hypothesized that disorganized speech is part of the broader social pragmatic communication difficulties seen in children with ASD and can be explained by a limited ability to tune one’s communication to the listeners’ needs, caused by social cognition problems, i.e., ToM [3,8,11,19,30,31]. Our results indicated children with profound disorganized speech (ILL+LA) showed more inappropriate initiations and less coherence. This small group ($n=12$) also showed more difficulties in ToM and cannot be explained by Verbal IQ.

Disorganized speech is part of social pragmatic communication difficulties (SPCD)

Our analyses revealed that disorganized speech was positively associated with several pragmatic subscales of the CCC, i.e., coherence and inappropriate initiation. Our findings extend previous studies that showed before that pragmatics or the appropriate use of language in social and communicative contexts are seriously impaired in children with ASD [8,32-

36].

According to the parents’ reports, children with profound disorganized speech showed more inappropriate initiations than children with mild or no disorganized speech. Children who demonstrate inappropriate initiation show impairments in reciprocal communication, no interest in others, and talk about subjects that are outside the interest of the listener [37,38]. Consequently, the listener is not able to follow their story. Also in another study, children with ASD were found to initiate less in their conversations with peers than the typically developing group [39]. In line with our hypothesis, severely disorganized speech, as tested with the KFTDS, was associated with more difficulties in social pragmatic communication among children with ASD. In our study, other aspects of social pragmatic communication skills did not show associations with disorganized speech, such as rapport. Rapport can be considered as mainly relying on nonverbal cues of social communication such as eye contact, facial expressions, or gestures. The KFTDS does not measure these nonverbal communication aspects. Therefore, these findings need to be confirmed by research using clinical observation as well as parent reports.

Are social cognition problems associated with disorganized speech?

Our second aim was to examine the differences in performance on the first-order false-belief task between groups. The hypothesis that poor ToM is associated with profound disorganized speech was not confirmed, as we only found a trend association if no corrections was made. Yet, after taking into consideration Verbal IQ as a covariate, groups of disorganized speech also showed not a significant predictor of performance on the ToM task. Children with severely disorganized speech showed more difficulties with social cognition as compared to children with mild or no disorganized speech, but this difference was mainly explained by their differences in Verbal IQ.

This can potentially be explained by the observation that some children with ASD seem to ‘imitate’ other people’s verbal behavior and therefore appear more verbally intelligent than they in fact are. In other words, their speech production appears better (i.e. abundant speech production with rich vocabulary) than their language perception is (i.e., in terms of semantics and syntaxis) which might alter the actual meaning of what is being said [40]. Underlying language difficulties can lead to poor performance on ToM tasks [40], which would explain the current findings.

Methodological considerations

The current study has several limitations that should be considered when interpreting the results. Firstly, a small and unequal number of children were included in each group of disorganized speech ($n=12$, $n=32$, $n=18$), which might have biased our findings (non-representative/selective inclusion). Secondly, our study compared three groups of children with ASD i.e., severe, mild, and no disorganized speech amongst each other, while previous studies compared children with ASD to typically developing children and/or other disorders [3,30]. In other words, children in our sample were more similar in diagnosis to each other as compared to the groups compared in other studies. Finally, our social pragmatic communication measurement was based on parents’ report only and not on direct observations of the children which may create bias in reporting about their child (i.e., under-or over report of problems based on the behaviors that parents observe mainly in the home context). However, [8] also showed associations between disorganized speech and social communication problems as observed with the ADOS-2.

Recommendations for future research

Since the current sample size was small for cross-tabulations, future research with bigger sample size is recommended. Second, the current findings show that disorganized speech seems to be associated with some aspects of social pragmatic communication skills and ToM. Longitudinal studies are also recommended because these could give more insight in

the development of disorganized speech over time and associations with developing ToM.

Conclusion and Implications of the Research findings for practice

In conclusion, the presence of disorganized speech in children with ASD should not be considered a pre-psychotic sign but rather, disorganized speech should be considered as a specific problem with social pragmatic communication. Moreover, severe disorganized speech (i.e., showing loose associations as well as illogical reasoning/abrupt topic shifts) may be related to difficulties with theory of mind, but mainly to Verbal IQ.

Although these findings are preliminary and should thus be replicated and considered in the light of the limitations of our study, readers could consider offering children with ASD and disorganized speech interventions that stimulate and train broader social pragmatic communication skills, social cognition, and verbal ability (i.e., conversational skills). An evidence-based intervention that meets these requirements is the Children Friendship Training. This intervention is designed for children in the age range of 6 to 12 years old. CFT explains rules of social skills explicitly which creates clear and concrete handles for children with ASD and disorganized speech. Children are encouraged to reflect on the perspectives of others and actively participate in social situations, which could help to generalize the obtained skills in social cognition and communication to other contexts. Differences in Verbal IQ may need to be considered as a moderator of treatment outcome.

Acknowledgement

The authors would like to thank Mart Eussen and Esther de Bruin for the support and commitment. We also would like to acknowledge the effort of Marleen de Vries for the data exploration and data cleaning at the beginning of the manuscript writing. We thank all our research colleagues for their helpful suggestions.

Conflicts of Interest

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

This study was supported financially by a grant from the Netherlands Organization for Scientific Research (NWO/ZonMw/OOG-100-002-006).

Availability of Data and Material

The datasets generated and/or analyzed during the current study are not publicly available but, are available from the corresponding author on reasonable request after the results are published.

Code Availability

Not applicable.

Authors' Contributions

All authors contributed to the study conception and design. Material preparation, data collection and analysis were performed by Sakinah Idris, Anneke Louwerse, Jan van der Ende, and Kirstin Greaves-Lord. The

first draft of the manuscript was written by Sakinah Idris and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Ethical Approval

All methods were carried out in accordance with the Declaration of Helsinki. The protocol was approved by the Ethics committee of the Erasmus University Medical Center in accordance with the Declaration of Helsinki.

Consent to Participate

All participants and their parents gave written informed consent prior to study participation.

Consent for Publication

Not applicable.

References

1. American Psychiatric Association. "Diagnostic and Statistical Manual of Mental Disorders, Fourth edition, Text Revision (DSM-IV-TR) (3rd ed.)." Washington, DC: *American Psychiatric Association Press* (2000).
2. Andreasen, N. C. "Thought, language & communication disorders." *Arch Gen Psychiatry* 36(1979): 1315-1321.
3. Norbury, CF, Gemmell T and Paul R. "Pragmatics abilities in narrative production: A cross-disorder comparison." *J Child Lang* 41(2014):485-510.
4. Solomon, M, Ozonoff S, Carter C and Caplan R. "Formal thought disorder and the autism spectrum: relationship with symptoms, executive control, and anxiety." *J Autism Dev Disord* 38(2008):1474-1484.
5. Swineford, LB, Thurm A, Baird G, Wetherby AM and Swedo S. "Social (pragmatic) communication disorder: a research review of this new DSM-5 diagnostic category." *J Neurodev Disord* 6(2014): 41.
6. Caplan, GD, Fish B, Tanguay PE and David LG. "The kiddie formal thought disorder rating scale: Clinical assessment, reliability and validity." *J Am Acad Child Adolesc Psychiatry* 28(1989):408-416.
7. Liddle, PF, Ngan ET, Caissie SL, Anderson CM and Bates AT, et al. "Thought and Language Index: an instrument for assessing thought and language in schizophrenia." *Br J Psychiatry* 181(2002): 326-330.
8. Eussen, MLJM, Louwerse A, Verhulst FC, de Bruin EI and Ende VD, et al. "Formal thought disorder in autism spectrum disorder predicts future symptom severity, but not psychosis prodrome." *Eur Child Adolesc Psychiatry* 24(2014):163-172.
9. Caplan, GD, Tang B, Komo S and Asarnow RF. "Thought disorder in childhood schizophrenia: Replication and update of concept." *J Am Acad Child Adolesc Psychiatry* 39(2000):771-778.
10. Simmons, ES, Paul R and Volkmar F. "Assessing pragmatic language in autism spectrum disorder: The yale in vivo pragmatic protocol." *J Speech Lang Hear Res* 57(2014):2162-2173.
11. Hale, CM and Tager FH. "Social communication in children with autism: The relationship between theory of mind and discourse development." *Autism* 9(2005):157-178.
12. Wimmer, H and Perner J. "Beliefs about beliefs: Representation and constraining function of wrong beliefs in young children's understanding of deception." *Cognition* 13(1983):103-128.

13. Baron-Cohen, S, Leslie AM and Frith U. "Does the autistic child have a "theory of mind?" *Cognition*, 21(1985):37-46.
14. Frith, UW. "Why we need cognitive explanations of autism." *Q J Exp Psychol* 65(2012):2073-2092.
15. Baren-Cohen, S. "The autistic child's theory of mind: A case of specific developmental delay." *J Child Psychiatry* 30(1989): 285-297.
16. Cummings, L, Capone FL, Piparo and Carapezza M. "Clinical pragmatics and theory of mind." *Perspec on Ling Pragmatics* 2(2013):23-59.
17. Tager, FH. "Evaluating the theory-of-mind hypothesis of autism." *Curr Dir Psychol Sci* 16(2007):311-315.
18. Capps, L, Kehres J and Sigman M. "Conversational abilities among children with autism and children with developmental delays." *Autism* 2(1998):325-344.
19. Capps, L, Losh M and Thurber C. "The frog ate the bug and made his mouth sad": Narrative competence in children with autism." *J Abnorm Child Psychol*, 28(2000):193-204.
20. De Bruin, EI, Verheij F, Wiegman T and Ferdinand RF. "Assessment of formal thought disorder: The relation between the kiddie formal thought disorder rating scale and clinical judgment." *Psychiatry Res* 149(2007):239-246.
21. De Bruin, EI. "The chaotic mind: A study of children with thought disorders and social contact problems." *Psychol* 7(2006):261-267
22. Van der, GRJ, Caplan R, Van EH, Loman F and Buitelaar JK. "A controlled study of formal thought disorder in children with autism and multiple complex developmental disorders." *J Child Adoles. Psychopharmacol* 15(2005):465-476.
23. Bishop, DVM. "Development of the children's communication checklist (CCC): A method for assessing qualitative aspects of communicative impairment in children." *J Child Psychol Psychiatry* 39(1998): 879-891.
24. Hartman, CA, Geurst HM, Bennink AC, Verte S and Roeyers H, et al. "De Drie C's: Children's communication checklist." (1998).
25. Geurts, HM, Hartman C, Verte S, Oosterlaan J and Roeyers H, et al. "Pragmatics fragmented: The factor structure of the Dutch Children's Communication Checklist (CCC)." *Int J Lang Commun Diso* 44(2009):549-574.
26. Walker, S. "Gender differences in the relationship between young children's peer-related social competence and individual differences in theory of mind." *J Genet Psycho*, 166(2005):297-312.
27. Lai, MC, Lombardo MV, Auyeung B, Chakrabarti B and Baron CS. "Sex/gender differences and autism: setting the scene for future research." *J Am Acad Child Adolesc Psych* 54(2015):11-24.
28. Schirvar, WM. "Investigating social competence in students with high intelligence: University of Minnesota Digital Conservancy." (2013) Retrieved from <http://hdl.handle.net/11299/154328>.
29. Field, A. "Discovering statistics using SPSS." Sage(2009).
30. Kuijper, SJM, Hartman CA, Bogaerds HSTM and Hendriks P. "Narrative production in children with ASD and children with ADHD: Similarities and differences." *J Abnorm Psychol* 126(2015):63-75.
31. Roberts, RJ and Patterson CJ. "Perspective taking and referential communication: The question of correspondence reconsidered." *Child Development* 54(1983):1005-1014.
32. Tager, FH, Paul P and Lord C. "Language and communication in autism." In F. R. Volkmar, A. Klin, R. Paul, & D. J. Cohen (Eds.), *Handbook of Autism and Pervasive Developmental Disorders, Section II: Development and Behavior* . (3 ed.). Hoboken, NJ: Wiley (2005).
33. Bishop, DVM. "Parent and teacher report of pragmatic aspects of communication: Use of the Children's Communication Checklist in a clinical setting." *Dev Med Child Neurol* 43(2001): 809-818.
34. Tager, FH. "Brief report: Current theory and research on language and communication in autism." *J Autism Dev Disord* 26(1996):169-172.
35. Volden, J, Coolican J, Garon N, White J and Bryson S. "Brief report: pragmatic language in autism spectrum disorder: relationships to measures of ability and disability." *J Autism Dev Disord* 39(2009):388-393.
36. Baird, G and Norbury CF. "Social (pragmatic) communication disorders and autism spectrum disorder." *Arch Dis Child* 101(2015):745-751.
37. Visser, S and Tops W. "Social pragmatic communication disorder and autism spectrum disorder: Two of a Kind? A narrative review." *Ann Clin Psychiatry* 5(2017):1121-1125.
38. Frankel, F, Myatt R, Sugar C, Whitham C and Gorospe CM, et al. "A randomized controlled study of parent-assisted Children's Friendship Training with children having autism spectrum disorders." *J Autism Dev Disord* 40(2010): 827-842.
39. Jones, CD and Schwartz IS. "When asking questions is not enough: An observational study of social communication differences in high functioning children with autism." *J Autism Dev Disord* 39(2009): 432-443.
40. McCann, J, Peppe S, Gibbon FE, O'Hare A and Rutherford M. "Prosody and its relationship to language in school-aged children with high-functioning autism." *Int J Lang Commun Disord* 42(2007):682-702.

How to cite this article: Idris, Sakinah, Louwse A, Van der Ende J and Hillegers M, et al. "Disorganized Speech in Autism Spectrum Disorder: Part of the Problems in Social Pragmatic Communication and Verbal Intelligence" *J Neurol Disord* 9(2022):484.