

Editorial Note on Impact of COVID-19 on Environment

Sukriti Mittal

Department of Environmental Studies, Central University of Jammu & Kashmir, India

Editorial Note

Across the nation lockdown because of the pandemic has prompted an extreme decrease of NO_2 discharges and diminished air contamination levels. It's significant this is kept up considerably after the lockdown lifts. In the midst of the overwhelming COVID-19 pandemic, an uncommon positive has been the noteworthy worldwide decline in air contamination levels. Basically, specialists have estimated nitrogen dioxide (NO_2), one of the six significant air poisons (notwithstanding particulate issue, carbon monoxide, sulfur dioxide, ground-level ozone, and lead). NO_2 has, as most different gases, normal and human sources.

Characteristic sources incorporate lightning, seas, and volcanoes. Be that as it may, in urban locales, regular sources of NO_2 represent a little part of the all-out NO_2 levels; as indicated by a 2005 report by Australia's Department of the Environment and Heritage; normal sources of NO_2 just record for 1 percent of by and large NO_2 levels in urban communities. Human action is as a rule answerable for NO_2 emanations in urban districts, with street transport being the main source. Planes, power plants and ships, all of which consume petroleum products, are likewise huge human sources of NO_2 . Given this present, it's obvious that during the severe worldwide lockdowns, NO_2 levels

have dropped altogether in urban territories, particularly in India's thickly populated urban communities.

Satellite symbolism from the European Space Agency's Copernicus Sentinel 5P satellite measure NO_2 levels comprehensively. These estimations precisely reflect emanations sources, in light of the fact that dissimilar to different gases that can travel a critical good ways from where they're radiated, NO_2 has a short life expectancy and kicks the bucket before it can move far. At the end of the day, if the Sentinel 5P satellite catches a hotspot of NO_2 over Delhi, it's almost certain it was transmitted from inside Delhi's region. Satellite symbolism is, in this way, an exceptionally dependable instrument for estimating NO_2 emissions, particularly if information with significant levels of cloud inclusion is prohibited.

The worldwide abatements in NO_2 levels were first found in China, where levels dove significantly following the exacting isolate measures implemented in late January. As countries in Europe and North America followed China's lead in late February and March, comparative patterns have been watched all around. India's across the nation lockdown, specifically, has significantly affected air contamination levels. With residents isolated at home, street transportation and force plant activities have gone to a crushing stop, and contamination levels in all over the nation, particularly in normally smoggy urban communities, have been observed to go down gradually.

How to cite this article: Sukriti Mittal. "Editorial Note on Impact of COVID-19 on Air Quality." *J Environ Anal Toxicol* 10:6 (2020) 1-1.

***Address for Correspondence:** Sukriti Mittal, Department of Environmental Studies, Central University of Jammu & Kashmir, India. Tel: 9155875649; E-mail: Sukritimittal2010@gmail.com.

Copyright: © 2020 Mittal S. This is an open-access article distributed under the terms of the creative commons attribution license which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Received 20 November, 2020; **Accepted** 25 November, 2020; **Published** 30 November, 2020