

Effect of Yoga on Depression among Alcoholics in Nainital

*Travis Fuller

Department of Nursing, University of Shri Venkateshwara, Amroha, India

Description

The highest consumption levels are recorded in Pondicherry, Chandigarh, Goa and Delhi in that order. Since all of these have lower taxes on alcohol than other states, their consumption numbers are almost certainly boosted by people from across the border buying cheap liquor. In the case of Pondicherry and Goa, the high volume of tourist relative to the local population could also be contributing to the high consumption.

The World Health Organization estimates that there are about 2 billion people worldwide who consume alcoholic beverages and 76.3 million with diagnosable alcohol use disorders [1]. Worldwide per capita consumption of alcoholic beverages equaled 6.3 liters. India is producing 70% of total alcoholic product in South East Asia and importing 10% of its alcoholic consumption. Tamil Nadu ranks first among states with large number of alcoholics. The highest consumption levels are recorded in Pondicherry, Chandigarh, Goa and Delhi in that order. Since all of these have lower taxes on alcohol than other states, their consumption numbers are almost certainly boosted by people from across the border buying cheap liquor [2]. In the case of Pondicherry and Goa, the high volume of tourist relative to the local population could also be contributing to the high consumption figure. Patterns and levels of alcohol consumption, alcohol dependency and alcohol abuse are determined by many factors; such as availability, income per capita, retail process, individual factors such as age of first use, family history, education, peer group pressure, psychosocial factors, cultural, historical context and government policies such as taxation, restrictions on advertisement and promotion [3]. There are 26 De-Addiction centers in Tamil Nadu and 14 centers are there in Coimbatore which provide treatment for detoxification and therapies like individual counselling, group therapy, are mainly spouse counselling, family counselling, yoga and rehabilitation for needy patients. Beyond this treatment modalities, Sudarshan Kriya Yoga helps the alcoholic patients to reduce the depression level, anxiety, stress and hypertension. The different

literatures regarding alcohol dependence suggests that they are facing many problems in society, occupational areas, emotional aspects and financial crisis [4]. Sudarshan Kriya Yoga helps to detoxifies the body and purifies the mind also helps in decreasing the symptoms of depression and anxiety [5]. The main benefit of yoga is to reduce depression and anxiety. Depression which affects large number of people in the world. Yoga has the direct effect of neurotransmitter of the brain. Practicing yoga will increase the serotonin which involved in depression.

The main aim of the study was to assess the effect of Yoga on depression among alcoholics. Quasi experimental Nonequivalent control group design was adopted in this study [6]. In the present study, 30 alcoholics were recruited in experimental group and 30 were in control group by using purposive sampling technique. Sudarshan Kriya Yoga was administered to the samples for a period of 7 days in a week in experimental group [7]. The depression of alcoholics was assessed before and after intervention by using beck depression inventory. The data were analyzed and the findings are discussed based on the objectives of the study. In the experimental group, (42%) Alcoholics belong to the age group between 31 years-40 years and in control group, (43.33%) Alcoholics belong to the age group between 31 years-40 years.

Educational status of Alcoholics, reveals that, most of them are completed higher secondary education in experimental (46%) and control group (56.6%). In the experimental group, majority, (54%) Alcoholics we are working under private sector and in control group, (56.6%) Alcoholics were working under the private sector. Data on monthly income which reveals that, most of them earning of about Rs 10,001- Rs 15000t in both experimental (44 %).

The data on religion which reveals that, majority were belongs to Hindu religion in both experimental (73%) and control group (68%). The data on residence which reveals that majority resides in urban area in both experimental (66%) and control group (63.33%).

*Address for Correspondence: Travis Fuller, Department of Nursing, University of Shri Venkateshwara, Amroha, India, E- mail: travisfuller211@gmail.com

Copyright: © 2022 Fuller T. This is an open-access article distributed under the terms of the creative commons attribution license which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Received: 23 November, 2019, Manuscript No. JIDM-22-4902; Editor assigned: 27 November, 2019, PreQC No. P-4902; Reviewed: 12 December, 2019, QC No. Q-4902; Revised: 25 August, 2022, QI No JIDM-22-4902. Manuscript No. R- 4902; Published: 22 September, 2022, DOI: 10.37421/2576-1420.2022.7.244

Marital status reveals that in experimental group, most of them (89%) were married, and in control group (76.7%) were married. The data on number of children explain that most of the alcoholics have two children both experimental (59.18%) and control group (56.17%). The data on number of times hospitalization due to alcoholism reveals that in majority of Alcoholics were hospitalized 2 times to the maximum in both experimental (66%) and control group (64%). Data on duration of alcohol intake reveals that most of them in both experimental (45%) and control group (41.3%) were consuming alcohol between 5 years-10 years. Data on age at first taken alcohol reveals that in both most of them had started to taking alcohol between 15 years-25 years of age in experimental (44%) and in control group (56.4%). Data on occasion of drinking reveals that most of them consuming alcohol throughout the day and only at the night time in both experimental (46%).

Conclusion

Data on of mode of introduction of alcohol reveals that most of them was introduced to alcohol by their friends in both experimental (56.1%) and control group (73.4%). Data on type of drink used among alcoholics reveals that in experimental group (66%) have used brandy and in control group (56.6%) used brandy, (26.6%) used beer and (46.6%) used whisky. Data on intake of alcohol with companionship reveals that majority take alcohol without companionship in both experimental (63%) and control group (67%). The level of depression among alcoholic's analysis of depression level among 40 Alcoholics shows, in experimental group (30%) had moderate level of depression and (70%) had severe level of depression. In control group, (44%) had moderate level of depression, (66%) had severe level of depression Level of Depression among experimental and control group before and after intervention Before intervention, (30%) Alcoholics from experimental group had moderate level of depression, and (70%) had severe level of depression. whereas after intervention, (66%) had mild level of depression and (44%) had moderate level of depression. In control group, pretest score showed that (44%) Alcoholics had moderate level of depression, (66%) had severe level of depression

and posttest score showed that, (70%) had moderate level of depression, (30%) had severe level of depression. Association between the level of depression and selected variables The association between the level of depression and selected variables among Alcoholics were calculated using Chi square for variables like age, educational status, occupation, Area of residence, marital status, duration of alcohol intake, Number of children, number of times admission in hospitalization due to alcoholism, age at first taken alcohol, occasion of drinking, mode of introduction, type of drink, intake of alcohol without companionship had no association with the level of depression among Alcoholics. It was found that there was no significant association.

References

1. Carey, MP, and Rao, P "Prevalence of Alcohol Abuse Associated Factors in Former Drinkers." *Scielo Publ Hlth* 41 (2004): 509.
2. BS, Chavan, Arun P, and Bhargava R. "Prevalence of Alcohol and Drug Dependence in Rural and Slum Population of Chandigarh." *Indian J Psychiatry* 49 (2009): 44-48.
3. Doria, S. "Anti-Anxiety Efficacy of Sudarshan Kriya Yoga in General Anxiety Disorder." *J Affect Disord* 12 (2015): 43-49.
4. FB, Grant. "Co-Occurrence of 12-Month Alcohol and Drug Use Disorders and Personality Disorders: Results from the National Epidemiologic Survey on Alcohol and Related Conditions." *Archiv Gen Psychiat* 29 (2004): 129-130.
5. T, Kevin, Larkin, and Claudia. "Anger Expression and Essential Hypertension: Behavioral Response to Confrontation." *J Psychos Res* 56 (2004): 113-118.
6. J, Hegde, and Shetty T. "Biological Effects of Sudarshan Kriya Yoga in Alcoholics." *NIMHANS J* 84 (2005): 647-653.
7. G, Jain, Rajak C, and Sanjeev. "Effect Of Sudarshan Kriya Yoga on Volunteers Having Anxiety along with Drinking Habit at AIIMS." *Ind J Psychiat* 69 (2009): 241-246.

How to cite this article: Fuller, Travis. "Effect of on Depression among Alcoholics in Nainital." *J Infect Dis Med* 7 (2022): 244.