

Energy Balance: Through the Looking Glass

Brilla LR^*

Western Washington University, Bellingham, USA

Corresponding author: Brilla LR, Western Washington University, Bellingham, USA, Tel: 360-650-3056; Fax: 360-650-7447; E-mail: Lorrie.Brilla@wwu.edu

Received date: June 06, 2016; Accepted date: June 06, 2016; Published date: June 15, 2016

Copyright: © 2016 Brilla LR. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Introduction

Human metabolic energy balance has been an area of intrigue and confusion for decades. An imbalance between energy intake and energy expenditure is considered the primary etiology for weight changes. Behavioral interventions, however, have generally resulted in a smaller than expected weight loss, which in part has been attributed to compensatory adaptations in other components contributing to energy balance [1,2]. Conversely, physically active individuals can exhibit negative energy balance without weight loss [3-5].

Energy balance is commonly used for understanding weight change and can be represented numerous ways, frequently by the equation [6]:

change in energy stores=energy intake-energy expenditure

Despite its use as the prevalent equation for energy, it lacks consideration regarding additional factors. Hence, simplistic notions based on "energy in and energy out" are largely ineffective. Human energy homeostasis is complex and there are many unanswered questions. This commentary reviews assumptions related to energy balance, especially emphasizing weight maintenance in exercising individuals. Additionally, paths for further studies are presented that may address some of the discrepancies in the current research literature.

Thermodynamics and Estimates

A common belief is that reducing one's caloric intake or increasing caloric expenditure should create an energy deficit, resulting in weight loss, thus complying with the first law of thermodynamics, which states that energy can be transformed from one form to another but cannot be created or destroyed. This accordance with the first law of thermodynamics is that energy within a closed system remains constant. As appealing as this ideology is, human physiology is not that simple, as humans are an open system within the biosphere.

Part of the noncompliance of data to support a simple energy equation is related to the ubiquitous measurement errors. Often, energy intake is estimated by food records or recall [7-10]. Underestimation of energy intake is a common finding. Redman and colleagues reported individuals significantly under reported energy intake (350 kcal/d; 15%), and underreporting by overweight individuals (~400 kcal.d⁻¹; 16%) was greater than that of normal-weight individuals (~270 kcal.d⁻¹; 12%) [9]. However, Ptomey and colleagues have reported on a new methodology for energy intake estimated by digital photography, which closely matched ($6.8\% \pm 28\%$) with the energy expenditure gold standard of doubly labeled water [8].

The other aspect of the energy balance determination is energy expenditure. Although calorimetric chambers and doubly labeled water are state-of-the-art techniques, they are not always available [9,11]. Thus, indirect calorimetry can be used for resting metabolic rate (RMR) or during exercise, while accelerometers and metabolic equivalents (METs) of physical activity are methods used during physical activity [12,13]. The value equating 1 MET to 3.5 ml O_2 Kg⁻¹.Min⁻¹ or 1 Kcal.Kg⁻¹. hr⁻¹ was derived from the resting O_2 consumption of one person, a 70-kg, 40-yr-old man. The MET value has come under scrutiny during rest and a set walking workload in 642 women and 127 men, 18-74 yr, 35 kg to 186 kg in weight, who were weight stable and healthy [13]. Resting values measured by indirect calorimetry were significantly lower than the commonly accepted 1 MET value. Body composition accounted for 62% of the variance.

Daily energy expenditure is the more difficult of the contributors to energy balance to obtain accurate measurements due to variability in putative compensatory effects [14]. These inconsistencies may be related to differences in RMR, exercise-related appetite changes, and reduced spontaneous physical activity. There is a large variability in resting energy expenditure (~250 kcal.d⁻¹) that is not explained by differences in body composition [1]. In "The Biggest Loser" weight loss competition, RMR decreased significantly more than expected based on measured body composition changes. The magnitude of this metabolic adaptation was correlated with the degree of energy imbalance (r=0.55) [15]. Exercise programs consistent with public health recommendations may promote modest weight loss (~2 kg) however responses are highly heterogeneous [16]. Lower energy compensation occurs with short-term exercise, and a much higher level of energy compensation accompanies long-term exercise interventions [14]. Longer duration studies are obviously needed. Additionally, in many studies, the energy deficit produced by the prescribed exercise may be far less than that produced by dietary restriction [17]. Further, it was reported that subjects increased 24 h energy expenditure by 209 ± 555 kcal over baseline as assessed by doubly labeled water, yet lost no weight. Although energy intake was measured periodically and did not increase, some compensatory change in calorie intake must have occurred for body weight to remain stable.

High-intensity exercise has a beneficial impact on 24 h energy balance, as measured in a calorimetric chamber, resulting in spontaneous decrease in energy intake [11]. In normal-weight and overweight men, increasing physical activity while keeping energy ad libitum over an 8-week period produces a prolonged negative energy balance [18]. In a longer study, over a 10-month period, exercise training did not significantly alter energy intake in young adults [19]. Energy intake drops upon initiating an increased exercise volume. Subsequently, intake begins to increase in order to provide compensation for about 30% of the energy expended in activity. This compensation is partial and incomplete over short term studies [20]. Exercise-induced alterations in appetite may be driven by complex changes in appetite-regulating hormones [21], among other factors. Conversely, it can be demonstrated that when active individuals are forced into a sedentary state, energy intake does not decrease to match the reduced energy expenditure.

There is minimal evidence to support that exercise training results in decreased non-exercise energy expenditure in healthy adults [12]. A 10-month aerobic exercise training program in previously sedentary, overweight and obese young adults was not associated with compensatory decreases in non-exercise energy expenditure. Results suggest that individuals do not become less physically active or spend more time in sedentary pursuits in response to exercise [22]. Generally, humans seem to be better equipped to defend against weight loss than avoid weight gain, but results also show a large individual variability. Therefore, individual differences should be explored to identify specific characteristics of compensator and non-compensators [23]. Dynamic energy imbalances definitely need further exploration [24].

Athletes' Defense to Weight Loss

Athletes have been reported to have energy imbalances. Discrepancies between energy intake and expenditure in physically active women was described in the 1990s in weight stable runners [3]. Daily energy intake equaled energy expenditure in non-runners, but energy expenditure in women runners significantly exceeded intake, a deficit of approximately 650 kcal.d⁻¹. The runners showed no evidence of compensating with decreased non-exercise energy expenditure. The findings suggest that women adapted to high levels of activity may possess mechanisms to maintain body weight without significantly increasing energy intake. Further, energy balance in weight stable athletes with and without menstrual disorders was assessed [25]. A small energy deficit was observed in runners with irregular menstrual function but not in in a comparable group of runners with normal menstrual function. Thyroid hormone was also measured with significantly lower levels of free thyroxin in runners demonstrating irregular menstrual function, which may indicate an adaptive lower RMR in these athletes.

Male endurance athletes may also exhibit energy imbalances, though they are weight stable. When RMR was assessed in low-energy intake and adequate-energy intake male endurance athletes who had been weight stable for 2 years, RMR was significantly lower in lowenergy intake athletes. The difference between groups was about 158 kcal.day⁻¹ [26]. The researchers postulated that lower RMR is one mechanism that contributes to weight maintenance when energy intake is less than energy expenditure. In another study that evaluated energy balance during high-volume and low-volume training periods among male endurance athletes, significant differences were reported in lower energy intake [27]. During the course of the study, there was no difference in body weight or composition. There was no compensation of increasing energy intake to match energy expenditure during the increased volume training.

Sex Incongruity

One major source of variability in energy balance might be sex. The studies on athletes demonstrated a reduced RMR in both females and males [25,26]. In comparison of non-athletic and athletic women, RMR per kilogram body weight was wide for nonathletic, but narrow for athletic women. The researchers postulated metabolic efficiency was highly variable in both lean and obese nonathletic women [28]. If higher efficiency is present, then it would predispose those women to developing body fat. However, the mean lower RMR in athletes may have some compensatory mechanism that defends against weight gain.

Conflicting evidence exists on RMR findings in response to variable energy expenditure. Long term exercise training studies have shown both that training increases RMR while other studies have failed to support these findings [5,29]. Intense exercise training may induce reductions in RMR, in spite of the increased lean tissue mass, similar to the changes observed in animals in response to flight [5]. There may be sex differences to account for the observed energy imbalances, as well. In 24 h energy expenditure, females had 5% to 10% lower values compared with males after adjusting for differences in body composition, age, and activity [30]. RMR was approximately 120 kcal.day⁻¹ lower in females than males. Assessed spontaneous physical activity was not significantly different between males and females in that study. In another study, total daily energy expenditure was about 20% higher in men than in women, which equated to 580 kcal.day⁻¹ [9]. Noting the differences in RMR and total daily energy expenditure from these studies, there must be some metabolic efficiency changes that contribute to the energy imbalance with stable weight.

In a 10-month study exercise intervention to promote weight loss, responders were defined as those who lost \geq 5% of body weight. Men who were non-responders had higher energy intake and lower nonexercise energy expenditure when compared with men who responded to the exercise, which may account for not losing as much weight. No significant differences in any parameters assessed were observed between women who did not meet the threshold weight loss compared to those losing \geq 5% of body weight. The researchers suggest that factors associated with the weight loss response to exercise in women warrant additional investigation [31].

Microbiota

A flurry of research has emerged on intestinal bacteria and how the microbiota may influence body weight in addition to promising health effects. However, adverse effects can also occur such as reported in a case study in a woman who was treated with fecal microbiota transplantation [32]. The young woman had been normal weight with a BMI of 26. After 16 months, her BMI had increased to 33; she had received the transplant from an obese donor.

Exercise effects on the microbiota have been reported [33], and limited animal and human research findings imply that exercise may have a beneficial role on microbiome characteristics [34]. In mice, exercise has a strong influence on gut integrity and host micro biome [35]. Heterogeneity in energy derived from energy intake has been the main focus of many recent studies. However, from the few studies to date, effects of habitual exercise on the intestinal micro biota have demonstrated positive health effects and support further rigorous research concerning energy balance.

Autonomic Influence

Some of the variance in energy balance may be related to autonomic system influences. A thermodynamic model demonstrates the sympathetic and parasympathetic nervous systems regulate energy [36]. The autonomic nervous system (ANS) plays a critical role in the control of energy balance and body weight [37]. The ANS is involved in the control through influences exerted on the production and loss of heat. Activation of the sympathetic discharge causes an increase in energy expenditure and a decrease in energy intake. Paradoxically, RMR is higher in the athletes than in sedentary women, despite the augmented parasympathetic activity that is usually related to lower energy expenditure.

Page 2 of 4

Research shows enhanced resting metabolic rate (RMR) in trained athletes is an acute effect of prior exercise induced by catecholamines, and not due other factors such as thyroxin levels [38]. Conversely, adaptation to exercise training is associated with a relative enhancement of vagal dominance [39-41]. In both male and female subjects, high vagal influence is negatively correlated with BMI [41,42]. These results provide evidence for a prominent role of the vagal tone in the modulation of the energy expenditure of the human. Further, intensive training resets towards sympathetic dominance [43,44]. Under similar training strain, male athletes showed constantly higher markers of sympathetic activity than female athletes [45]. Sympathetic influence on reduced energy intake may contribute to observations during multi-stage exercise events of high volume and intensity, such as the Tour de France. Exercise effects on the ANS influence on energy balance needs to be explored more fully. Further studies should be addressed to reveal new aspects of the control exerted by the autonomic nervous system on body weight.

Exercise Perspective on Energy Balance

Energy balance is a complex phenomenon. The thermodynamic equation for energy balance is an oversimplification. The body is not a closed system, since there are energy fluxes with the biosphere. In addition to the promising areas of research briefly presented in this paper, other influences include genetics and hormonal balances, among others. The list of peripheral anorexigenic and orexigenic physiological factors is intimidating and expanding. Exercise may modify responses in a direction expected to enhance satiety [46], and yet permit weight maintenance. The possibility of a threshold level beyond which increased exercise energy expenditure fails to produce a more negative energy balance and potential sex differences in the energy intake response to increased levels of exercise are potentially important. Much research focus has been on negative energy balance factors to address the expanding prevalence of obesity in society. However, in examining energy balance from the exercise perspective, through a looking glass, may provide understanding that otherwise may be not be achieved. This brief commentary emphasizes the need for more research with simultaneous measurements of all major components contributing to energy balance to enhance understanding of the complex regulation.

References

- Hall KD, Heymsfield SB, Kemnitz JW, Klein S, Schoeller DA, et al. (2012) Energy balance and its components: implications for body weight regulation. Am J Clin Nutr 95: 989-994.
- 2. Hill JO, Wyatt HR, Peters JC (2012) Energy balance and obesity. Circulation 126: 126-132.
- 3. Mulligan K, Butterfield GE (1990) Discrepancies between energy intake and expenditure in physically active women. Br J Nutr 64: 23-36.
- 4. Thompson J, Manore MM, Skinner JS (1993) Resting metabolic rate and thermic effect of a meal in low- and adequate-energy intake male endurance athletes. Int J Sport Nutr 3: 194-206.
- 5. Speakman JR, Selman C (2003) Physical activity and resting metabolic rate. Proc Nutr Soc 62: 621-634.
- Galgani J, Ravussin E (2008) Energy metabolism, fuel selection and body weight regulation. Int J Obes (Lond) 32: S109-S119.
- Kowalkowska J, Slowinska MA, Slowinski D, Dlugosz A, Niedzwiedzka E, et al. (2013) Comparison of a full food-frequency questionnaire with the three-day unweighted food records in young Polish adult women: implications for dietary assessment. Nutrients 5: 2747-2776.

- Ptomey LT, Willis EA, Honas JJ, Mayo MS, Washburn RA, et al. (2015) Validity of energy intake estimated by digital photography plus recall in overweight and obese young adults. J Acad Nutr Diet 115: 1392-1399.
- Redman LM, Kraus WE, Bhapkar M, Das SK, Racette SB, et al. (2014) Energy requirements in nonobese men and women: results from CALERIE. Am J Clin Nutr 99: 71-78.
- Voss S, Kroke A, Klipstein-Grobusch K, Boeing H (1998) Is macronutrient composition of dietary intake data affected by underreporting? Results from the EPIC-Potsdam Study. European Prospective Investigation into Cancer and Nutrition. Eur J Clin Nutr 52: 119-126.
- 11. Thivel D, Isacco L, Montaurier C, Boirie Y, Duché P, et al. (2012) The 24-h energy intake of obese adolescents is spontaneously reduced after intensive exercise: a randomized controlled trial in calorimetric chambers. PLoS One 7: e29840.
- Washburn RA, Lambourne K, Szabo AN, Herrmann SD, Honas JJ, et al. (2014) Does increased prescribed exercise alter non-exercise physical activity/energy expenditure in healthy adults? A systematic review. Clin Obes 4: 1-20.
- 13. Byrne NM, Hills AP, Hunter GR, Weinsier RL, Schutz Y. (2005) Metabolic equivalent: one size does not fit all. J Appl Physiol (1985) 99: 1112-1119.
- Riou MÈ, Jomphe-Tremblay S, Lamothe G, Stacey D, Szczotka A, et al. (2015) Predictors of energy compensation during exercise interventions: A systematic review. Nutrients 7: 3677-3704.
- 15. Knuth ND, Johannsen DL, Tamboli RA, Marks-Shulman PA, Huizenga R, et al. (2014) Metabolic adaptation following massive weight loss is related to the degree of energy imbalance and changes in circulating leptin. Obesity (Silver Spring) 22: 2563-2569.
- Swift DL, Johannsen NM, Lavie CJ, Earnest CP, Church TS (2014) The role of exercise and physical activity in weight loss and maintenance. Prog Cardiovasc Dis 56: 441-447.
- Catenacci VA, Wyatt HR (2007) The role of physical activity in producing and maintaining weight loss. Nat Clin Pract Endocrinol Metab 3: 518-529.
- Melzer K, Renaud A, Zurbuchen S, Tschopp C, Lehmann J, et al. (2016) Alterations in energy balance from an exercise intervention with ad libitum food intake. J Nutr Sci 5: e7.
- Washburn RA, Honas JJ, Ptomey LT, Mayo MS, Lee J, et al. (2015) Energy and macronutrient intake in the Midwest Exercise Trial 2 (MET-2). Med Sci Sports Exerc 47: 1941-1949.
- 20. Blundell JE, Stubbs RJ, Hughes DA, Whybrow S, King NA (2003) Cross talk between physical activity and appetite control: does physical activity stimulate appetite? Proc Nutr Soc 62: 651-661.
- Larson-Meyer DE, Palm S, Bansal A, Austin KJ, Hart AM, et al. (2012) Influence of running and walking on hormonal regulators of appetite in women. J Obes 2012: 730409.
- 22. Willis EA, Herrmann SD, Honas JJ, Lee J, Donnelly JE, et al. (2014) Nonexercise energy expenditure and physical activity in the Midwest Exercise Trial 2. Med Sci Sports Exerc 46: 2286-2294.
- Drenowatz C (2015) Reciprocal compensation to changes in dietary intake and energy expenditure within the concept of energy balance. Adv Nutr 6: 592-599.
- Hall KD, Sacks G, Chandramohan D, Chow CC, Wang YC, et al. (2011) Quantification of the effect of energy imbalance on bodyweight. Lancet 378: 826-837.
- Tomten SE, Høstmark AT (2006) Energy balance in weight stable athletes with and without menstrual disorders. Scand J Med Sci Sports 16: 127-133.
- Thompson J, Manore MM, Skinner JS (1993) Resting metabolic rate and thermic effect of a meal in low- and adequate-energy intake male endurance athletes. Int J Sport Nutr 3: 194-206.
- Drenowatz C, Eisenmann JC, Carlson JJ, Pfeiffer KA, Pivarnik JM (2012) Energy expenditure and dietary intake during high-volume and lowvolume training periods among male endurance athletes. Appl Physiol Nutr Metab 37: 199-205.

- Owen OE, Kavle E, Owen RS, Polansky M, Caprio S, et al. (1986) A reappraisal of caloric requirements in healthy women. Am J Clin Nutr 44: 1-19.
- Poehlman ET, Melby CL, Badylak SF (1988) Resting metabolic rate and postprandial thermogenesis in highly trained and untrained males. Am J Clin Nutr 47: 793-798.
- Ferraro R, Lillioja S, Fontvieille AM, Rising R, Bogardus C, et al. (1992) Lower sedentary metabolic rate in women compared with men. J Clin Invest 90: 780-784.
- Herrmann SD, Willis EA, Honas JJ, Lee J, Washburn RA, et al. (2015) Energy intake, nonexercise physical activity, and weight loss in responders and nonresponders: The Midwest Exercise Trial 2. Obesity (Silver Spring) 23: 1539-1549.
- 32. Alang N, Kelly CR (2015) Weight gain after fecal microbiota transplantation. Open Forum Infect Dis 2: ofv004.
- Cronin O, Molloy MG, Shanahan F (2016) Exercise, fitness, and the gut. Curr Opin Gastroenterol 32: 67-73.
- 34. Cook MD, Allen JM, Pence BD, Wallig MA, Gaskins HR, et al. (2016) Exercise and gut immune function: evidence of alterations in colon immune cell homeostasis and microbiome characteristics with exercise training. Immunol Cell Biol 94: 158-163.
- Campbell SC, Wisniewski PJ, Noji M, McGuinness LR, Häggblom MM, et al. (2016) The effect of diet and exercise on intestinal integrity and microbial diversity in mice. PLoS One 11: e0150502.
- 36. Recordati G (2003) A thermodynamic model of the sympathetic and parasympathetic nervous systems. Auton Neurosci 103: 1-12.
- Messina G, De Luca V, Viggiano A, Ascione A, Iannaccone T, et al. (2013) Autonomic nervous system in the control of energy balance and body weight: personal contributions. Neurol Res Int 2013: 639280.
- Herring JL, Molé PA, Meredith CN, Stern JS (1992) Effect of suspending exercise training on resting metabolic rate in women. Med Sci Sports Exerc 24: 59-65.

- Hautala AJ, Mäkikallio TH, Kiviniemi A, Laukkanen RT, Nissilä S, et al. (2004) Heart rate dynamics after controlled training followed by a homebased exercise program. Eur J Appl Physiol 92: 289-297.
- 40. Pal R, Singh SN, Chatterjee A, Saha M (2014) Age-related changes in cardiovascular system, autonomic functions, and levels of BDNF of healthy active males: role of yogic practice. Age (Dordr) 36: 9683.
- Rosenwinkel ET, Bloomfield DM, Arwady MA, Goldsmith RL (2001) Exercise and autonomic function in health and cardiovascular disease. Cardiol Clin 19: 369-387.
- 42. Koenig J, Jarczok MN, Warth M, Ellis RJ, Bach C, et al. (2014) Body mass index is related to autonomic nervous system activity as measured by heart rate variability--a replication using short term measurements. J Nutr Health Aging 18: 300-302.
- 43. Baumert M, Brechtel L, Lock J, Hermsdorf M, Wolff R, et al. (2006) Heart rate variability, blood pressure variability, and baroreflex sensitivity in overtrained athletes. Clin J Sport Med 16: 412-417.
- 44. Iellamo F, Legramante JM, Pigozzi F, Spataro A, Norbiato G, et al. (2002) Conversion from vagal to sympathetic predominance with strenuous training in high-performance world class athletes. Circulation 105: 2719-2724.
- 45. Schäfer D, Gjerdalen GF, Solberg EE, Khokhlova M, Badtieva V, et al. (2015) Sex differences in heart rate variability: a longitudinal study in international elite cross-country skiers. Eur J Appl Physiol 115: 2107-2114.
- 46. Lean ME, Malkova D (2016) Altered gut and adipose tissue hormones in overweight and obese individuals: cause or consequence? Int J Obes (Lond) 40: 622-632.