

Guidelines-Recommended Blood Pressure Goals and the Incidence of Cardiovascular and Renal Events

Niti Taylor*

Department of Hypertension, Eastern University, St Davids, PA 19087, USA

Introduction

Essential or primary hypertension, the world's top risk factor for disease burden, is predicted to account for more than half of the estimated 17 million deaths caused by Cardiovascular Disease (CVD) each year. Hypertension, defined as a rise in Blood Pressure (BP) above 140/90 mm Hg, is significantly linked to negative outcomes such as stroke, ischemic heart disease, heart failure and end-stage renal disease. The challenges of controlling hypertension and preventing the development of these latter effects are unlikely to abate anytime soon; by 2025, the global burden of hypertension is expected to rise by 60%, affecting nearly 1.6 billion adults. We focus on this highly widespread illness in this month's subject review in circulation: cardiovascular quality and outcomes.

Description

Overcoming the burden of hypertension still has a long way to go. First, due to the insidious nature of hypertension's development, it is commonly undetected, making early detection prior to the onset of end-organ damage difficult. Second, many individuals who have been correctly diagnosed with hypertension fail to meet the treatment goals set forth in guidelines. This underscores the significant difficulty of long-term risk factor management and adequate antihypertensive therapy adherence. Third, the optimal BP therapy target for high-risk patients is yet unknown. While a target blood pressure of 140/90 mm Hg is generally suggested, people at high risk of CVD, such as those with diabetes, should have a lower target of 130/80 mm Hg. It's unclear whether such aggressive blood pressure lowering leads to better outcomes. Fourth, even receiving appropriate therapy, a percentage of individuals remain resistant to treatment, despite taking various drugs. These patients with resistant hypertension are at a high risk of developing complications. Renal artery de-innervation could be the start of a new and effective treatment option for these patients [1-3]

The issues raised are relevant to a wide range of people. In the context of limited healthcare resources, many low- and middle-income nations, the majority of which are in the midst of the epidemiological transition, the incidence of hypertension is fast increasing. The rate of hypertension diagnosis and treatment in these nations is alarmingly low. As a result, developing novel and cost-effective technologies to enhance hypertension diagnosis and control remains a top focus. Despite good access to treatment, a plethora of information surrounding lifestyle modification and the presence of very efficacious anti-hypertensive medicines, less than half of patients in the United States have optimal blood pressure control. Indeed, on-going deficits

have inspired national initiatives like the Healthy People 2020 campaign and the Million Hearts initiative, which aim to improve awareness, treatment, and, ultimately, outcomes of this common condition [4,5]

Conclusion

When blood pressure surpasses 140/90 mm Hg or 130/80 mm Hg in people at high risk of Cardiovascular (CV) events, guidelines prescribe therapy. However, whether this lower treatment threshold in high-risk patients results in fewer CV events is debatable, with a few trials finding little benefit or perhaps damage from excessive BP lowering. With a median follow-up of 56 months, the on-going Telmisartan Alone and in Combination with Ramipril Global Endpoint Trial (ONTARGET) compared telmisartan with telmisartan + ramipril for the management of hypertension.

Acknowledgement

None.

Conflict of Interest

None

References

1. Forouzanfar, Mohammad H, Patrick Liu, Gregory A. Roth and Marie Ng et al. "Global burden of hypertension and systolic blood pressure of at least 110 to 115 mm Hg, 1990-2015." *JAMA* 317 (2017): 165-182.
2. Klag, Michael J., Paul K. Whelton, Bryan L. Randall and James D. Neaton et al. "Blood pressure and end-stage renal disease in men." *N Engl J Med* 334 (1996): 13-18.
3. Gnatiuc, Louisa, William G. Herrington, Jim Halsey and Jaakko Tuomilehto, et al. "Sex-specific relevance of diabetes to occlusive vascular and other mortality: A collaborative meta-analysis of individual data from 980 793 adults from 68 prospective studies." *Lancet Diabetes Endocrinol* 6 (2018): 538-546.
4. Rapsomaniki, Eleni, Adam Timmis, Julie George and Mar Pujades-Rodriguez, et al. "Blood pressure and incidence of twelve cardiovascular diseases: Lifetime risks, healthy life-years lost and age-specific associations in 1·25 million people." *Lancet* 383 (2014): 1899-1911.
5. Stamler, Jeremiah, Rose Stamler and James D. Neaton. "Blood pressure, systolic and diastolic and cardiovascular risks: US population data." *Arch Intern Med* 153 (1993): 598-615.

*Address for Correspondence: Niti Taylor, Department of Hypertension, Eastern University, St Davids, PA 19087, USA; E-mail: taylor9852@yahoo.com

Copyright: © 2022 Taylor N. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Received: 23 December, 2022; Manuscript No. jhoa-23-87815; Editor Assigned: 26 December, 2022, PreQC No. P-87815; Reviewed: 09 January, 2023, QC No. Q-87815; Revised: 16 January, 2023, Manuscript No. R-87815; Published: 24 January, 2023, DOI: 10.37421/2167-1095.2023.12.386

How to cite this article: Taylor, Niti. "Guidelines-Recommended Blood Pressure Goals and the Incidence of Cardiovascular and Renal Events." *J Hypertens* 12 (2023): 386.