

Malone Procedure Combined with Mitrofanoff Using a Divided Appendix and Vesicular Enlargement Description of the Surgical Technique and Case Report

Jose P*, Justino LC, José Manuel GG, Marco AVG, Heriberto CS, Roberto Antonio García Vasquez, and Julissa BA

Secretaría de Salud de Mexico, Hermosillo, Sonora, Mexico

Abstract

Surgical treatment for neurogenic bladder has evolved notably. A great number of patients with neurogenic bladder also present fecal incontinence and treatment for these patients is discouraging. With the induction and use of the Malone procedure through appendicostomy, the practitioner may offer the confection of a stoma for coecal catheterization, allowing easy access for daily enemas. The objective of this report is to present one surgical technique, where the coecal appendix was divided on the proximal portion created a stoma for antegrade enema (Malone principle) and the distal portion created a stoma for urinary catheterization in an augmented bladder with an intestinal segment (Mitrofanoff principle).

Introduction

The function of the inferior urinary tract depends on the neuromuscular coordination of the bladder, the urethra and the striated muscle responsible for the sphincter mechanism. A neurological or muscular disorder in these structures causes dysfunction. Void. The neurogenic bladder resulting from a neurological control disorder of continence and urination includes all problems of micturition whose origin is in the central nervous system [1]. The primary cause of dysfunctional neurogenic bladder in children is occult spina bifida; however, it may also be produced by sacral agenesis, tumors and inflammatory diseases affecting the spinal cord or spinal cord trauma [2]. It is essential to carry out urodynamic and imaging studies to determine the function of the high and low urinary tract in order to establish a treatment plan for each patient, and then to carry out an adequate follow-up. In the neurogenic bladder, the urodynamic findings basically determine: dysinergic bladder, synergic bladder and bladder with absence of activity [3]. The management of children with neurogenic bladders must be comprehensive, multidisciplinary and has three main objectives: Preserve renal function, seek urinary and fecal continence, and ensure that the patient becomes a self-sufficient and useful person in society [4]. The treatment seeks to ensure that the patient reaches urinary continence and that the upper urinary tract remains protected from the progressive alterations that occur as a result of increased pressures in the lower urinary tract. To achieve this, interventions must be performed to convert a bladder of low capacity and/or high pressure into a reservoir with adequate capacity at a low pressure, allowing an easy and complete evacuation as well as a hypotonic sphincter in a competent mechanism That does not allow the involuntary exit of urine. The ability to spontaneously urinate through the urethra should also be preserved, which is more difficult to achieve. Management of the neurogenic bladder should begin as soon as the diagnosis is made in order to prevent progressive deterioration of the upper urinary tract. The usual surgical procedures are: bladder augmentation, continual urinary stomas, including Mitrofanoff (coecal appendix reimplanted from the bladder to the abdominal wall) and the Monti tube (tubularization of a segment of the intestine and artificial sphincters, as well as urethral sling, submucosal injection of the bladder neck, reconstruction of the bladder neck and urethral cerclage to improve continence [5]. Surgical treatment can be complemented with pharmacological therapy, which consists of the administration of anticholinergic drugs that produce detrusor relaxation. The majority of patients with neurogenic bladder present alterations in colonic and anorectal function that cause fecal incontinence. Therefore, in some patients, procedures are performed to

allow antegrade enemas, such as the Malone technique [6]. The main complication that occurs with the stomates is the stenosis for which different techniques have been designed that use flaps to avoid it or to correct it.

Case Report

This is a 14-year-old male patient, the first-born of a mother in her first pregnancy, who had a course of urinary tract infection with multiple antibiotic management, with term pregnancy at birth diagnosed with myelomeningocele and Arnold Chiari type II. He was corrected for myelomeningocele and peritoneal ventricle bypass. In treatment from birth by orthopedics by bilateral equinovaro foot. An open reduction of hip joint was performed, as well as surgery of the medullary cone by ulceration of the filum terminale. Patient with neurogenic bladder that had been managed as a medium-risk bladder with clean intermittent catheterization, antibiotic prophylaxis and anticholinergic, begins its urological evaluation by our service with the following complementary studies: bilateral renal US in which bilateral hydro nephrosis is seen being more important on the right side with a degree III hydro nephrosis (Figure 1). The left side with grade I hydro nephrosis (Figure 2). When documenting hydronephrosis, we determined to perform a voiding urethrociogram (Figure 3) in which we observed the classic bladder in a Christmas tree, with irregular walls, with diverticular formations and absence of vesicoureteral reflux. It complements the assessment with a urodynamic study (Figure 4) which reflects a decreased bladder capacity of 140 ml (29%), without unconstrained contractions, with a leakage point of 107 ml. Bladder with pressure increase up to 60 cmH₂O with decreased compliance, an accommodation of 5 ml/cm.sup.20 and an average filling pressure of 38 cmH₂O. With the results of the studies already mentioned, it is possible to catalog as a high-risk

*Corresponding author: Jose Padilla, Secretaría de Salud de Mexico, Hermosillo, Sonora, Mexico, Tel: +525550621600; E-mail: drpadillajose@gmail.com

Received February 03, 2017; Accepted April 13, 2017; Published April 20, 2017

Citation: Jose P, Justino LC, José Manuel GG, Marco AVG, Heriberto CS, et al. (2017) Malone Procedure Combined with Mitrofanoff Using a Divided Appendix and Vesicular Enlargement Description of the Surgical Technique and Case Report. J Clin Case Rep 7: 952. doi: [10.4172/2165-7920.1000952](https://doi.org/10.4172/2165-7920.1000952)

Copyright: © 2017 Jose P, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Figure 1: US Right renal with grade III hydronephrosis.




Figure 2: Left renal ultrasound where mild ectasia is seen.

bladder, thus being a candidate for bladder enlargement with continent catheterization. He is offered to perform at the same time the procedure of Malone, due to the antecedent of constipation present in this case.

Preoperative laboratory studies were performed obtaining Hb 13.2 H to. 36.2 Platelets 320000 TP 13.1 TPT 30.3 Creatinine 0.9.

The surgical technique that was developed was the following: an umbilical mid incision was made, the abdominal cavity was entered and

the bladder was separated from the abdominal cavity; Approximately 15 cm of colonic segment was taken, resection was performed with whole-integer anastomosis in two colon-colonic planes and bladder enlargement was performed with this segment with meso (Figure 5). The appendix was removed, which was approximately 15 cm in length (Figure 6). The mesoappendix was divided, half of the appendix was used to perform the procedure of Malone that went towards the right iliac fossa; The other segment of the appendix went to the umbilical scar



Figure 3: Urethrocytogram.




Figure 4: Urodynamic study showing a neurogenic bladder.




Figure 5: Illustration of colono-colonic anastomosis and appendectomy performed. 1) Sigmoidocystoplasty, 2) Malone stoma with catheter, 3) Mitrofanoff stomata, 4) Mesoappendix, 5) Ileocecal valve, 6) End-to-end intestinal anastomosis, 7) Ascending colon, 8) Iliac Cup, 9) Left ureter, 10) Cystostomy.




Figure 6: Photo showing the appendicular size to be used for the catheterized ducts 15 cm.




Figure 7: Photo showing the separate appendix, one part was used for the Mitrofanoff and the other segment for the Malone.




Figure 8: Making the final Mitrofanoff bag.

and the Mitrofanoff was made into the bladder (Figure 7). The final results of the surgical procedure were: 100 mL bleeding, surgical time 210 minutes, 14-day in-hospital stay.

The patient was exited with silicon probes by mitroffanof stents (Figure 8) and functional malone, the drains were removed until the expenditure was less than 50 ml and the probes were removed at 21 postoperative days.

Discussion

The final outcome of the Mitrofanoff technique can be evaluated in terms of urinary continence, absence of complications of the stoma, surgical reinterventions due to complications and protection of the upper urinary tract. In general, it can be said that the results regarding continence are good, although in the series with longer follow-up there is a high frequency of surgical examinations of the duct, by filtration and stenosis according to what was published by Arango et al. [7].

In 1990 Malone et al. first described the continent-wide antegrade enema (EAC) technique for the treatment of severe fecal incontinence of neurogenic origin in children and has subsequently been successfully used also in adults and in patients with constipation who do not respond to medical treatment. The procedure consists of the formation of a continent duct, generally an appendectomy that allows intermittent catheterization for irrigation and the rapid and controlled cleaning of the colon described by Elsenberg [8]. This technique combines the principles of antegrade lavage of the colon and the catheterizable antireflux duct described by Mitrofanoff for the treatment of the neurogenic bladder. The goal is to achieve a complete cleaning of the colon to prevent fecal impaction and to avoid the episodes of incontinence described by Monti [9]. In the original description, the appendix was sectioned at its base and vascularization was preserved. From the moment this principle was exposed, the appendix was constituted as the tissue of choice for the development of the duct, in this case the Malone surgery was performed to improve fecal continence with periodic lavage through the continent stoma. The choice of the type of canal to be used depends on several factors including availability, type of surgery and basic diagnosis according to Landero et al. [10]. The defect in the cecal wall was closed and the appendix 180° was rotated, with reimplantation in one of the blind's tapeworms through a 5 cm long submucosal tunnel to avoid the reflux described by Landero and Gladman [10,11]. Modifications of the original technique have been described in recent years, which in some cases simplify the procedure. In our patient, the surgical techniques of Mitrofanoff and Malone were performed, with a long coecal appendix of 15 cm, divided into two segments, as reported by Kajbafzadeh et al. being the first report in the literature [12]. They concluded that the appendix greater than 9 cm is ideal to develop this surgical technique in conjunction with V-flap technique for the stoma and to reduce its main complication than stoma stenosis.

Conclusion

The technique of Malone should be taken into account in patients with severe constipation and fecal impaction, since it facilitates regular colon cleansing and reduces encopresis, which significantly limits the quality of life of the patient.

Successful treatment of urinary and fecal incontinence secondary to neuropathic disorders can improve the quality of life of children, achieving greater socialization and well-being. The development of the catheterizable vesicocutaneous duct technique described by Mitrofanoff has been the main contribution in achieving adequate control of urinary continence and has been of great importance for periodic bladder emptying.

The management of the neurogenic bladder requires a complete assessment of the patient's conditions, basic diagnosis and associated pathologies; This will depend on the prognosis of our patients.

Surgical treatment of the bladder with neurogenic bladder prevents renal damage and generates advantages in terms of continence, comfort and quality of life.

References

1. Malone PS, Ransley PG, Kiely EM (1990) Preliminary report: The antegrade continence enema. *Lancet* 336: 1217-1218.
2. Bica D (1996) Apendicectomia ortotopica continente, 5 casos. *Rev de Cir Infantil* 4.
3. Orabi A, Paterson HM, Goncette L, Danse E, Saey JP (2011) Malone appendicostomy: A an unexpected complication: A case report. *Tech Coloproctol* 15: 81-83.
4. Esteves E (1999) Simultaneous surgical treatment of neurogenic bladder and fecal incontinence in children. *Rev by Cir Infantil* 2.
5. Traxel E, Minevich EA, Noh PH (2010) A review: The application of minimally invasive surgery to pediatric urology: Lower urinary tract reconstructive procedures. *Urology* 76: 115-121.
6. Pera M (2006) Treatment of severe chronic constipation by means of the continental antegrade enema technique. *Cir Esp* 80: 403-405.
7. Arango M (2009) Results of the use of the Mitrofanoff technique in patients with neurogenic bladder dysfunction: Analysis of the experience at the Hospital Universitario San Vicente de Paul. *Urological Acts Spanish* 33: 69-75.
8. Elsenberg L (2012) Pilot Study of the vesicocutaneous continent catheterizable stoma (Mitrofanoff) in adults-high complication rates. *Urologys* 79: 222-226.
9. Cain M (2008) Updated experience with the monti catheterizable channel. *Urology* 72: 782-785.
10. Landero (Malone technique for the continent's antegrade enema in children with intestinal and urinary neurogenic disease *Rev Mex Urol* 69: 268-272. Gladman M, Knowels CH (2008) Surgical treatment of patients with constipation and fecal incontinence. *Gastroenterol Clin N Am* 605-625.
11. Kajbafzadeh M (2001) Simultaneous Malone antegrade continent enema and Mitrofanoff principle using dividen appendix: Report of a new technique for prevention of stoma complication. *Journal of Urology* 2404-2409.