

Open Access

Middle-Late Pleistocene Paleo-Climate and Paleo-Altimetry of the Centre of Tibetan Plateau Indicated by the Sporopollen Record of Well QZ-4

Jianglin He^{1,2*}, Jian Wang^{1,2}, Weipeng Li^{3,4} and Wei Sun^{1,2}

¹Chengdu Institute of Geology and Mineral Resources, Ministry of Land Resources, Chengdu 610082, PR China

²Key Laboratory for Sedimentary Basin and Oil and Gas Resources, Ministry of Land Resources, Chengdu 610081, PR China

³Land and Resources College, China West Normal University, Nanchong, 637002, PR China

⁴Key Laboratory of Mountain Environment Evolution and Regulation, Institute of Mountain Hazards and Environment, Chinese Academy of Sciences, Chengdu 610041, PR China

Abstract

The core sample from well QZ-4 is an important climate archive for the central Tibetan Plateau in the middle-late Pleistocene. In this work, a detailed pollen analysis of it is carried out to provide a preliminary insight into the paleoclimate and paleo-altimetry change in the central Tibetan Plateau. It can be concluded that the pollen assemblage can be obviously divided into two pollen zones, Pollen zone I (251.1-314 m in depth, 120.0-345.8 ka BP) and Pollen zone II (200-251.1 m in depth, 105.4-120 ka BP). The paleo-climate during pollen zone I deposition period was comparatively colder and wetter than it was during the pollen zone II deposition period. After Gonghe movement, the center of Tibetan Plateau was uplifted about 300 m (from 3500-3700 m to 3800-4000 m in elevation). The wind was changed from horizontal or downward direction to upward direction, in the study area. In the central of Tibetan Plateau, the climate change seems to be mainly driven by global climate change, and that tectonic uplift may have been a subordinate influence at the middle-late Pleistocene.

Keywords: Tibet Plateau; Mid-pleistocene; Palaeoclimate; Paleoaltimetry; Gonghe movement

Introduction

The Tibetan Plateau is not only the largest and highest mountain Plateau on Earth, but it is also the site of the first successful terrestrial gas hydrate exploration in China [1]. In recent years, considerable attention has been focused on the study of the correlation between the uplift of the Tibetan Plateau and quaternary climate evolution [2], which is not only beneficial to the study of global climate change, but it is also beneficial for the study of permafrost surveys and the exploration of gas hydrates in the Tibetan Plateau. However, the existing results come from the surrounding area [3,4], and little study in the central Tibetan Plateau has been reported (Figure 1). Previous studies have shown that, globally, the hottest period since the middle Pleistocene was at approximately 120 ka BP, which is recorded in the ice sheet above Lake Vostok in the Southern hemisphere [5] and in Greenland [6] in the Northern hemisphere. North Greenland Eemian Ice Drilling (NEEM) surface temperatures after the onset of the Eemian (126 ka) peaked at $8 \pm 4^{\circ}$ C above the mean of the past millennium [6]. In the central Tibetan Plateau, the thermoluminescence (TL) dating of well QD-1 [7], the ¹⁴C and ²³⁰Th/²³⁴U dating of well TS-95 [8,9] and the Electronic Spin Resonance (ESR) dating of well QZ-4 show that the transition section of the colour sequence in the three wells was deposited at approximately 120 ka (Figure 2). Furthermore, the last rapid uplift of the Tibetan Plateau started at 150 ka BP, which is referred to as the Gonghe movement [10]. After the Gonghe movement (Figure 3), the average elevation reached almost 4,000 m, at which the East Asian monsoon was apparently affected by the Tibetan Plateau [10,11]. The paleoclimate study based on the cores of well QZ-4 will benefit the study of the correlation between climate changes in the central Tibetan Plateau and global climate change in the middle-late Pleistocene, which will contribute to the study of the correlation between the uplift of the Tibetan Plateau and quaternary climate evolution.

In this work, the thickest quaternary sedimentary formation in the central Tibetan Plateau was drilled in well QZ-4 (Figure 1), and the most complete core was collected. The cores are characterized by a colour sequence with three colour features: (1) the yellow subsequence (with dark yellowish-brown and brownish-red colour) at the depth from 272.48 m to 314 m, (2) the grey subsequence (with dark grey and whitish-grey colour) at the depth from 198 m to 240 m, and (3) the transition section (with light greyish-green colour) interbedded with the brown and grey clay at the depth from 240 m to 272.48 m, which is consistent with the colour sequences observed in wells QD-1 [7] and TS-95 [9], which are located in the central and Northwestern Tibetan Plateau, respectively (Figure 2). The cores drilled in well QZ-4 are desirable archives for the study of climate change in the central Tibetan Plateau. In this paper, we present a detailed pollen analysis of the well QZ-4 in an effort to provide preliminary insight into the climate and the elevation change recorded in it. The data of the MIS curve are come from the website: https://commons.wikimedia.org/wiki/File:Vostok_Petit_data.svg?uselang=zh-sg

Sample and Methods

Well QZ-4 is located in the central Tibetan Plateau, approximately 50 km East of Shuanghu County at 89°14′E and 33°5′N, with an elevation of 4932 m. The surrounding area is a flat terrain, in which the maximum elevation difference is smaller than 50 m. The rivers in this area are seasonal rivers, which are formed during the rainy season and flow West into Dirangbicuo Lake and Caiduochaka Lake (Figure 1). The bedrock around the QZ-4 well is the Tumengela Formation, which is composed of quartz-lithic sandstone and was deposited in a deltaic floodplain environment in the Late Triassic. The total depth of this well is 314 m. The cores in the well were well collected at 190-314 m depth, partly

*Corresponding author: Jianglin He, Chengdu Institute of Geology and Mineral Resources, Key Laboratory for Sedimentary Basin and Oil and Gas Resources, Ministry of Land Resources, Chengdu 610081, PR China, Tel: +86 02883231771; E-mail: 5hjl998@163.com

Received July 07, 2017; Accepted July 13, 2017; Published July 18, 2017

Citation: He J, Wang J, Weipeng Li, Sun W (2017) Middle-Late Pleistocene Paleo-Climate and Paleo-Altimetry of the Centre of Tibetan Plateau Indicated by the Sporopollen Record of Well QZ-4. J Environ Anal Toxicol 7: 490. doi: 10.4172/2161-0525.1000490

Copyright: © 2017 He J, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

J Environ Anal Toxicol, an open access journal ISSN: 2161-0525

Page 2 of 8





Figure 2: The lithological association and the ESR, TL and ¹⁴C and ²³⁰Th /²³⁴U dating results of this color sequence in well QZ-4, well QD-1 [7] and well TS-95 [8,9] with depth scale. The core images of this color sequence in well QZ-4.

J Environ Anal Toxicol, an open access journal ISSN: 2161-0525

collected at 0-110 m depth, and absent at 110-190 m depth. Basing on the core samples, core recovery, detritus and indicators of sedimentary environments (e.g., colour, lithologic characters, grain size, grain shape, particle roundness, sedimentary structures, and fossil assemblage) in the core samples, the stratigraphic sequence drilled in well QZ-4 can be subdivided into two sedimentary sequences:

The first sequence, at 0-198 m in depth, is a regressive depositional sequence, which consists of semi-deep lake facies, shallow lake facies, lake shore facies, meandering stream facies and flood plain from the deep to shallow layers in sequence. These facies are characterized by black grey clay and grey clay, light grey silty clay, light grey, yellow sand, and a yellow-brown sandy gravel layer, respectively. The second sequence, at 198-314 m depth, consists of several transgressiveregressive depositional sequences, which consist of shallow lake facies, lake shore facies and semi-deep lake facies. In this sequence, the semideep lake facies is characterized by black grey clay, grey clay, dark yellowish-brown clay and brownish-red clay, with horizontal bedding at depths of 198-199.9 m, 203.18-221.68 m, 227.18-240.8 m, 247.88-260.78 m and 281.33-293.98 m. The shallow lake facies is characterized by yellow-brown silty clay interlayered with thin light-grey silty clay, brown clay, and white grey clay at depths of 199.9-203.18 m, 22168-227.18 m, 240.8-247.88 m, 260.78-278.31 m, 293.98-296.18 m and 302.4-313.0 m. The lake shore facies is characterized by yellow-brown sand interlayered with white grey sand and brown clay at depths of 278.31-281.33 m, 296.18-301.7 m and 313.0-314 m (Figure 2).

A total of 20 samples with 5 m intervals were collected from the well QZ-4 for palynological analysis. Samples weights of 200 g were used to prepare pollen residues for the upper 244 m depth, and samples weights of 300 g were used to the lower depths. Samples were mostly prepared using standard techniques [12], involving alkali digestion, treatment with 10% cold HCl, 10% hot KOH, 46% hot HF, and Erdtman's acetolysis, staining with safranin, dehydration with tertiary butyl alcohol, and mounting in silicone oil. Prior to alkali digestion, moss polsters and soils were dispersed in distilled water and sieved (200 mm) to remove coarse detritus and sand. Lycopodium tablets were added to the samples to allow for estimation of pollen concentrations [13]. All samples have been studied under microscopes at Nanjing Institute of Geology and Palaeontology, Academic Sinica (Nanjing, China) and The Institute of Hydrogeology and Environmental Geology, Chinese Academy of Geological Sciences (Shijiazhuang, China). Leica microscope at a magnification of 400 was used for pollen identification and counting. More than 100 pollen grains were counted for each sample. Pollen abundance was expressed as percentages calculated using the total pollen sum.

Results

Most of the studied samples yielded rich, well-preserved palynomorph assemblages, with the identified sporomorphs representing about 75 taxa, including arboreal taxa,

e.g. Pinus, Picea, Picea, Tsuga, Cupressaceae, Betula, Carpinus, Alnus, Juglans, Quercus, Fagus, Ailanthus, Ulmus, Tilia, Moraceae, Salix, Corylus, Ilex, Oleaceae, as well as scrubby and herbaceous taxa, e.g. Lonicera, Sambucus, Rosa, Spiraea, Rhododendron, Caragana, Reaumuria, Hippophae, Nitraria, Ephedra, Gramineae, Chenopodiaceae, Compositae, Chrysanthemum, Artemisia, Aster, Xanthium, Ranunculaceae, Ranunculus, Thalictrum, Polygonum, Rumex, Labiatae, Prunella, Euphorbia, Rosaceae, Potentilla, Sauguisorba, Papilionatae, Primula, Rubiaceae, Scrophulariaceae, Lythraceae, Cruciferae, Araliaceae, Geranium, Clerodendron, Gentiana, Solanaceae, Sedum, Umbelliferae, Saxifraga, Lilium, Arenaria, Macleaya, Urticaceae, Cyperaceae, Typha, and some fern spores, including Lycopodium, Sellaginella, Adiantaceae Pteris, Athyrium, Polypodiaceae, Polypodium, Lepisorus, Filicale. Some algal remains (e.g. Pediastrum) were also detected. The total spore and pollen sum was used to calculate the percentage of various spore and pollen taxa. As shown in Figure 3, the pollen assemblage can obviously divided into two pollen zones:

Pollen zone I (251.1-314 m in depth, 120.0-345.8 ka BP) is characterised by single species and low pollen concentrations. The average pollen concentration is around 0.4 grain/g, except the sample at 295.0 m in depth with 3.1 grain/g. In this pollen zone, the pollen species are particularly characterized by high abundance of tree, a few herb, shrubs and fern spores. The maximum proportion of tree pollen reaches 86.5%, with an average of 66.2%. The tree pollen principally includes Pinus (10.5-71.9%, av. 45.31%), Picea (5.6-29.0%, av. 17.4%) (Figure 3). Tsuga, Cupressaceae, Betula, Quercus etc. are occasionally seen in some samples. The average content of herb pollen is 19.2%. The herbs principally include Ephedra (0.9-18.9%, av. 7.0%), Chenopodiaceae (1.1-13.9%, av. 5.4%), Cyperaceae (0.2-6.8%, av. 2.1%) (Figure 3). Gramineae, Artemisia, Labiatae, Rosaceae, Typha etc. are occasionally seen in some samples, especially the Typha composes 25.9% of the total pollen sum at 295.0 m in depth. The maximum proportion of shrub pollen reaches 46.4%, with an average of 9.7%. The shrub principally includes Nitraria (0.5-12.7%, av. 4.2%) (Figure 3). Lonicera, Rosa etc. are occasionally seen in some samples, especially the Lonicera composes 38.8% of the total pollen sum at 256.9 m in depth. The fern spores are sporadically present in a few samples. The maximum proportion of fern spores is smaller than 0.9%.

Pollen zone II (200-251.1 m in depth, 105.4-120 ka BP) is characterised by multiple species and high pollen concentrations. The average pollen concentration is around 2.8 grain/g. In this pollen zone, the pollen species are particularly characterized by a high abundance of herb, a few tree, shrubs and fern spores. The maximum proportion of herb pollen reaches 80.3%, with an average of 60.9%. The herb principally includes Thalictrum (4.2-50.5%, av. 28.8%), Chenopodiaceae (0.5-19.5%, av. 6.8%), Ephedra (0.7-12.9%, av. 4.7%), Gramineae (1.3-5.4%, av. 3.1%) (Figure 3). Artemisia, Ranunculus, Labiatae, Rosaceae etc. are occasionally seen in some samples. The maximum proportion of tree pollen reaches 61.9%, with an average of 24.5%. The trees principally are Pinus (0.5-35.6%, av. 16.11%) (Figure 3). Picea, Tsuga, Cupressaceae, Betula, Quercus etc. are occasionally seen in some samples. The maximum proportion of shrub pollen reaches 24.1%, with an average of 13.6%. The shrub principally include Nitraria (0.5-21.4%, av. 9.0%), Rosa (0.4-8.6%, av. 2.9%) (Figure 3). Lonicera, Spiraea, Hippophae, Reaumuria etc. are occasionally seen in some samples. The maximum proportion of fern spores reaches 1.7%, with an average of 1.0%. The fern spores principally is Athyrium (0.2-0.8%, av. 0.4%), Pteris, Polypodium, Lepisorus etc. are sporadically present in a few samples.

Discussion

Ecologic environment

The sporopollen assemblage of pollen zone I is characterized by high percentages of arboreal taxa, e.g. *Pinus, Picea* and *Abies*, and low percentages of scrubby and herbaceous taxa. However, the sporopollen assemblage of pollen zone II is characterized by high percentages of shrub species and herbaceous taxa, e.g. *Thalictrum, Rosaceae* and *Labiatae*, and low percentages of arboreal taxa (Figure 3). Many studies of the representation of Picea/Abies-pollen in modern





sporopollen assemblages indicate that such assemblages are highly indicative of autochthonous vegetation [14,15]. In pollen zone I, the average percentage of *Picea* and *Abies* is 17.4% (5.6-29.0%), which is suggesting that spruce and fir forests developed widely in the central of Tibetan Plateau. On the Tibetan Plateau today, *Picea* and *Abies* are the main representatives of sub-alpine dark coniferous forests and are principally distributed Gongga Mountain, Himalaya, Nyainqentanglha and Hengduan Mountains [14,16]. However, in the pollen zone II, *Picea* and *Abies* are occasionally seen in some samples, with average percentage of 0.8% (0-4.9%). The pollen zone II is mainly composed of shrub species and herbaceous taxa, such as *Labiatae* [16], *Thalictrum* [16] and *Rosaceae* which are the major herb species of alpine shrub meadow in Tibetan Plateau.

In brief, during pollen zone I deposition period, the zonal vegetation in the central of Tibetan Plateau should have been subalpine dark coniferous forests. However, it has been alpine shrub meadow during pollen zone II deposition period.

Paleo-temperature interpretation

The study of Cordova *et al.* [17] shows that *Quercus-Ulmus-Poaceae* pollen assemblage zone and the development of a chernozem soil suggest cool-dry climate conditions. In well QZ-4, the *Quercus-Ulmus-Poaceae* pollen assemblage is observed in the pollen zone II, and the chernozem soil section is observed in this zone too (Figure 3). Meanwhile, it is known that *Pinus* prefers cool weather [18]. In the well QZ-4, the *Pinus* is observed in the both pollen zones. It is indicated that both the two pollen zones are deposited in the cool temperature regime. On the other hand, *Chenopodiaceae, Cyperaceae* and *Lamiaceae* are suggesting dry and cold conditions [19]. In pollen zone II, *Chenopodiaceae, Cyperaceae* and *Lamiaceae* are richer than them in the pollen zone I. Meanwhile, the modern mean annual temperature is generally above 4°C in the

sub-alpine dark coniferous forests [14] and the is no higher than -1°C in the alpine shrub meadow in Tibetan Plateau [20]. Hence, it can be concluded that both pollen zone I and II are deposited in a cool climate environment. The climate is comparatively cooler during the pollen zone II depositional period than the pollen zone I depositional period. In the central of Tibetan Plateau, the temperature was slightly declined in the middle-late Pleistocene.

Paleo-humidity interpretation

The study of Ji et al. [21] shows that the *Thalictrum* is normally in arid climates. In well QZ-4, *Thalictrum* is rich in pollen zone II (4.2-50.5%, av. 28.8%), but absent in the in the pollen zone I, which is indicating that regional climate is comparatively drier during the pollen zone II depositional period than the depositional period of pollen zone I.

Ephedra is typical desert plants and is often used as indicator of dry climate [22]. It is a perennial shrub and it is a very effective sandbinder. In Saudi Arabia, it is associated with sand dunes formation, especially the mobile, non-saline and low moisture content ones [23]. In the pollen zone I, the *Ephedra* principally is rich in the sand strata and absent in the clay strata. However, in pollen zone II, it is obviously richer in clay strata than sand strata (Figure 3). It is indicated that the *Ephedra* is principally associated with sand dunes formation in the pollen zone I. Nevertheless, in pollen zone II, the climate is so dry that the *Ephedra* is principally associated with the moisture. The regional climate is comparatively drier during the pollen zone I.

Nitraria is one of dominant species in desertified steppe and desert [24]. *Gramineae* is better represented in dry than in moist forest [25], and it is taken as dry climate [26]. *Chenopodiaceae* characterize dry environments [22,27]. *Rosaceae* and *Rosa* are suitable for cool-dry conditions [28]. The *Nitraria, Chenopodiaceae, Gramineae, Rosaceae*

and *Rosa* are enriched in pollen zone II (Figure 3), which are indicated that regional climate is comparatively drier during the pollen zone II depositional period than the depositional period of pollen zone I. In brief, climate is comparatively drier during the pollen zone II depositional period than the depositional period of pollen zone I. In the central of Tibetan Plateau, the climate was dried in the middle-late Pleistocene.

Paleo-altimetry interpretation

Traditionally, air temperature decline with increasing elevation in free air (lapse rate) is considered the primary factor determining the position and composition of altitude-related vegetation zones [29]. The approaches basing on the correlation of temperature and altitude are applied to the reconstruction of paleo-altimetry [14], such as isotopes [30], atmospheric pressure, enthalpy, pCO₂, fossils, the δD of leafwaxes and sporopollen assemblages. However, the practice shows that surface temperatures do not depend simply on elevation. For example, there is no trees growing in the area higher than 4500 m in the central of Tibet, such as the mean annual temperature is about -0.9°C in the Bange county where the elevation is about 4500 m (Figure 4). However, multiple species of plant grow in the cold region such as the Mohe country in the Northeast of China. The mean annual temperature is at about -5.5°C in Mohe country where the elevation is only about 400 m (Figure 4). Song et al. [29] suggested that palaeo-elevation can be deduced by enthalpy which is a combination of both temperature and humidity at a constant pressure. In fact, the air pressure is declining with increasing of elevation. The quantitative study of paleo-altimetry would be much more complex than existing research. Up to now, in the study area, there have been no quantitative temperature data (such as U_{a7}^{k} and TEX86), humidity data and air pressure data in middlelate Pleistocene. In present, the analogy analysis is likely more credible



Figure 4: The vegetation distribution is not depend simply on elevation, such as the present vegetation landscape in Mohe country is obviously different from it in the Bange country.

than quantitative estimation of palaeo-elevation which is based on the temperature, humidity and air pressure.

Fossil pollen is almost ubiquitous and this makes it attractive as the basis of a possible palaeoaltimeter. Based on their presumed Nearest Living Relatives, the climate and altitude can be studied through their taxonomic affinity and assumptions about past environmental tolerances [14,29,31]. In the Eastern Tibetan Plateau, the distribution of modern pollen assemblages had been well study by Li et al. [16], Zhang et al. [32] and Cheng and Luo [33] in the altitudinal transect from 1100 to 4500 m (Table 1). Based on previous work, the paleo-elevation of the pollen zone can be semi-quantitatively estimated by its pollen assemblages respectively (Figure 5).

Pollen zone I is dominated by *Pinus* (38.6%, 1200-4200 m), *Picea* and Abies (16.8%, 2700-3700 m), *Ephedra* (13.1%, 2400-5300 m), *Lonicera* (10.8%, 1200-5200 m), *Chenopodiaceae* (7.3%, 2000-5500 m), *Gramineae* (2.1%, 650-5600 m), *Thalictrum* (1.9%, 2800-4000 m), *Cyperaceae* (1.5%, 3700-4500 m), *Cupressaceae* (1.3%, 3700-4500 m), *Rosa* (1.1%, 860-5600 m) and *Potentilla* (1.0%, 3700-4500 m). It is showing that the pollen zone I was mainly came from the belt in the elevation of 3500-3700 m. Meanwhile, the pollen zone II is dominated by *Thalictrum* (33.9%), *Pinus* (19.5%), *Chenopodiaceae* (8.1%), *Ephedra* (5.5%), *Betula* (4.4%), *Gramineae* (3.6%), *Labiatae* (3.4%), *Rosaceae* (3.4%), *Cyperaceae* (3.4%), *Potentilla* (1.2%), *Ulmus* (1.2%), *Lonicera* (1.0%) and *Rosa* (1.0%). It is showing that the pollen zone II mainly came from the belt in the elevation of 3800-4000 m.

In the Eastern Tibetan Plateau, the subalpine dark coniferous forests are distributed in the belt range from 2800 to 3700 m above the sea level (a.s.l.) of Gongga Mountain [16]. The frigid dark coniferous forests are distributed in the belt range from 2800 to 3600 m a.s.l. of Gongga Mountain. Mountain dark coniferous forests are mainly distributed in the Northern slopes of some mountains at an elevation of about 3100-3500 m a.s.l. in the Southeast and 2800-3800 m a.s.l. in the Northwest of Tibetan Plateau [34-39]. As mention above, the pollen zone I is mainly deposited in the subalpine dark coniferous forests environment. Hence, it is reasonable that the pollen zone I was mainly came from the belt in the elevation from 3500 to 3700 m.



Figure 5: Plot of altitude ranges of presumed nearest living relatives of pollen taxa recovered from the Eastern Tibetan Plateau. The pollen percentages of the pollen zone I and II is colored with orange and grey respectively. The distribution belt of the pollen zone I and II is expressed with the yellow and grey horizontal bars respectively.

	Pollen/Spore	Content in Pollen zone II%	Content in Pollen zone I%	Credible upper limit (m)	Maximum reported (m)	Minimum reported (m)	Credible lower limit (m)	References
1	Pinus	19.54	38.63	4200	4200	1200	1200	[29]
2	Picea/Abies	0.97	16.76	3700	4200	2300	2700	[16,29]
3	Tsuga	0.52	0.22	3200	3200	2800	2800	[29]
4	Cupressaceae	0.48	1.32	4500	5300	1700	3700	[16,29]
5	Betula/Carpinus	4.41	0.62	3700	2800	780	2800	[16,29]
6	Alnus	0.38	0.18	2800	2800	780	1900	[29]
7	Juglans	0.66	0.13	2700	3300	1300	1900	[29]
8	Quercus	0.68	0.92	2400	4000	1900	1900	[16,29]
9	Ulmus	1.24	0.04	2800	2800	2100	2100	[29]
10	Tilia	0.02	0.00	2900	2900	700	700	[16,29]
11	Salix	0.05	0.09	4000	5100	2300	2800	[16,29]
12	Corylus	0.43	0.04	2800	2800	2600	2600	[29]
13	llex	0.00	0.13	3600	3600	900	900	[29]
14	Oleaceae	0.07	0.00	4000	4000	860	860	[29]
15	Lonicera	0.93	10.78	3800	5200	1200	1500	[29,34]
16	Rosa	3.28	1.10	5600	5600	860	860	[29]
17	Spiraea	0.43	0.09	4800	4800	3800	3800	[29]
18	Rhododendron	0.05	0.00	4500	5490	450	3700	[16,29]
19	Hippophae	0.48	0.13	5200	5200	1000	1000	[29]
20	Ephedra	5.47	13.07	5300	5300	2400	3500	[29,35,36]
21	Gramineae	3.57	2.11	5600	5600	650	3800	[29,37,38]
22	Chenopodiaceae	8.10	7.26	5500	5500	2000	2000	[29]
23	Artemisia	0.43	0.31	1900	5600	1400	1400	[29]
24	Thalictrum	33.88	1.85	4000	5200	2000	2800	[16,29]
25	Polygonum	0.36	0.00	5100	5100	900	900	[29]
26	Labiatae	3.37	0.26	3700	5500	600	1400	[29]
27	Rosaceae	3.39	0.44	860	860	560	560	[29]
28	Potentilla	1.29	0.92	4500	5600	2100	3700	[16,29]
29	Gentiana	0.50	0.04	5700	5700	2100	2100	[29]
30	Solanaceae	0.50	0.04	5200	5200	780	780	[29]
31	Sedum	0.34	0.18	5400	5400	1000	1000	[29]
32	Umbelliferae	0.00	0.13	5300	5300	800	800	[29]
33	Lilium	0.77	0.70	5400	5400	800	800	[29]
34	Cyperaceae	3.42	1.50	4500	5600	800	3700	[12,16,29]

Table 1: Modern altitudinal ranges of taxa and the percentage content of taxa in each pollen zone.



The alpine shrub meadows are distributed at an elevation of about 3700-4000 m a.s.l. in Gongga Mountain [16], 3800-4800 m a.s.l. in the Southern slope of mid-Himalaya Mountains [40], above 3800 m in the Eastern margin of the Tibetan Plateau [41]. As mention above, the pollen zone I is mainly deposited in the subalpine dark coniferous forests environment. Hence, it is reasonable that the pollen zone I was mainly came from the belt in the elevation of 3800-4000 m. The study of

the Zhou et al. [42] is shows that the elevation of the Plateau surface was raised about 300 m, from 4270 to 4570 m after the Gonghe Movement. Although the absolute altitude is inconsistence with our study, the rising height of the plateau surface is consistence with ours [42]. Meanwhile, our study is consistence with the paleo-elevation inferred from oxygen isotope of lacustrine deposits in the Kunlun Mountain Pass too. In the Pliocene period, the paleo-elevation was 3426 m from oxygen isotope of marlstone sample collected at 94°03'58"E and 35°38'18"N [43]. In the Pleistocene period, the paleo-elevation was 3773 m from oxygen isotope of marlstone sample collected at 94°05'24"E and 35°38'42"N [43]. The paleo-elevation was 3769 m from oxygen isotope of marlstone sample collected at 94°05'44"E and 35°38'51"N [43]. The elevation of Kunlun Mountain Pass was raised about 340 m (from 3430 to 3770 m) during the period from Pliocene to Pleistocene [43]. The modern elevation of Kunlun Mountain Pass is approximately 4780 m a.s.l. The modern elevation of the study area is approximately 4932 m, which is approximately 150 m higher than the Kunlun Mountain Pass. Hence, it is reasonable that the elevation of study area was raised about 300 m (from 3500-3700 m to 3800-4000 m) during the middle-late Pleistocene.

The wind direction interpretation

Usually, air temperature is declining with the increasing elevation. As shown above, the elevation of study area was raised about 300 m

J Environ Anal Toxicol, an open access journal ISSN: 2161-0525

(from 3500-3700 m to 3800-4000 m) during the middle-late Pleistocene. It seems to be contrary to the paleo-temperature interpretation that the climate is comparatively cooler during the pollen zone I depositional period than the pollen zone II depositional period. However, in the pollen zone I, part of the pollen (Cupressaceae, Gramineae, Cyperaceae and Potentilla) is come from higher altitude, but little is come from the lower altitude localities (Figure 5). It is indicated that the study area was located in the advection wind or down wind direction. In the pollen zone II, the pollen species are not only more numerous than them in Pollen zone I, but also the pollen assemblages in it are more complex. A little pollen (Potentilla) is coming from higher altitude localities, while a lot of pollen is come from the lower altitude localities (Tsuga, Alnus, Juglans, Quercus, Ulmus, Salix, Artemisia, Rosaceae) (Figure 5). It is indicated that the study area was located in the upwind direction. It can be concluded that the wind direction in the central of Tibetan Plateau was changed from horizontal or downward direction to upward direction at middle-late Pleistocene. During the pollen zone II depositional period, the pollen may be transported on to the high and relatively cold Plateau from lower altitude warm temperate forests by upwelling wind (summer monsoon) [29], which may be resulted the characteristics that the climate is comparatively cooler during the pollen zone I depositional period than the pollen zone II depositional period.

Influence of the Gonghe movement

It is known that the hottest period since the middle-Pleistocene was at approximately 120 ka BP both in the Southern hemisphere [44,45] and Northern hemisphere [6]. Meanwhile the Gonghe movement was started at 150 ka BP and persisted for about 50 ka (Figure 6) [10,11]. Hence, many studies regard the Plateau uplift as a possible driving force for the simultaneous enhancement of East Asian winter and summer monsoons as well as for the Mid-Pleistocene transition occurrence [46,47]. However, some authors suggest that the mid-Pliocene climate changes in East Asia are unlikely to be a response to Plateau uplift [48].

As shown above, the elevation of study area was raised about 300 m (from 3500-3700 m to 3800-4000 m) during the middle-late Pleistocene. As the elevation increased, the temperature would be decreased about 1.8°C calculated with an air temperature gradient of 6 K/km. However, the MIS (marine isotope stage) curve shows that the annual temperature raised about 8°C after the Gonghe movement (Figure 3). In the central of Tibetan Plateau, the temperature was slightly elevated in the middle-late Pleistocene. The climate change in the central of Tibetan Plateau was corresponding with the global climate change at the middle-late Pleistocene.

In another face, the paleo-climate during pollen zone I deposition period was comparatively wetter than it during the pollen zone II, which is shown that the barrier effect of Tibetan Plateau to moisture is stronger and stronger with the elevation increasing of Tibetan Plateau. Hence, it can be concluded that the climate change in the central of Tibetan Plateau at the middle-late Pleistocene documented by the well QZ-4 core seems to be mainly driven by global climate change, and that tectonic uplift may have been a subordinate influence.

Conclusions

The comprehensive paleo-climate and paleo-altimetry studies of the pollen samples from the well QZ-4, in the centre of Tibetan Plateau, lead to the following conclusions:

1) In the well QZ-4, the pollen assemblage can obviously divided into two pollen zones. Pollen zone I (251.1-314 m in depth, 120.0-345.8 ka BP) is characterised by single species and low pollen concentrations.

Pollen zone II (200-251.1 m in depth, 105.4-120 ka BP) is characterised by multiple species and high pollen concentrations.

2) The paleo-climate during pollen zone I was comparatively colder and wetter than it during the pollen zone II.

3) The palaeo-elevation of pollen zone I is concentrated on the belt in the elevation from 3500 to 3700 m. The palaeo-elevation of pollen zone II is mainly come from the belt in the elevation from 3800 to 4000 m. The study area had been uplifted about 300 m after the Gonghe Movement.

4) In the centre of Tibetan Plateau, the wind had changed from horizontal or downwelling wind to upwelling wind, in the middle-late Pleistocene.

5) In the central of Tibetan Plateau, the climate change seems to be mainly driven by global climate change, and that tectonic uplift may have been a subordinate influence in the middle-late Pleistocene.

Acknowledgements

This study was supported by the National Natural Science Foundation of China Nos. 41302092. Grateful acknowledgements are made to the Xizang Geothermal Geological Brigade, which provided the field help during the drilling process. Guobang Tong and Songmei Tong, coming from Institute of Hydrogeology and Environmental Geology, Chinese Academy of Geological Sciences, provided assistance for the identification of pollen. Jinhui Cheng, coming from Nanjing Institute of Geology and Palaeontology, Academic Sinica gave much help during sample preparation.

Highlights

- I. The center of Tibetan Plateau uplifted about 300 m after Gonghe movement
- II. Barrier effect of Tibetan Plateau to moisture was enhanced after Gonghe movement
- III. Wind changed from horizontal or down welling to upwelling after Gonghe movement
- $\operatorname{IV}.$ Paleo-climate change in Tibetan Plateau is mainly driven by global climate change

References

- He J, Wang J, Fu X, Zheng C, Chen Y (2012) Assessing the conditions favorable for the occurrence of gas hydrate in the Tuonamu area Qiangtang basin, Qinghai–Tibetan, China. Energy convers Manage 53: 11-18.
- Zhang D, Tingyun YA (2015) Environmental impacts of hydraulic fracturing in shale gas development in the United States. Petroleum Explor Develop 42: 876-883.
- Hoorn C, Straathof J, Abels HA, Xu Y, Utescher T, et al. (2012) A late Eocene palynological record of climate change and Tibetan Plateau uplift (Xining Basin, China). Palaeogeography Palaeoclimatol Palaeoecol 344: 16-38.
- Xiao G, Guo Z, Dupont-Nivet G, Lu H, Wu N, et al. (2012) Evidence for northeastern Tibetan Plateau uplift between 25 and 20Ma in the sedimentary archive of the Xining Basin, Northwestern China. Earth Planetary Sci Lett 317: 185-195.
- Yamamoto K, Yamashita F, Adachi M (2005) Precise determination of REE for sedimentary reference rocks issued by the Geological Survey of Japan. Geochem J 39: 289-297.
- Dahl-Jensen D, Albert MR, Aldahan A, Azuma N, Balslev-Clausen D, et al. (2013) Eemian interglacial reconstructed from a Greenland folded ice core. Nature 493: 489-494.
- 7. Yi H, Zhu X, Zhu Y (2006) Lake level change recorded by core of the Quaternary lacustrine sediment in the central Tibetan plateau and its climatic implications. Earth Sci Front p: 300-307.
- Shijie L, Hongliang Z, Yafeng S (2008) A high resolution MIS 3 environmental change record derived from lacustrine depositi of Tianshuihai lake, Qinghai-Tibet Plateau. Quaternary Sci 28: 122-31.
- 9. Yu SH, Zhu ZY, Li SJ, Li B, Zhou H, Sun Y (1997) Environmental records of

Page 8 of 8

variation of iron oxides in drill core from Tianshuihai lake on the southern flanke of west Kunlun mountains. Geochemica 26: 88-98.

- 10. Jijun L (1991) The environmental effects of the uplift of the Qinghai-Xizang Plateau. Quaternary Sci Rev 10: 479-483.
- Qiao Y, Qi L, Liu Z, Wang Y, Yao H, et al. (2014) Intensification of aridity in the eastern margin of the Tibetan Plateau since 300ka BP inferred from loess-soil sequences, western Sichuan Province, southwest China. Palaeogeography Palaeoclimatol Palaeoecol 414: 192-199.
- Lu H, Wu N, Liu KB, Zhu L, Yang X, et al. (2011) Modern pollen distributions in Qinghai-Tibetan Plateau and the development of transfer functions for reconstructing Holocene environmental changes. Quaternary Sci Rev 30: 947-966.
- Zhao K, Li X, Dodson J, Atahan P, Zhou X, et al. (2012) Climatic variations over the last 4000calyr BP in the western margin of the Tarim Basin, Xinjiang, reconstructed from pollen data. Palaeogeography Palaeoclimatol Palaeoecol 321: 16-23.
- Wu F, Herrmann M, Fang X (2014) Early Pliocene paleo-altimetry of the Zanda Basin indicated by a sporopollen record. Palaeogeography Palaeoclimatol Palaeoecol 412: 261-268.
- Xiao X, Shen J, Wang S (2011) Spatial variation of modern pollen from surface lake sediments in Yunnan and southwestern Sichuan Province, China. Rev Palaeobotany Palynol 165: 224-234.
- 16. Li Q, Ge Q, Tong G (2012) Modern pollen-vegetation relationship based on discriminant analysis across an altitudinal transect on Gongga Mountain, eastern Tibetan Plateau. Chinese Sci Bull 4: 1-9.
- Cordova CE (2007) Holocene Mediterranization of the southern Crimean vegetation: Paleoecological records, regional climate change, and possible non-climatic influences. NATO Science Series IV–Earth and Environmental Sciences, edited by: Yanko-Hombach, V., Gilbert, A., Panin, N., and Dolukhanov, P., Springer, Berlin, New York, USA, p: 319-344.
- 18. Good RE, Good NF (1975) Growth characteristics of two populations of Pinus rigida Mill. from the Pine Barrens of New Jersey. Ecology 56: 1215-1220.
- Jaimand K, Nejad SK, Monfared A, Akbarzadeh M (2014) Study of the chemical composition of essential oil of Teucrium chamaedrys at the different distillation in Mazandaran province. J Med Plants By-Prod 3: 193-198.
- 20. Li Y, Sun X, Zhao X, Zhao L, Xu S, et al. (2006) Seasonal variations and mechanism for environmental control of NEE of CO2 concerning the Potentilla fruticosa in alpine shrub meadow of Qinghai-Tibet Plateau. Science in China Series D: Earth Sci 49: 174-185.
- Ji S, Xingqi L, Sumin W, Matsumoto R (2005) Palaeoclimatic changes in the Qinghai Lake area during the last 18,000 years. Quaternary Int 136: 131-140.
- 22. Zhao Y, Yu Z, Chen F, Liu X, Ito E (2008) Sensitive response of desert vegetation to moisture change based on a near-annual resolution pollen record from Gahai Lake in the Qaidam Basin, northwest China. Global Planetary Change 62: 107-114.
- Wafa'a A, Al-Qarawi AA, Alsubiee MS (2010) Effect of water stress by Polyethylene Glycol 8000 and Sodium Chloride on germination of Ephedra alata Decne seeds. Saudi J Biol Sci 17: 253-257.
- 24. Zhang J, Wu B, Zhu Y, Li Y, Lu Q, et al. (2013) Responses of Nitraria tangutorum to water and photosynthetic physiology in rain enrichment scenario. Acta Ecologica Sinica 33: 172-177.
- 25. Bullock SH (1995) Seasonally dry tropical forests. Cambridge University Press, UK.
- Fujii R, Sakai H (2002) Paleoclimatic changes during the last 2.5 myr recorded in the Kathmandu Basin, Central Nepal Himalayas. J Asian Earth Sci 20: 255-266.
- Herzschuh U (2007) Reliability of pollen ratios for environmental reconstructions on the Tibetan Plateau. J Biogeography 34: 1265-1273.
- Chuanhu ZH, Xing CH (2010) Climatological significance of Pollen factors revealed by pollen-climate response surface functions in Dajiuhu, Shennongjia. J Meteorol Res 24: 699-712.

- 29. Song XY, Spicer RA, Yang J, Yao YF, Li CS (2010) Pollen evidence for an Eccene to Miocene elevation of central southern Tibet predating the rise of the High Himalaya. Palaeogeography Palaeoclimatol Palaeoecol 297: 159-168.
- Xu Q, Ding L, Zhang L, Cai F, Lai Q, et al. (2013) Paleogene high elevations in the Qiangtang Terrane, central Tibetan Plateau. Earth Planetary Sci Lett 362: 31-42.
- Sun J, Xu Q, Liu W, Zhang Z, Xue L, et al. (2014) Palynological evidence for the latest Oligocene- early Miocene paleoelevation estimate in the Lunpola Basin, central Tibet. Palaeogeography Palaeoclimatol Palaeoecol 399: 21-30.
- 32. Xianhao Z, Wenjing Z, Ji L (1999) The characteristics of the mountain ecosystem and environment in the Gongga mountain region. Ambio.
- Cheng G, Luo J (2001) Successional features and dynamic simulation of subalpine forest in the Gongga Mountain, China. Acta Ecologica Sinica 22: 1049-1056.
- 34. Viraj M, Verma RK, Chauhan NS, Kapoor KS (2012) Assessment of floristic diversity in Phulang valley of Lippa-Asrang Wildlife Sanctuary of district Kinnaur, Himachal Pradesh. Environ Ecol 30: 226-233.
- 35. Maher LJ (1964) Ephedra pollen in sediments of the Great Lakes region. Ecology 45: 391-395.
- Liu N, Wang Q, Chen J, Zhu Y, Tashi T, et al. (2010) Adaptive evolution and structure modeling of rbcL gene in Ephedra. Chinese Sci Bull 55: 2341-2346.
- 37. Liu SL, Zhao HD, Su XK, Deng L, Dong SK, et al. (2015) Spatio-temporal variability in rangeland conditions associated with climate change in the Altun Mountain National Nature Reserve on the Qinghai-Tibet Plateau over the past 15 years. Rangeland J 37: 67-75.
- Deng T, Li Y (2005) Vegetational ecotype of the Gyirong Basin in Tibet, China and its response in stable carbon isotopes of mammal tooth enamel. Chinese Sci Bull 50: 1225-1229.
- Lu H, Wu N, Gu Z, Guo Z, Wang L, et al. (2004) Distribution of carbon isotope composition of modern soils on the Qinghai-Tibetan Plateau. Biogeochemistry 70: 275-299.
- 40. Lu C (2014) Spatial distributing characteristics of land use in the southern slope of mid-Himalaya Mountains. InLand Surface Remote Sensing II p: 9260.
- 41. Zhou W, YU SY, Burr GS, Kukla GJ, Jull AJ, et al. (2010) Postglacial changes in the Asian summer monsoon system: a pollen record from the eastern margin of the Tibetan Plateau. Boreas 39: 528-539.
- Zhou S, Wang X, Wang J, Xu L (2006) A preliminary study on timing of the oldest Pleistocene glaciation in Qinghai–Tibetan Plateau. Quaternary Int 154: 44-51.
- 43. Wu ZH, Zhao X, Ye PS, Wu ZH, Hu DG, et al. (2007) Paleo-elevation of the Tibetan Plateau inferred from carbon and oxygen isotopes of lacustrine deposits. Acta Geol Sin 81: 1277-1288.
- 44. Vimeux F, Cuffey KM, Jouzel J (2002) New insights into Southern Hemisphere temperature changes from Vostok ice cores using deuterium excess correction. Earth Planetary Sci Lett 203: 829-843.
- 45. Yamamoto Y, Kitahara N, Kano M (2012) Long memory effect of past climate change in Vostok ice core records. Thermochimica Acta 532: 41-44.
- 46. Liu D, Fang X, Song C, Dai S, Zhang T, et al. (2010) Stratigraphic and paleomagnetic evidence of mid-Pleistocene rapid deformation and uplift of the NE Tibetan Plateau. Tectonophysics 486: 108-119.
- 47. Han W, Fang X, Berger A (2012) Tibet forcing of mid-Pleistocene synchronous enhancement of East Asian winter and summer monsoons revealed by Chinese loess record. Quaternary Res 78: 174-184.
- 48. Ge J, Dai Y, Zhang Z, Zhao D, Li Q, et al. (2013) Major changes in East Asian climate in the mid-Pliocene: Triggered by the uplift of the Tibetan Plateau or global cooling? J Asian Earth Sci 69: 48-59.