

Mucocele of the Rectum

Shivakumar SP* and Shanmugam RP

Department of Surgical Gastroenterology, Kumaran Hospital Pvt. Ltd., No. 214, EVR Periyar Salai, Kilpauk, Chennai - 600 010, India

Abstract

Mucocele of the rectum causing intestinal obstruction is a rare complication of retained long rectal stump. We present one such case.

Keywords: Mucocele of rectum; Long rectal stump; Intestinal obstruction

Introduction

The mucosa of rectum secretes mucus, which must drain out otherwise it will accumulate over time. Injury, scarring and stenosis of the anal canal in a proximally closed rectum can cause mucocele, which progressively enlarges in size.

Case Report

A 45 yr old male presented with vomiting, low grade fever, diffuse pain & distension of abdomen with constipation of one week duration. History of road traffic accident in October 1991 for which a sigmoid end colostomy with closure of distal bowel loop with supra pubic cystostomy done as emergency surgical procedure for pelvic fracture with damage to lower rectum & anal canal including sphincter.

On examination the patient was febrile 100 F, Tachypneic pulse rate: 96/min & mildly dehydrated. Abdomen was distended & tense. Colostomy was seen in the left iliac fossa and SPC catheter in the supra pubic area. Bowel sounds were absent. Colostomy examination was normal. Examination of perineum showed complete stenosis of the anal canal.

Plain X ray abdomen showed dilated large bowel loops & no bowel gas seen in the pelvic area. Ultra sound abdomen revealed grossly dilated fluid filled bowel loops suggestive of sub acute intestinal obstruction (Figure 1). Blood routine was normal except low serum potassium (2.9 mEq /lit).




Figure 1: Ultra sound abdomen revealed grossly dilated fluid filled bowel loops suggestive of sub acute intestinal obstruction.

The patient was initially managed conservatively for two days with nil per oral, IV fluids, correction of hypokalemia, Ryle's tube aspiration & enema through the colostomy. Since the pain and distension increased, laparotomy was decided. Through left Para median incision abdomen opened. A huge cystic mass arises from the pelvis to the supra umbilical region seen and small bowel adherent to the cystic swelling. The cystic swelling was compressing on the colostomy loop causing obstruction. On mobilizing the swelling it was found to be the long rectal stump that was closed during the earlier procedure.

The inter bowel adhesions released. 800 ml of mucous fluid aspirated from the cystic swelling & sent for culture & sensitivity. The swelling shrunk completely relieving the pressure on the colostomy site. The end was brought out as a mucous fistula on the right iliac fossa. Post operative period was uneventful.

Discussion

The mucosal lining of the rectum secretes mucus, which must drain otherwise it will accumulate over time. Inflammation, scarring, epithelialization & stenosis with obstruction of the anal canal can produce a mucocele, which can increase in size enormously.

Mucocele are commonly associated with the appendix, retained esophagus, gall bladder neck obstruction and cranial sinuses. Rectal mucoceles are rare. The first reported pediatric case was an extraluminal pelvic mucocele in a patient with an imperforate anus following an endorectal pull-through operation with incomplete mucosal excision [1]. There are case reports of rectal mucoceles following Hartmann's procedure [2] for ulcerative colitis, and secondary to high anal sphincter tone following spinal trauma [2,3]. Mucoceles have developed in defunctioned colon after pull-through surgery for Hirschsprungs disease [5] due to stenosis and retraction of the colonic stump [5,6]. Scarring of the mucus fistula, or failure to create a mucus fistula are other described contributing factors. Treatment requires effective drainage or removal of the mucus-producing epithelium.

References

1. Hitch DC, Patil UB, Panicek DM (1987) Mucocele after endorectal pull-through for imperforate anus. J Pediatr Surg 22: 1023-1024.

*Corresponding author: Dr. S.P. Shivakumar, Kumaran Hospital Pvt. Ltd., No. 214, E.V.R. Periyar Salai, Kilpauk, Chennai - 600 010, India, E-mail: drshiv74@hotmail.com

Received May 05, 2012; Accepted July 14, 2012; Published July 16, 2012

Citation: Shivakumar SP, Shanmugam RP (2012) Mucocele of the Rectum. J Clin Case Rep 2:170. doi:10.4172/2165-7920.1000170

Copyright: © 2012 Shivakumar SP, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

-
2. Creagh MF, Chan TY (1991) Case report: rectal mucocoele following Hartmann's procedure. Clin Radiol 43: 358-359.
 3. Elsmann BH, Van Vroonhoven TJ (1995) Pelvic mucocoele. A late complication after treatment of ulcerative colitis: case report Eur J Surg 161: 373-374.
 4. Hsu WM, Wang W, Chen CC (2000) Enteric mucocoele formation after endorectal pull-through: report of a case. J Formos Med Assoc 99: 650-652.
 5. Witte JT, Harms BA (1989) Giant colonic mucocoele after diversion colostomy for ulcerative colitis. Surgery 106: 571-574.