

## NT Pro-BNP can be used as a Risk Predictor of Clinical Atrial Fibrillation with or without Left Atrial Enlargement

Zhao X<sup>1</sup>, Li H<sup>2</sup>, Liu C<sup>1</sup>, Ren Y<sup>3</sup> and Sun C<sup>4\*</sup>

<sup>1</sup>Health Science Center, Xi'an Jiaotong University, Shaanxi Province, P.R. China

<sup>2</sup>Department of Rehabilitation and Treatment, The First Affiliated Hospital of Xi'an Jiaotong University, Shaanxi Province, P.R. China

<sup>3</sup>Medical Science Center, Yan'an University, Shaanxi Province, P.R. China

<sup>4</sup>Cardiovascular Department, The First Affiliated Hospital of Xi'an Jiaotong University, Shaanxi Province, P.R. China

\*Corresponding author: Sun C, Doctoral Supervisor, Cardiovascular Department, The First Affiliated Hospital of Xi'an Jiaotong University, Shaanxi Province, P.R. China, Tel: + 8618191273192; E-mail: chaofengsxjtu@163.com

Received: May 17, 2019; Accepted: May 29, 2019; Published: June 06, 2019

Copyright: © 2019 Zhao X, et al. This is an open-access article distributed under the terms of the creative commons attribution license, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Abstract

**Background:** NT Pro-BNP is a blood marker secreted from myocardial. The main stimulus for cardiac NT Pro-BNP secretion is myocardial stretch. NT Pro-BNP is a significant risk factor for stroke of cardiac insufficiency and pulmonary embolism. Patients with normal levels of NT-pro-BNP have low risks for death. So does atrial stretch.

**Retraction note:** Publisher took the decision to retract the article from the journal.

**RETRACTED**