

Psychosocial Factors among Patients with Diabetes

Shiva Mani

Department of Medicine & Advanced Technology, Seoul National University Hospital, Seoul, Korea

Editorial

Diabetes is a growing problem that poses a significant public health challenge around the world. According to the International Diabetes Federation (IDF) and the World Health Organization (WHO), diabetes currently affects more than 8% of the global population (415–420 million people), with prevalence expected to rise to 10.4% (642 million) by 2040. Diabetes affects an estimated 9.3 percent of the population (29.1 million people) in the United States. Type-2 diabetes is the most common type of diabetes worldwide, accounting for 90 percent of cases [1].

Diabetes is the fourth or fifth leading cause of death in the majority of high-income countries, posing a significant burden on public health systems. Diabetes health spending accounted for 12% (USD 673 billion) of global health expenditure in 2015, and it may account for up to 20% of national health-care budgets in some countries. Furthermore, the indirect costs of diabetes, such as reduced labour force participation and lower economic productivity, are significant.

Psychosocial factors and diabetes risk

Emotional distress: Depression is the most commonly studied factor in diabetes research. According to the findings of two meta-analyses of longitudinal studies, depression is associated with a 37–60% increased risk of developing diabetes. Prospective evidence also suggests that higher levels of depressive symptoms, as well as clinical depression, are associated with an increased risk of diabetes [2]. After controlling for diabetes risk factors such as BMI, family history of diabetes, smoking, physical activity, diet, and alcohol consumption, the associations reported in these studies remained significant.

One possibility is that diabetes and depression share common etiological factors such as physical inactivity or inflammation, which statistical adjustments may not completely eliminate. Preclinical diabetes, on the other hand, may increase the likelihood that an individual will report depression, resulting in a reverse causal process [3].

Psychological distress is characterised by a number of comorbid psychological factors, including depressive and anxiety symptoms, general stress, and sleep disturbance. In a UK study of 9514 people, psychological distress at baseline was linked to incident diabetes after adjusting for age, gender, education, and income. However, after adjusting for health-related factors, the relationship was no longer significant.

Exposure to life stress: Chronic exposure to external stressors has also been linked to the onset of diabetes. Until now, the majority of research has focused on the link between work stress and incident diabetes. Job strain, defined as the combination of high job demands and limited job control, is a

well-studied work stress construct. A large meta-analysis that looked at the relationship between job stress and diabetes development gathered data from 13 prospective European cohort studies [4]. Over a 10.3-year average follow-up, job stress was associated with a 1.15-fold increased risk of incident diabetes. This association was found to be independent of a variety of covariates, and it extends previous pooled cross-sectional associations. A large number of studies on the link between long work hours and diabetes have also been conducted. It appears that working 55 h or more a week also increases the risk of developing diabetes but only in low socioeconomic status (SES) groups.

Early life adversity: Early life adversity has not been widely investigated as a risk factor for future diabetes onset, though it appears to be a significant issue for health-related processes such as telomere length and inflammation in adult life.

Personality characteristics: Personality factors in relation to diabetes have received little attention. Hostility is typically defined as a negative cynical attitude toward others, with a proclivity for anger or aggression [5]. This trait has been linked prospectively to elevated fasting glucose and cross-sectionally to insulin resistance, glycated haemoglobin (HbA1c), and diabetes prevalence.

Conflict of Interest

None.

References

1. Ramirez, Julio A, Timothy L Wiemken, Paula Peyrani, and Forest W Arnold, et al. "Adults Hospitalized With Pneumonia in the United States: Incidence, Epidemiology, and Mortality." *Clin Infect Dis* 65 (2017): 1806-1812.
2. Jain, Seema, Wesley H Self, Richard G Wunderink, and Sherene Fakhraian, et al. "Community-Acquired Pneumonia Requiring Hospitalization among U.S. Adults." *N Engl J Med* 373 (2015): 415-427.
3. Keys, HM, BN Bundy, FB Stehman, and LI Muderspach, et al. "Cisplatin, radiation, and adjuvant hysterectomy compared with radiation and adjuvant hysterectomy for bulky stage IB cervical carcinoma." *N Engl J Med* 340 (1999): 1154 -1161.
4. Rotman, Marvin, Alexander Sedlis, Marion R Piedmonte, and Brian Bundy, et al. "A phase III randomized trial of postoperative pelvic irradiation in Stage IB cervical carcinoma with poor prognostic features: follow-up of a gynecologic oncology group study." *Int J Radiat Oncol Biol Phys* 65 (2006): 169-176.
5. Kaye, Walter H, Guido K Frank, Ursula F Bailer, and Shannan E Henry, et al. "Serotonin alterations in anorexia and bulimia nervosa: new insights from imaging studies." *Physiol Behav* 85 (2005): 73-81.

***Address for Correspondence:** Shiva Mani, Department of Medicine & Advanced Technology, Seoul National University Hospital, Seoul, Korea, Tel: +9232716844; E-mail: Shivaman1939@gmail.com

Copyright: © 2022 Mani S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Received: 05 March, 2022, Manuscript No: jhmi-22-58005; **Editor Assigned:** 07 March, 2022, PreQC No: P-58005; **Reviewed:** 10 March, 2022, QC No: Q-58005; **Revised:** 15 March, 2022, Manuscript No: R-58005; **Published:** 20 March, 2022, DOI: 10.37421/jhmi.2022.13.408

How to cite this article: Mani, Shiva. "Psychosocial Factors among Patients with Diabetes." *J Health Med Informat* 13 (2022): 408.