

## Remote Primary Intraventricular Haemorrhage in Case of a Ruptured MCA Aneurysm

Redmond JG\* and Pattavilakom SA

Department of Neurosurgical, Princess Alexandra Hospital, Australia

### Abstract

This is the first documented case report of a simultaneous middle cerebral artery (MCA) aneurysmal subarachnoid hemorrhage (aSAH) and remote third ventricular clot. The occurrence of simultaneous subarachnoid hemorrhage and remote hypertensive intracerebral hemorrhage (ICH) is a rare occurrence and has only been reported three previous times in the literature. Intraventricular hemorrhage (IVH) post aneurysmal subarachnoid hemorrhage is common, where peri-ventricular intracerebral clot has ruptured the ependymal lining. However primary intraventricular hemorrhage (PIVH) without associated parenchymal clot is rare. Migration of blood via outlet foramina of the fourth ventricle through CSF pulsation and flow is thought to be the mechanism for this occurrence and would account for small volumes in the subacute setting. Here, we present a case of a 52-year-old woman with a CT demonstrated spontaneous middle cerebral artery (MCA) bifurcation aneurysmal subarachnoid hemorrhage and a large acute third ventricular clot spanning the foramen Monroe.

**Keywords:** Aneurysm; Cerebral hemorrhage; Hypertension; Subarachnoid haemorrhage

### Introduction

Intraventricular hemorrhage is associated with significant mortality and morbidity. The incidence of primary intraventricular hemorrhage (PIVH) without parenchymal clot is reported as 3.1% of all intracranial hemorrhage presentations. Most cases of IVH result from the extension of parenchymal hemorrhage or rupture of an aneurysm. Although intraventricular hemorrhage and its association with aneurysmal rupture is well described in the literature, remote hemorrhages are poorly differentiated and understood. The simultaneous occurrence of remote hypertensive intracerebral hemorrhage (ICH) following aneurysmal rupture has been published three previous times in the literature [1-3]. However, to our knowledge this is the first documented case of remote PIVH associated with aSAH. We performed a literature review of both primary intraventricular bleeds and other rare hemorrhages associated with aneurysmal subarachnoid hemorrhage (aSAH). This case serves to provide an opportunity to examine a CT demonstrated hemorrhage that led to a delayed diagnosis of an aneurysmal rupture (Figure 1).

### Case Report

#### History

A 52-year-old woman presented with an episode of generalized tonic-clonic seizure witnessed by her husband. The seizure lasted one minute and was associated fecal and urinary incontinence. Following a five-minute post-ictal period of unresponsiveness, she regained consciousness and remained confused. She had no significant past medical or surgical history, took no regular medications and was a non-smoker with no significant family history.

Initially, the patient was transported by ambulance to a non-neurosurgical hospital. Examination by an ED consultant revealed mild hypertension (BP 157/88) and otherwise normal vital signs (GCS 15, PEARL, HR 66 BP 157/88 RR 16 Sa 100% Temp 36.8). Her central nervous system and review of systems examinations were unremarkable.

The initial CT performed at the peripheral hospital showed acute subarachnoid hemorrhage with fresh blood tracking along the course of the left middle cerebral artery into the Sylvian fissure (Figures 2). A hyperdense circumscribed lesion anteriorly in the third ventricle measuring 16 mm × 13 mm × 12 mm in diameter was reported. This lesion appeared to be continuous with a further component present in



**Figure 1:** CT demonstrated hemorrhage that led to a delayed diagnosis of an aneurysmal rupture.



**Figure 2:** CT showed hemorrhage with fresh blood tracking along the course of the left middle cerebral artery into the Sylvian fissure.

the anterior horn of the left lateral ventricle 1.91 cm in diameter. The lesion was hyperdense posteriorly but isodense to the adjacent brain. No layering of blood was identified on the lateral or third ventricles.

**\*Corresponding author:** Joseph Garcia Redmond, Department of Neurosurgical, Princess Alexandra Hospital, Australia, Tel: +61 7 3176 2111; E-mail: [joevg36@hotmail.com](mailto:joevg36@hotmail.com)

**Received** February 07, 2018; **Accepted** March 29, 2018; **Published** March 31, 2018

**Citation:** Redmond JG, Pattavilakom SA (2018) Remote Primary Intraventricular Haemorrhage in Case of a Ruptured MCA Aneurysm. J Clin Case Rep 8: 1093. doi: 10.4172/2165-7920.10001093

**Copyright:** © 2018 Redmond JG, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

On arrival to the neurosurgical department, the patient was GCS 13 E3 V4 M6 PEARL. She was drowsy orientated to person, denied a headache and reported feeling sleepy. Her power was globally reduced to 4/5. She was begun on treatment for aSAH despite the negative CT angiogram. A repeat non-contrast CT brain showed an increase in the subarachnoid hemorrhage and new intraventricular extension into the posterior horns of both lateral ventricles (Figure 3). The apparent mass in the third ventricle seen on the afternoon CT was noted to have increased in size and density indicating that it could be a clot (Figure 4).

Formal angiogram found a multi-lobulated aneurysm projecting infero-laterally from the left MCA bifurcation, which approximated 6.5 mm × 4 mm. Two daughter sacs were evident. No other aneurysms or arteriovenous malformations (AVM) identified (Figure 5).

The patient received craniotomy and aneurysmal clipping on day two of her presentation and localized to the tube post-clipping. Post-operative CTA brain day three demonstrated the aneurysm was excluded from circulation. With no other occult aneurysms identified. The patient remained in ICU for six days. She remained a GCS 14 and MSQ 2-3 post-extubation. She remained shunt-dependent and received a ventriculoperitoneal shunt day twenty-one. She discharged on day

twenty-five with further out-patient rehabilitation. Two years on she has made full recovery return to driving with no major neurological deficits.

## Discussion

Primary intraventricular hemorrhage (PIVH) was first described by Sanders in 1881 as the flooding of the ventricle by blood without the presence of any rupture or laceration in the ventricular wall [4]. Venous angioma has been suggested to be the most likely cause when a parenchymal injury is not present [5]. Other studies have reported hypertension followed by vascular malformation as the most common risk factors for PIVH [1].

The incidence of IVH associated with aSAH varies from 10% [6] to 70% [7] depending on whether the parenchymal clot is included. Jabarli et al. reported an incidence of 23% IVH associated with MCA aneurysms (2016). The range in incidence also reflects the wide variation in mortality rates as noted by Rosen et al. This suggests that the etiology of a ventricular clot in the setting of aSAH may also be quite varied. In the presented CT images there is no continuity between the patient's subarachnoid hemorrhage (SAH) and the intraventricular clot. It is reasonable to assume that the bleeds are independent in origin. Despite an extensive literature review this appears to be the first case reported of a remote PIVH with aSAH.

Middle cerebral artery aneurysms have been more commonly associated with ICH while anterior communicating artery aneurysms and posterior circulation aneurysms with IVH [8]. Rarely aneurysmal rupture presents without SAH [8,9]. There are three case reports of remote ICH following aSAH. Hypertension is postulated to be the most reasonable cause as a reflex of the initial aneurysmal bleed or independent of the bleed [3]. The location of IVH without parenchymal clot has previously been reported not to correlate with either the clinical status of the patient or the nature, site or size of the bleeding lesion [6]. However, recent studies report increased mortality 64-65% following SAH with IVH associated with poor clinical status [10,11]. Patient age, location, and size of an aneurysm are described as predictors for development of IVH post aSAH [12]. Other studies reported a smaller size of aneurysm to be predictive of occurrence and severity of ICH [7]. This reflects the progression in our understanding of predictors for development of IVH/ICH in aSAH.

The presence of IVH in the setting of aSAH is a risk factor for the development of hydrocephalus [12]. Location of an aneurysm may also predict shunt dependency when IVH is not present [13]. Placement of an EVD is the procedure of choice for the treatment of acute hydrocephalus in patients with SAH [14]. This was performed bilaterally as a trapped ventricle was suspected. This patient had a Graeb score of 6 and while clot itself is a predictor of shunt dependency severity of or size of clot is not [12]. Jabbarli et al. found no impact of IVH severity on shunt dependency, while other recent studies report a Graeb score of 6 or greater to be significantly associated with the development of hydrocephalus [15]. The development of predictive scores may help with early assessment of shunt dependency [16].

A seizure occurs in approximately 26% of patients following aSAH. Early onset seizures are thought to negatively affect the grading of the aSAH. This subset of patients may have significantly improved recovery given the early seizure distorting the grading. Current opinion is that post resuscitation neurological status is the best predictor of outcome. However, patients that are WFNS grade I initially tend to do better than those patients that improve to a grade I [17,18].

Rupture of the lamina terminalis may be one cause of IVH in




Figure 3: CT brain showed an increase in the subarachnoid hemorrhage and new intraventricular extension.



Figure 4: CT was noted to have increased in size and density.



Figure 5: Angiogram brain.

aSAH. However, this is more often associated with an ACA/ACOM aneurysm where the membranous layer offers little resistance against bleeding forces. The ascent of blood products through CSF outlet foraminae via elastic recoil of brain tissue following cardiac systole has also been suggested as a reason for IVH [5,6]. Neither theory proffered adequately serves to explain the combination of the radiographic findings presented here. Certainly, the delay in observing the mass as a clot is reasonable given in the hyper-acute setting (first 4-6 h), which during this time, CT imaging does not reveal hemorrhage well, because it appears isodense to the brain [8]. In this case, the delay to radiological diagnosis did not halt pre-emptive aneurysmal treatment or the need for a further radiological investigation.

## Conclusion

This report illustrates a case where the location of hemorrhage and CT findings do not marry leading to delayed radiological diagnosis. Whether IVH represents an isolated clot or passage of blood via CSF spaces is difficult to delineate in the literature and may represent different etiologies. The link between aSAH and IVH, is described in the literature, however, it is difficult to find examples of studies that reflect the presentation or course of this patient. There are no previous reports of a link between PIVH and aSAH. The presentation of this case aims to show that while IVH in aSAH is known that the etiology behind the IVH may be varied. It is our hypothesis that this case is representative of how reflexive hypertensive hemorrhages may occur in aSAH and that these hemorrhages may occur in varied locations. Whether this itself impacts patient presentation and outcome will require further investigation.

## References

1. Bulgular K, Ev S (2009) Spontaneous primary intraventricular hemorrhage in adults: Clinical data, etiology and outcome. *Turkish Neurosurgery* 19: 338-344.
2. Rosen DS, Macdonald RL, Huo D, Goldenberg FD, Novakovic RL, et al. (2007) Intraventricular hemorrhage from ruptured aneurysm: clinical characteristics, complications, and outcomes in a large, prospective, multicenter study population. *J Neurosurg* 107: 261-265.
3. Lee JK, Lee HK, Kim IY, Kim ST, Jung S, et al. (2002) Simultaneous Occurrence of subarachnoid hemorrhage due to ruptured aneurysm and remote hypertensive intracerebral hemorrhage. *J Korean Med Sci* 17: 144-146.
4. Sanders E (1881) A study of primary, immediate, or direct hemorrhage into the ventricles of the brain. *Am J Med Sci* 82: 85-128.
5. Donauer JRE, Khalaf B, Menedoht EF, Faubert (1993) Intraventricular haemorrhage caused by aneurysms and angiomas. *Acta Neurochirurgica* 122: 23-31.
6. West CGH, Forbes WC (1985) Intraventricular blood without parenchymal clot following spontaneous subarachnoid haemorrhage. *Neuroradiology* 27: 254-258.
7. Kramer AH, Mikolaenko I, Deis N, Dumont AS, Kassell NF, et al. (2010) Intraventricular hemorrhage volume predicts poor outcomes but not delayed ischemic neurological deficits among patients with ruptured cerebral aneurysms. *Neurosurgery* 67: 1044-1052.
8. Thai QA, Raza SM, Pradilla G, Tamargo RJ (2005) Aneurysmal rupture without subarachnoid hemorrhage: Case series and literature review. *Neurosurgery* 57: 225-229.
9. Song TW, Kim SH, Jung SH, Kim TS, Joo SP (2016) Rupture of distal anterior cerebral artery aneurysm presenting only subdural hemorrhage without subarachnoid hemorrhage: a case report. *Springer Plus* 5: 73.
10. Mohr G, Khan M, Malloy D, Watts R, Benoit B, et al. (1983) Intraventricular haemorrhage from ruptured aneurysm. Retrospective analysis of 91 cases. *J Neurosurg* 58: 482-487.
11. Nieuwkamp DJ, Verweij BH, Rinkel GJ (2009) Massive intraventricular haemorrhage from aneurysmal rupture: patient proportions and eligibility for intraventricular fibrinolysis. *J Neurol* 257: 354-358.
12. Jabbarli R, Reinhard M, Roelz R, Shah M, Niesen WD, et al. (2016) The predictors and clinical impact of intraventricular hemorrhage in patients with aneurysmal subarachnoid hemorrhage. *Int J Stroke* 11: 68-76.
13. Pietila TA, Pallese H, Brock M (1995) Influence of aneurysm location on the development of chronic hydrocephalus following SAH. *Acta Neurochirurgica* 137: 70-73.
14. Kirmani AR, Sarmast AH, Bhat AR (2015) Role of external ventricular drainage in the management of intraventricular hemorrhage; its complications and management. *Surg Neurol Int* 6: 188.
15. AlShardan MM, Mubasher M, Orz Y, Yamany MA (2015) Factors that predict hydrocephalus following intraventricular hemorrhage. *Br J Neurosurg* 29: 225-228.
16. Jabbarli R, Bohrer AM, Pierscianek D, Muller D, Wrede KH, et al. (2016) The CHES score: a simple tool for early prediction of shunt dependency after aneurysmal subarachnoid hemorrhage. *Eur J Neurol* 23: 912-918.
17. Fung C, Balmer M, Murek M, Z'Graggen WJ, Abu-Isa J, et al. (2015) Impact of early-onset seizures on grading and outcome in patients with subarachnoid hemorrhage. *J Neurosurg* 122: 408-413.
18. Giraldo EA MJ, Rubin MN, Dupont SA, Zhang Y, Lanzino G, et al. (2012) Timing of clinical grade assessment and poor outcome in patients with aneurysmal subarachnoid hemorrhage. *J Neurosurg* 117: 15-19.