

Severe Hypertriglyceridemia-Induced Pancreatitis in Young Female Managed with Plasmapheresis: A Case Report

Ahmed A Khalifa¹, Sheref A Elseidy^{1*} and Ahmed M Khalifa²

¹Department of Rheumatology, Ain Shams University, Egypt

²Department of Forensic Medicine and Toxicology, Ain Shams University, Egypt

Abstract

Background: Hypertriglyceridemia (HTG) is postulated to be a rare cause of acute pancreatitis accounting for approximately 4% of causes. A serum triglyceride (TG) level above 1,000 mg/dl in the setting of pancreatitis and absent of other major causes is necessary to describe hypertriglyceridemia as a cause of acute pancreatitis. The mechanism involves the degradation of triglycerides by pancreatic lipase to release free fatty acids that induces free radical damage to the tissues. The major cornerstone in management is by reducing the triglyceride level to below 1000 mg/dL to achieve both the subsiding of the ongoing attack and the effective prevention of further episodes of pancreatitis. This can be achieved by a spectrum of treatment modalities ranging from includes dietary restriction of fat, administration of lipid-lowering agents to Experiences with plasmapheresis and effective lipid pheresis however, the latter modality hasn't been sufficiently discussed regarding therapeutic effects.

Case presentation: This case reports a young female, 22-year-old female patient non-alcoholic, non-diabetic, non-obese with negative history for gall stones or significant drug use diagnosed with hypertriglyceridemic acute pancreatitis associated with acute kidney injury, and received one session of hemodialysis then plasmapheresis was initiated and she received 8 sessions with full dose of hypolipidemics, and responded only to plasmapheresis.

Keywords: Acute pancreatitis; Hyperlipidemia; Hypertriglyceridemia; Plasmapheresis

Introduction

Hypertriglyceridemia (HTG) is the third most common cause of acute pancreatitis (AP) episodes [1-3]. Workup of AP caused by hyperlipidemia is the same as in all types of pancreatitis. In addition to exclusion of heavy alcohol consumption, gallstone disease and secondary causes as drugs and iatrogenic causes as post ERCP pancreatitis. A positive family history of signs and symptoms of hypertriglyceridemia is necessary to steer the diagnosis. Current guidelines of management must include full dose hypolipidemic mainly fibrates, minimization of dietary fat consumption, insulin and/or heparin, administration of plasmapheresis and/or lipid pheresis have been found to be the most useful in reduction of TG levels, improving of clinical signs and symptoms during the active phase of pancreatitis also effectively preventing further episodes of pancreatitis [4,5]. In this case report a female patient with positive family history for hyperlipidemia, no past significant history of diabetes or history of alcoholism or gall stones or significant drug history, was diagnosed with hyperlipidemia induced pancreatitis and received one session of hemodialysis then plasmapheresis was initiated and she received 8 sessions with full dose of hypolipidemics. She responded only to plasmapheresis which reversed the acute attack and exponentially reduced her TGs levels, but recurrence occurred although multiple sessions of plasmapheresis were done [4].

Case Report

A young 22-year-old female patient presented to our ER by epigastric pain, vomiting and fever. O/E: she was fully conscious, feverish, with epigastric tenderness. Pelvic-abdominal ultrasound showed bulky heterogeneous pancreas with surrounding rim of free peripancreatic fluid collection and was diagnosed with pancreatitis. While withdrawing her blood for laboratory tests, her blood looked milky white. Laboratory results showed: elevated kidney functions (creatinine 3.4 mg/dL) and serum amylase 209 IU. TGs level was (1335 mg/dL) and serum cholesterol was (750 mg/dL). She did not improve on full dose of hypolipidemics, then developed acute kidney injury and her serum blood urea nitrogen was 73 mg/dL and serum creatinine was 6.2 mg/dL and received one

session of hemodialysis followed by 8 sessions of plasmapheresis with full dose hypolipidemics. She responded only to plasmapheresis. Her TGs and cholesterols started to decrease and reached 545 mg/dL and 194 mg/dL respectively and she clinically started to improve with a serum amylase of 11 IU. Plasmapheresis was stopped. Unfortunately, she deteriorated again with more severe symptoms and revelation of TGs and cholesterol. computerized tomography abdomen revealed bulky heterogeneous pancreas with high density fluid collection and multiple air foci suggestive of necrotizing pancreatitis with moderate ascites and bilateral pleural effusion. the patient then had 2 attacks of generalized tonic clonic fits followed by deterioration of conscious level. she developed disseminated intravascular coagulopathy (DIC) and acute respiratory distress syndrome (ARDS) and unfortunately passed away.

Discussion

Hyperlipidemia as a cause of acute pancreatitis was first discussed in 1865 by Speck, the relationship then has been thoroughly studied in numerous papers and reports. The mechanism behind which hyperlipidemia leads to pancreatitis is thought to be that serum chylomicrons or triglycerides are degraded by the pancreatic lipase to generate free fatty acids that causes pancreatic capillary inflammation and thrombosis within the pancreatic capillary bed leading to pancreatitis. Free fatty acids also either exerts a toxic effect on pancreatic acinar cells or damages the capillaries directly via free radical damage. Elevated concentrations of free fatty acids in the pancreatic capillaries causes capillary inflammation and thrombosis leading to ischemia,

*Corresponding author: Sheref A. Elseidy, Department of Rheumatology, Ain Shams University, Egypt, Tel: +2001001571545; E-mail: drsherif.elseidy@gmail.com

Received September 26, 2017; Accepted November 23, 2017; Published November 28, 2017

Citation: Khalifa AA, Elseidy SA, Khalifa AM (2017) Severe Hypertriglyceridemia-Induced Pancreatitis in Young Female Managed with Plasmapheresis: A Case Report. J Clin Case Rep 7: 1051. doi: [10.4172/2165-7920.10001051](https://doi.org/10.4172/2165-7920.10001051)

Copyright: © 2017 Khalifa AA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

infarction and acidosis, also free fatty acids in this highly acidic environment activate pancreatic enzymes and thus initiating acute pancreatitis [6-9]. The two major signs of hyperlipidemic induced pancreatitis are high serum triglyceride level above 1,000 mg/dL and hyperchylomicronemia at the early stage of pancreatitis in a non-obese non-diabetic non-alcoholic patient with no other secondary cause for pancreatitis. Current management guidelines include: hypolipidemic agents, fat restriction, heparin infusion, insulin and plasmapheresis. However, with heparin the effect on triglyceride reductions is transient and the patient experience re-accumulation of TGS with long term infusion [10,11], on the other hand insulin use in non-diabetics has no clear evidence [11]. Patients with HTG-AP are more susceptible to systemic inflammatory response syndrome and end organ damage than in other types of pancreatitis [8] making therapeutic plasma exchange most beneficial to reduce serum triglyceride levels, as it was proposed that its efficacy is superior especially when there is evidence of shock or end-organ failure due to systemic inflammatory response syndrome [5]. In our case report the young patient presented with severe form of hyperlipidemic pancreatitis in the acute phase with triglyceride levels at first of (1335 mg/dL), management with a trial of hypolipidemics was unsatisfactory and only Plasma exchange lowered the lipid level and TGs with a value reaching (934 mg/dl) after the first session and also improved the clinical status of the patient. But even though eight sessions were sufficient to control active disease they did not prevent recurrence in this patient with a more severe picture and superadded ARDS and DIC.

Conclusion

Plasmapheresis is the best treatment modality to reverse the acute phase of hyperlipidemic induced pancreatitis. However, the

role of plasmapheresis administration to prevent recurrence of acute pancreatitis remains unclear and further research is indicated.

References

1. Tsuang W, Navaneethan U, Ruiz L, Palascak J, Gelrud A (2009) Hypertriglyceridemic pancreatitis: Presentation and management. *Am J Gastroenterol* 104: 984-991 .
2. Yadav D, Pitchumoni C (2003) Issues in hyperlipidemic pancreatitis. *J Clin Gastroenterol* 36: 753-758.
3. Fortson M, Freedman S, Webster P (1995) Clinical assessment of hyperlipidemic pancreatitis. *Am J Gastroenterol* 90: 2134-2139.
4. Lennertz A, Parhofer K, Samtleben W, Bosch T (1999) Therapeutic plasma exchange in patients with chylomicronemia syndrome. complicated by acute pancreatitis therapeutic. *Apheresis* 3: 227-233.
5. Click B, Ketchum A, Turner R (2015) The role of apheresis in hypertriglyceridemia-induced acute pancreatitis: A systematic review. *Pancreatol* 15: 313-320.
6. Havel R (1969) Pathogenesis, differentiation and management of hypertriglyceridemia. *Adv Intern Med* 15: 117-54.
7. Valdivielso P, Ramirez-Bueno A, Ewald N (2014) Current knowledge of hypertriglyceridemic pancreatitis. *Eur J Intern Med* 25: 689-694.
8. Nawaz H, Koutroumpakis E, Easler J, Slivka A, Whitcomb DC, et al. (2015) Elevated serum triglycerides are independently associated with persistent organ failure in acute pancreatitis. *Am J Gastroenterol* 110: 1497-1503.
9. Dominguez MJ, Malfertheiner P, Ditschneit H (1991) Hyperlipidemia in acute pancreatitis. Relationship with etiology, onset and severity of the disease. *Int J Pancreatol* 10: 261-267 .
10. Näsström B, Olivecrona G, Olivecrona T, Stegmayr BG (2001) Lipoprotein lipase during continuous heparin infusion: Tissue stores become partially depleted. *J Lab Clin Med* 138: 206-213.
11. Zhang K, Cox K, Sellers Z (2017) Plasmapheresis for hypertriglyceridemia-induced acute pancreatitis in a child: A case report and brief review of the literature. *Pancreas* 46: e58-e59 .