

Single I.V.-ketamine Augmentation of Newly Initiated Venlafaxine Hydrocortisone for Treatment Resistant Depression: Randomized, Double blind Controlled Study Protocol

Yun Du, Ruobing Feng, Wenle Zhang, Jiexin Fang, Ke Ma, Encong Wang and Yongdong Hu*

Department of Clinical Psychology, Beijing Chao-Yang Hospital, Capital Medical University, China

Abstract

Background: The critical issue of treating Treatment Resistant Depression (TRD) is to improve antidepressant effect rapid onset and the recovery rates. Significant antidepressant effects of ketamine onset rapidly, but decays rapidly and repeated dosing increases the risks. This study aimed to explore the efficacy and safety of single intravenous ketamine, combined with venlafaxine's hydrochloride for TRD.

Methods: 32 patients with treatment resistant depression will be randomly selected to undergo a four weeks double blind treatment with single dose ketamine (0.5 mg/kg over 40 min) and venlafaxine hydrochloride (4 weeks) or midazolam maleate (0.045 mg/kg over 40 min) and venlafaxine hydrochloride (4 weeks). Depressive symptoms will be measured by the Montgomery-Asberg Depression Rating Scale (MADRS) and the 16 item Quick Inventory of Depressive Symptomatology Self Report (QIDS-SR16). Adverse effects will be measured by the Brief Psychiatric Rating Scale-4 (BPRS-4), Young Manic Rating Scale (YMRS), the Clinician Administered Dissociative States Scale (CADSS), Frequency and Intensity of Side Effects Rating/Global Rating of Side Effects Burden (FISER/GRSEB) and Patient Rated Inventory of Side Effects (PRISE). Patients will be assessed at baseline, after intervention 1, 2, 4, 24, 48, 72 hours and 7, 14, 21 and 28 days.

Discussion: This study will provide important information on whether this new combined intervention can rapidly relieve the symptoms of depression and improve the remission rates of treatment resistant depression.

Keywords: Ketamine • Treatment resistant depression • Venlafaxine hydrochloride • Randomized controlled group

Background

Major Depressive Disorder (MDD) affects more than 300 million people worldwide and has an increased risk of suicide and mortality [1,2]. 12%-30% of patients with MDD are Treatment Resistant Depression (TRD) which have a failed response to antidepressant therapies [3-5]. The patients with TRD have a higher risk of suicide [6,7]. Although electroconvulsive therapy (ECT) is the most reliably effective therapy for TRD some patients forgo ECT due to adverse effects [8-11]. Moreover, oral antidepressants onset at least two weeks, and about two thirds of patients have an insufficient response [12-14]. Important issues in the treatment of TRD are antidepressant effect rapid onset and improve the recovery rates of TRD.

The rapid antidepressant actions of ketamine have been proven in many clinical studies [15-18]. Ketamine was approved to be effective by the U.S. Food and Drug Administration (FDA) for the use of an aesthetic in 1970. It has become a revolutionary discovery in the last few decades, and researchers have developed a keen interest in its use since the onset of rapid antidepressant effects of ketamine was discovered [19,20]. Ketamine improves depressive symptoms rapidly by blocking the N-methyl-D-aspartate

receptor's (NMDAR) dependent bursting activity of the lateral habenula (LHb) neurons to disinhibit reward centres [21]. Ketamine onset within hours after injection significantly relieves the symptoms of depression, and reduces the risk of acute suicide with a maximum efficacy at 24 hours after injection, with the effects of ketamine lasting seven days [22,23]. Moreover, when other traditional treatments have failed ketamine improves depressive symptoms [24].

On March 5, 2019, the U.S. FDA approved intranasal es-ketamine for TRD [25]. Previous studies have assessed the efficacy and safety between es-ketamine and ketamine in patients with TRD, and it shows that both medicines are effective, relatively safe and well tolerated [26,27]. Compared with nasal spray, intravenous ketamine is preferred for greater dosing accuracy and ease of administration [28].

The antidepressant effect of single dose ketamine is maintained for about seven days, to maintain the antidepressant effect have to use multiple doses ketamine [29-31]. The researchers have limited longer term studies 0.5mg/kg ketamine on safety, and the potential neurocognitive impairment and urinary difficulties by chronic and repeated use of ketamine are the main safety concerns [13,24,32-35]. To that end, the purpose of our study is to prolong the antidepressant effect of ketamine, and avoid to intravenous multiple doses ketamine.

Previous studies attempted to use ketamine with other antidepressants or treatments [16,36-39] to find a therapeutic dose which can balance safety and effectiveness. Intravenous ketamine has been used in combination with various antidepressants, such as escitalopram, sertraline duloxetine or venlafaxine extended release to enhance the antidepressant effect. In these studies, on the variety of oral antidepressants, venlafaxine, a serotonergic and noradrenergic reuptake inhibitor (SRNI) has had an increased efficacy on treatment than selective serotonin reuptake inhibitors (SSRIs) [40,41]. It has been well tolerated in patients with TRD [42,43].

Previous studies are shown that patients responded to an antidepressant

***Address for Correspondence:** Yongdong Hu, Department of Clinical Psychology, Beijing Chao-Yang Hospital, Capital Medical University, China, E mail: angusle19980328@gmail.com

Copyright: © 2022 Yun Du, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Received: 02-February-2022, Manuscript No. jnd-22-53882; **Editor assigned:** 04- February-2022, PreQC No: P-53882 (PQ); **Reviewed:** 18- February-2022, QC No: Q-53882; **Revised:** 23- February-2022, Manuscript No: R-53882 (R); **Published:** 02-March-2022, DOI:10.4172/2329-6895.10.2.266

with a placebo, using an active placebo in the control group is essential to improve the reliability of research [44,45]. Midazolam maleate is a psychoactive anaesthetic agent, the half-life and partial clinical response for ketamine and midazolam maleate are similar, and it is not proven to have an antidepressant effect [46,47]. In previous studies about the effectiveness of ketamine, midazolam was used as an active placebo control for nasal spray esketamine and infusion of ketamine [48-52]. Therefore, we use midazolam as active placebo instead of saline.

Objectives

In the study we use single I.V.-ketamine augmentation of newly initiated venlafaxine hydrochloride for TRD to improve the recovery rate of TRD, prolong the antidepressant effect of ketamine, and avoid the side effects of intravenous multiple doses ketamine.

Methods

Study design

This study is a randomized, double blind parallel controlled and clinical experiment. The control group will have an active medical control in the form of a single I.V. injection of midazolam maleate.

Participants

The samples will consist of 32 outpatients with TRD recruited from the Department of Clinical Psychology, Beijing Chaoyang Hospital, and Capital Medical University. Participants will be randomly divided into a study group and a control group, with 20 in each group.

Inclusion criteria

1. Patients who meet the diagnostic criteria for depression in the Diagnostic and Statistical Manual of Mental Disorders-IV (DSM-IV) without psychotic symptoms diagnosed by a psychiatrist;
2. Meet the criteria for TRD. The treatment was ineffective after a sufficient and complete course of treatment with at least two antidepressants (over six weeks);
3. The total score is greater than or equal to 17 on the 17 item Hamilton Rating Scale for Depression (HAM-D);
4. 25-64 years old, both genders;
5. No antidepressants, antipsychotics, mood stabilizers, electrical shock treatments in the last two weeks;
6. Have the ability to understand the content of the scale and cooperate with the assessment;
7. Sign research informed consent and be able to follow up.

Exclusion criteria

1. Patients with a previous history of substance use disorders or acute poisoning; patients with psychotic disorder, bipolar disorder, obsessive compulsive disorder;
2. The primary diagnosis is not depression;
3. History of inefficacy or intolerance to Venlafaxine;
4. Women who are pregnant or breastfeeding;
5. Patients who are at high risk of suicide: suicide attempts, recent suicide attempts; no family caregivers;
6. Patients who are contraindicated in using ketamine, or midazolam maleate, or venlafaxine hydrochloride;
7. Patients who have used ECT or NMDA receptor antagonists within the past six months;

8. There are currently patients with respiratory disease, hypertension, or other physical severe diseases, such as patients with severe cardiovascular disease and hyperthyroidism, patients with intracranial hypertension or cerebral haemorrhage and glaucoma.

Participant withdrawal and termination criteria

1. The patient has a concomitant physical illness or serious adverse effect and others, who are unable or unwilling to continue to complete the study protocol;
2. The patient of lost follow up;
3. The patient switch to mania, hypomania during the study;
4. The patient suicide or attempted suicide occurred during the study;
5. The patient no longer wishes to be part of the trial.

Safety considerations

The anesthesiologist and nurse will give the injection treatment in the operating room. During the injection process, the blood pressure, oxygen hemoglobin saturation and other significant physiological indicators will be monitored in real time. The psychiatrist will evaluate and record the side effects throughout the injection treatment. Any discomfort of the patient during the whole treatment process will be reported to the psychiatrist and anesthesiologist, who will act to alleviate this and record the side effects. Four hours after the injection treatment, the patient can be discharged from the hospital, only if they do not have discomfort and have been safely evaluated without side effects.

Block randomization will be used; the length of the block is four, which means there are 4 study subjects in each block. SAS software will be used to generate a random allocation list. Based on the random allocation list, the research subjects in the block group will be randomly allocated to the experimental group and the control group at a ratio of 1: 1. The random allocation is sealed with an opaque random concealed letter printed by a pinhole and given to a designated person (not involved in clinical observation assigned by the project center for keeping).

Blind

The double blind method will not be used in this study; neither the experimenter nor do the participants know which treatment is received. We chose midazolam maleate instead of the placebo used in previous experiments -0.5% saline, because ketamine can cause some unique drug interactions within patients. Participants are randomly assigned to a single intravenous infusion of I.V.-ketamine 0.5 mg/kg or midazolam maleate 0.045 mg/kg. Questionnaires will be scored by professional raters who do not know which kind of drugs participants have been administered.

Primary outcome measures

The time required to reach the clinical effective standard and clinical cure standard will be measured. The effective standard is the remission rates of the total Montgomery-Asberg Depression Rating Scale (MADRS) score $\geq 50\%$ compared with the baseline, and the clinical cure standard is defined as the total MADRS score ≤ 10 .

Secondary outcome

16 item Quick Inventory of Depressive Symptomatology Self Report (QIDS-SR16). Patients will be assessed at baseline, after I.V.-ketamine at 1, 2, 4, 24, 48, 72 hours, 7, 14, 21 and 28 days.

Side effects and safety

Record vital signs (blood pressure, heart rate, and oxygen saturation) every 5 minutes in the duration I.V.-ketamine.

Adverse psychopathological effects will be measured by/with the Brief Psychiatric Rating Scale-4 (BPRS-4), Young Manic Rating Scale (YMRS), the Clinician Administered Dissociative States Scale (CADSS), Frequency

and Intensity of Side Effects Rating/Global Rating of Side Effects Burden (FISER/GRSEB), and Patient Rated Inventory of Side Effects (PRISE). Patients will be assessed at baseline, after I.V.-ketamine at 1, 2, 4, 24,

48, 72 hours, 7, 14, 21 and 28 days. The schedule of assessments and procedures is summarized in (Table 1).

Table 1. Study procedures and schedule of assessments experiments

| Study Period | | | | | | | | | | | | |
|--|---------------------|--------------------|---------|----------|----------|---------|---------|---------|---------|---------|----------|----------|
| | Screening (Visit 0) | BaseLine (Visit 1) | Visit 2 | Visit 3 | Visit 4 | Visit 5 | Visit 6 | Visit 7 | Visit 8 | Visit 9 | Visit 10 | Visit 11 |
| After Infusion | | | | | | | | | | | | |
| Time Point | | Visit 4 | 1 Hours | 2 Hours | 4 Hours | Day 1 | Day 2 | Day 3 | Day 7 | Day 14 | Day 21 | Day 28 |
| Informed Consent | | Visit 5 | | | | | | | | | | |
| Demographic and Clinical characteristics | | Visit 6 | | | | | | | | | | |
| Physicalexamination | | Visit 7 | x | x | x | x | x | x | x | x | x | x |
| Major depression disorder (DSMV) | | Visit 8 | Visit 9 | Visit 10 | Visit 11 | | | | | | | |
| Treatment- resistant depression | x | | | | | | | | | | | |
| Clinical examination | | x | | | | | | | | | | x |
| HAMD-17 | x | | | | | | | | | | | |
| Vital signs | x | x | x | x | x | x | x | x | x | x | x | x |
| MADRS | | x | x | x | x | x | x | x | x | x | x | x |
| BPRS-4 | | x | x | x | x | x | x | x | x | x | x | x |
| CADSS | | x | x | x | x | x | x | x | x | x | x | x |
| YMRS | | x | x | x | x | x | x | x | x | x | x | x |
| QIDS- SR 16 | | x | x | x | x | x | x | x | x | x | x | x |
| PRISE | | x | x | x | x | x | x | x | x | x | x | x |
| FISER/GRSEB | | x | x | x | x | x | x | x | x | x | x | x |

Intervention

Participants will meet the criteria and sign informed consent. Participants will undergo a psychotropic medications washout period of two weeks (fluoxetine hydrochloride for four weeks).

The injection treatment is performed on the first day of the intervention. Preparation before treatment: The patient fasts for 12 hours and drinks water for 4 hours. Control Groups: establish an intravenous infusion channel in the operating room, and inject physiological saline at a rate of 20-30 drops/min. Connect a physiological monitor to monitor BP, SpO2, ECG. Ketamine Group: 0.5 mg/kg ketamine hydrochloride injection, diluted with a physiological saline injection to 50 ml intravenous for 40 minutes. Control group: midazolam maleate injection, a dose of 0.045 mg/kg, diluted with a saline injection to 50 ml intravenous injection for 40 minutes. Both groups: Blood pressure, heart rate, and oxygen saturation will be recorded every 5 minutes during drug injection. Observe the patient's consciousness and mental state for 4 hours during and after administration.

Both groups are treated with venlafaxine hydrochloride 150 mg/day (the initial dose was 75 mg/d and increased to 150 mg/d one week later) from the first day of the intervention for four weeks.

In the study, if the subject has severe insomnia (cannot fall asleep half an hour after going to bed), zolpidem tartrate can be temporarily administered 5-10 mg, ≤ 3 times a week. Do not use systemic psychotherapy, electroconvulsive therapy, antidepressants, and antipsychotics other than research medications.

Sample size

Sample size calculation: using the superiority hypothesis test, setting unilateral $\alpha=0.05$, $\beta=0.2$. With reference to the previous study setting, the treatment time of the control group that reached the clinical cure standard was set to 28.0 ± 0.0 days, and the mean treatment standard deviation of the experimental group was 15.8 ± 13.0 days [53]. Based on previous literature and the time required for clinical cure, the optimal effect cut off value $\Delta=-5$ days was calculated using PASS14.0 software. 13 samples were required for each group [4-6]. Considering the 15% missed follow up rate and random block capacity, the number of expanded samples is 16 in each group, and 32 samples are required for the two combinations.

Statistical analytical plan

Primary outcome analysis: Kaplan-Meier survival analysis will be used as the time required reaching the effective standard and clinical cure standard. Cox proportional-hazards regression models are to be used to compare the estimated time to respond and to remission between the two groups, controlling the baseline effects.

Secondary outcome analysis: Baseline demographics, clinical feature analysis, effectiveness, and clinical cure rate between the two groups will be measured using two independent samples'-test or Mann-Whitney U test or χ^2 test or Fisher's exact test. The significance level will be set at 0.05 (in a two-tailed test).

Data management and monitoring strategies

Researchers fill in the data on the case report form in time after each

visit, and this is to be consistent with the original record. The data will be collected on Epidata to build the database. Special management and double verification are adopted to ensure the accuracy of data entry. According to the Intention to Treat (ITT), an intent analysis data set (subjects who took at least one drug after enrollment) will be established to analyze the research data statistically.

For subjects who did not observe the full effect evaluation, a mixed model will be used for data processing.

Discussion

In recent years, although oral antidepressants have improved in terms of side effects and effectiveness, patients still face the dilemma of the long onset of antidepressants and limited treatment effect [54,55]. Since the discovery of rapid antidepressant effect, ketamine has been the focus of treatment research. However, there are always some problems that need to be solved within clinical applications. The antidepressant effects of ketamine decay rapidly after a single dose, while repeated dosing increases the risks includes physical injury and substance use disorders [56,57].

In order to avoid the risks of repeated dosing, the current study only uses a single injection of I.V. ketamine at the beginning of treatment. Initiated at the beginning of oral antidepressants solves the dilemma in efficacy decreasing after a single injection of ketamine. This combination therapy reduces patient's suicide risk during these weeks before oral antidepressants onset. In our study, it is first important to select venlafaxine, a SNRI, as an oral antidepressant in combination with I.V. ketamine. We will set follow up after injection I.V. ketamine at 1, 2, 4, 24, 48, 72 hours and 7, 14, 21 and 28 days. Follow up at multiple time points can indicate the law of ketamine decay, to accurate evolution the time of ketamine onset and recover. From compared ketamine and control in this study, we explore more precisely enhancement of venlafaxine to ketamine for accelerating the reduction of symptom.

The study is the first to verify the effectiveness and safety of ketamine with TRD in Chinese. In the past studies, there are few RCT studies use ketamine in Chinese with TRD, who are needs more effective and safer treatment options due to the higher risk of suicide and mortality within this group.

The clinically meaningful treatment effect of therapy will be evidenced by the reduction of MADRS and QIDS-SR16 total score: the most commonly and valid standard clinician rating scale for depression [58,59]. In previous studies, either the MADRS or QIDS-SR16 has been used alone. Because there are individual differences in emphases and evaluation methods between these scales, we will use both scales in our trial to more accurately assess depressive symptoms. Both scales will be used to examine depressive symptoms comprehensively due to their different characteristics in evaluations. Previous research has shown that minor side effects briefly experienced by patients include symptoms like dissociation and elevated blood pressure after using ketamine, but these side effects usually happened during the initial dosing and the first few hours [60,61]. BPRS-4, YMRS, CADSS, FISER/GRSEB, PRISE total score and the physical sign during injection will be monitored for side effects. This data can then provide information on the safety of ketamine combined with venlafaxine.

Conclusion

This clinical trial is intended to study the efficacy and safety of I.V. ketamine initiated venlafaxine for TRD and our findings will provide valuable information for clinical applications, which may assist psychiatrist to use treatments more effectively in the future in China.

Availability of Data and Materials

All the data and materials are availability.

Competing Interests

The authors declare they have no competing interests.

Funding

The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: Capital Health Development Scientific Research Project (No.: First Issue 2018-2-2033). 2018 Beijing Municipal Key Clinical Specialty Construction Project (Psychiatric). Beijing Municipal Key Laboratory Open Project (No. 2018JSJB01). Beijing Ronghe Medical Development Foundation.

Authors' Contributions

Y Du as the first author designed the whole study and wrote this manuscript. R-B Feng and W-L Zhang helped with the statistical analysis and language modification. J-X Fang, K Ma and E-C Wang went through and revised the manuscript. Y-D Hu supervised the design of the study through the writing of the manuscript.

Acknowledgements

Not applicable

References

1. Depression: Key facts. World Health Organization (WHO) (2018).
2. Walker, ER, Mcgee RE and Druss BG. "Mortality in mental disorders and global disease burden implications: A systematic review and meta-analysis". *JAMA Psychiatry* 72(2015):334-41.
3. Mrazek, DA, Hornberger JC, Altar CA and Degtiar I. "A review of the clinical, economic, and societal burden of treatment-resistant depression:1996-2013". *Psychiatr Serv* 65(2014):977-987.
4. Murrough, JW, Abdallah CG and Mathew SJ. "Targeting glutamate signalling in depression: Progress and prospects". *Nat Rev Drug Discov* 16(2017):472-486.
5. Rush, AJ, Trivedi MH and Wisniewski SR. "Acute and longer-term outcomes in depressed outpatients requiring one or several treatment steps". *Am J Psychiatry* 163(2006):1905-1917.
6. Fava, M. "Diagnosis and definition of treatment-resistant depression". *Biol Psychiatry*, 2003, 53(8):649-659.
7. Amital, D, Fostick L, Silberman A and Beckman M. "Serious life events among resistant and non-resistant MDD patients". *J Affect Disord* 110(2008):260-264.
8. Grunebaum, MF, Galfalvy HC and Choo TH. "Ketamine for rapid reduction of suicidal thoughts in major depression: A midazolam-controlled randomized clinical trial". *The Am J Psychiatry* 175(2018):327.
9. Cusin, C and Dougherty DD. "Somatic therapies for treatment-resistant depression: ECT, TMS, VNS, DBS". *Biol Mood Anxiety Disord* 2(2012):14.
10. Pagnin, D, De Queiroz V, Pini S and Cassano GB. "Efficacy of ECT in depression: A meta-analytic review". *J Ect* 20(2004):13-20.
11. Regenold William, T, Noorani Robert J and Piez Deborah, et al. "Nonconvulsive electrotherapy for treatment resistant unipolar and bipolar major depressive disorder: A proof-of-concept trial". *Brain Stimul* 8(2015): 855-61.
12. Papakostas, GI, Perlis RH and Scalia MJ, et al. "A meta-analysis of early sustained response rates between antidepressants and

- placebo for the treatment of major depressive disorder". *J Clin Psychopharmacol* 26(2006):56-60.
13. Posternak, MA and Zimmerman M. "Is there a delay in the antidepressant effect? A meta-analysis". *J Clin Psychiatry* 66(2005):148.
 14. Volonteri, G, Cerveri A, Colasanti I and De Gaspari M. "Clinical outcome and tolerability of duloxetine in the treatment of major depressive disorder: A 12-week study with plasma levels". *J Psychopharmacol* 24(2008):23.
 15. Zarate, CA, Singh JB and Carlson PJ. "A randomized trial of an n-methyl-d-aspartate antagonist in treatment-resistant major depression". *Arch Gen Psychiatry* 63(2006):856.
 16. Hu, YD, Xiang YT and Fang J X. "Single i.v. ketamine augmentation of newly initiated escitalopram for major depression: Results from a randomized, placebo-controlled 4-week study". *Psychol Med* 46(2016):623-635.
 17. Murrough, JW, Iosifescu DV and Chang L. "Antidepressant efficacy of ketamine in treatment-resistant major depression: A two-site, randomised controlled trial". *Am J Psychiatry* 23(2013): S411-S412.
 18. Andrade, C. "Ketamine for depression, 1: Clinical summary of issues related to efficacy, adverse effects, and mechanism of action". *J Clin Psychiatry* 78(2017): e415.
 19. Berman, RM, Cappiello A, Oren, DA and Heninger, GR, et al. "Antidepressant effects of ketamine in depressed patients". *Biol Psychiatry* 47(2000): 351-354.
 20. Newport D, Jeffrey, Carpenter Linda L, McDonald William M and Potash James B, et al. "Ketamine and other nmda antagonists: Early clinical trials and possible mechanisms in depression". *Am J Psychiatry* 172(2015):950-966.
 21. Yang, Y, Cui Y and Sang K. "Ketamine blocks bursting in the lateral habenula to rapidly relieve depression". *Nature* 554(2018):317-322.
 22. Coyle, CM and Laws KR. "The use of ketamine as an antidepressant: A systematic review and meta-analysis". *Hum Psychopharmacol* 30(2015):152-163.
 23. Fond, G, Loundou A, Rabu C and Macgregor A, et al. "Ketamine administration in depressive disorders: A systematic review and meta-analysis". *Psychopharmacology (Berl)* 231(2014):3663-76.
 24. Sanacora, G, Frye MA and McDonald W. "A consensus statement on the use of ketamine in the treatment of mood disorders". *JAMA Psychiatry* 74(2017):399.
 25. US Food and Drug Administration. FDA approves new nasal spray medication for treatment-resistant depression; available only at a certified doctor's office or clinic.
 26. Correia-Melo, Fernanda, Leal Gustavo C, Vieira Flávia and Jesus-Nunes Ana Paula, et al. "Efficacy and safety of adjunctive therapy using esketamine or racemic ketamine for adult treatment-resistant depression: A randomized, double-blind, non-inferiority study". *J Affect Disord* 264(2020):527-534.
 27. Molero, P, Ramos-Quiroga JA and Martin-Santos R. "Antidepressant efficacy and tolerability of ketamine and esketamine: A critical review". *CNS Drugs* 32(2018):411-420.
 28. Ruberto Valerie, L, Jha Manish K and Murrough James W. "Pharmacological treatments for patients with treatment-resistant depression". *Pharmaceuticals (Basel)* 13(2020):116.
 29. Aan Het Rot, M, Zarate Jr CA and Charney DS. "Ketamine for depression: Where do we go from here". *Biol Psychiatry* 72(2012):537-547.
 30. Coyle, CM and Laws KR. "The use of ketamine as an antidepressant: A systematic review and meta-analysis". *Hum Psychopharmacol* 30(2015):152-163.
 31. Lee, EE, Della Selva MP and Liu A. "Ketamine as a novel treatment for major depressive disorder and bipolar depression: A systematic review and quantitative meta-analysis". *Gen Hosp Psychiatry* 37(2015):178-184.
 32. Lee, EE, Della Selva MP and Liu A. "Ketamine as a novel treatment for major depressive disorder and bipolar depression: A systematic review and quantitative meta-analysis". *Gen Hosp Psychiatry* 7(2015):178-184.
 33. Wilkinson, ST and Sanacora G. "Considerations on the off-label use of ketamine as a treatment for mood disorders". *JAMA* 318(2017):793-794.
 34. Morgan, CJA and Curran HV. "Ketamine use: A review". *Addiction* 107(2011):27-38.
 35. Xu Y, Hackett M and Carter G. "Effects of low-dose and very low-dose ketamine among patients with major depression: A systematic review and meta-analysis". *Int J Neuropsychopharmacol* 19(2016):124.
 36. Daly EJ, Singh JB and Fedgchin M. "Efficacy and safety of intranasal esketamine adjunctive to oral antidepressant therapy in treatment-resistant depression: A randomized clinical trial". *Jama Psychiatry* 75(2017):139-148.
 37. McGirr, A, Berlind MT and Bond DJ. "Adjunctive ketamine in electroconvulsive therapy: Updated systematic review and meta-analysis". *Br J Psychiatry* 116(2017):195826.
 38. Ren, Li, Deng Jie, Min Su and Peng Lihua, et al. "Ketamine in electroconvulsive therapy for depressive disorder: A systematic review and meta-analysis". *J Psychiatr Res* 104(2018):144-156.
 39. Short, B, Fong J and Galvez V. "Side-effects associated with ketamine use in depression: A systematic review." *Lancet Psychiatry* 5(2017):65-78.
 40. Undurraga, J and Baldessarini RJ. "Randomized, placebo-controlled trials of antidepressants for acute major depression: Thirty-year meta-analytic review". *Neuropsychopharmacology* 37(2012):851-864.
 41. Rush AJ, Trivedi MH and Wisniewski SR. "Bupropion-sr, sertraline, or venlafaxine-xr after failure of ssris for depression". *N Engl J Med* 354(2006):1231-1242.
 42. Rot, MAH, Collins KA and Murrough JW. "Safety and efficacy of repeated-dose intravenous ketamine for treatment-resistant depression". *Biol Psychiatry* 67(2010):139-145.
 43. Murrough, JW, Perez AM and Pillemer S. "Rapid and longer-term antidepressant effects of repeated ketamine infusions in treatment-resistant major depression". *Biol Psychiatry* 74(2013):250-256.
 44. Cipriani, A, Furukawa TA and Salanti G. "Comparative efficacy and acceptability of 21 antidepressant drugs for the acute treatment of adults with major depressive disorder: A systematic review and network meta-analysis". *Lancet* 391(2018):1357-1366.
 45. Walsh, BT, Seidman SN and Sysko R. "Placebo response in studies of major depression". *JAMA* 287(2002):1840-7.
 46. Papakostas, GI, Søren D Østergaard and Iovieno N. "The nature of placebo response in clinical studies of major depressive disorder". *J Clin Psychiatry* 76(2015):456-466.
 47. Grunebaum, MF, Ellis SP and Keilp JG. "Ketamine versus midazolam in bipolar depression with suicidal thoughts: A pilot midazolam-controlled randomized clinical trial". *Bipolar Disord* 19(2017):176-183.
 48. Fan, Wei, Yang HaiKou, Sun Yong and Zhang Jun, et al. "Ketamine rapidly relieves acute suicidal ideation in cancer patients: A randomized controlled clinical trial". *Oncotarget* 8(2015):2356-2360.
 49. Grunebaum, MF, Ellis SP and Keilp JG. "Ketamine versus midazolam in bipolar depression with suicidal thoughts: A pilot midazolam-controlled randomized clinical trial". *Bipolar Disord* 19(2017):176-183.
 50. Grunebaum, MF, Galfalvy HC and Choo TH. "Ketamine for rapid reduction of suicidal thoughts in major depression: A midazolam-controlled randomized clinical trial. *Am J Psychiatry* 175(2018):327-

- 335.
51. Gálvez, Verónica, Li A and Huggins C. Repeated intranasal ketamine for treatment-resistant depression – the way to go? Results from a pilot randomised controlled trial. *J Psychopharmacol* 32(2018):397-407.
52. Phillips Jennifer, L, Norris Sandhaya, Talbot Jeanne and Birmingham Meagan, et al. "Single, repeated, and maintenance ketamine infusions for treatment-resistant depression: A randomized controlled trial". *Am J Psychiatry* 176(2019):401-409.
53. Doherty, T, Wajs E and Melkote R. "Cardiac safety of esketamine nasal spray in treatment-resistant depression: Results from the clinical development program". *Cns Drugs* 2020(10):1-12.
54. Garay RP, Zarate CA and Charpeaud T. "Investigational drugs in recent clinical trials for treatment-resistant depression". *Expert Rev Neurother* 17(2017):1-17.
55. Thase, ME, Friedman ES and Howland RH. "Venlafaxine and treatment-resistant depression". *Depress Anxiety* 12(2010):55-62.
56. Sir, A, D'Souza RF and Uguz S. "Randomized trial of sertraline versus venlafaxine in major depression". *J Clin Psychiatry* 66(2005):1312-1320.
57. Murrough, JW, Perez AM and Pillemer S. "Rapid and longer-term antidepressant effects of repeated ketamine infusions in treatment-resistant major depression". *Biol Psychiatry* 74(2013):250-256.
58. Grunebaum, MF, Galfalvy HC and Choo TH. "Ketamine for rapid reduction of suicidal thoughts in major depression: A midazolam-controlled randomized clinical trial". *Am J Psychiatry* 175(2018):327-335.
59. Bernstein, IH, Rush AJ and Stegman D. "A comparison of the qids-c16, qids-sr16, and the madrs in an adult outpatient clinical sample". *CNS Spectrums* 15(2010):458-468.
60. Jauhar, Sameer and Morrison Paul. "Esketamine for treatment resistant depression". *BMJ* 366(2019):155-172.
61. Katalinic, N, Lai R and Somogyi A. "Ketamine as a new treatment for depression: A review of its efficacy and adverse effects". *Aust N Z J Psychiatry* 47(2013):710-727.

How to cite this article: Yun, Du, Ruobing Feng, Wenle Zhang and Jiexin Fang et al. "Single I.V.-ketamine Augmentation of Newly Initiated Venlafaxine Hydrocortisone for Treatment Resistant Depression: Randomized, Double blind Controlled Study Protocol" *J Neurol Disord* 10(2022): 266