

Spectrophotometric Determination and Commercial Formulation of Tebuconazole Fungicide after Derivatization

Kafeel Ahmad Khan* and Farzana Haider

Bacha Khan University, Charsadda, KPK, Pakistan

Abstract

A spectrophotometric method has been developed for the determination of tebuconazole fungicide formulation. The method is based on the complexation reaction of ferric chloride with tebuconazole fungicide in acidic media and the purple colored complex was studied at 530 nm. The reaction conditions and other analytical parameters were optimized. A linear calibration curve between absorbance and concentration over the range from 0.2 to 20 $\mu\text{g mL}^{-1}$ was obtained with molar absorptivity of $2.1 \times 10^4 \text{ L mol}^{-1} \text{ cm}^{-1}$. The limit of detection (LOD) and limit of quantification (LOQ) were found to be 0.05 and 0.2 $\mu\text{g mL}^{-1}$ respectively. The proposed method has been successfully applied for the analysis of commercial formulation of tebuconazole fungicide. The recoveries of the method were found to be in the range of $92.16 \pm 0.06\%$ to $96.66 \pm 0.18\%$.

Keywords: Ferric chloride; Sodium hydroxide; Tebuconazole fungicide; UV/Visible spectrophotometer

Introduction

The production of good quality of crops is essential for the existence of human being. Many micro-organisms have spoiled crops which lead to worldwide food shortage. Among these micro-organisms fungi play a critical role in decaying crops [1-3]. Fungicides work in a variety of ways, but most of them damage fungal cell membranes or interfere with energy production within fungal cell [4,5]. Fungicides have changed the nature of agriculture. Researchers all around the world are involved in the synthesis of fungicides of new formulation in order to meet the modern world needs [4-6]. Tebuconazole is a fungicide commonly used in agriculture. Tebuconazole [1-(4-Chlorophenyl)-4,4-dimethyl-3-(1,2,4-triazol-1-ylmethyl)pentan-3-ol] is a triazole fungicide. The triazole (tebuconazole) fungicide have increased the persistence of disease control, an effect that leads to prolonged green leaf retention and extended grain fill compared to traditional fungicides [7-9]. Tebuconazole has novel mode of action, and are very safe from environmental point of view. Due to its fungicidal action the inhibition of mitochondrial respiration in fungi occurs and stopping their energy supply [10,11]. The plants for which tebuconazole are more effective are cereals, vines, fruits and vegetables, rice, turf and ornamentals [2,3]. Various studies on tebuconazole have been investigated by using the HPLC/MS, gas chromatography in combination with electron capture detection, gas chromatography [12,13], micellar electro kinetic chromatography [14,15], spectrophotometric methods reported for the metribuzin determination is based on its complexation with copper [16]. The present method is based on spectrophotometric determination of tebuconazole fungicide by reacting with ferric chloride in alkaline media through complexation reaction.

Apparatus

UV/Vis Spectrophotometer (Model SP-1800 plus, Optima Tokyo Japan), with matched 1 cm quartz cells.

Reagents

All chemicals used were of analytical reagent grade (NaOH and HCl (37%), Ferric chloride (99.98%) [Merck, Germany], Methanol

(99.98%) [Merck Darmstadt, Germany], Commercial formulated tebuconazole 70% w/w of active ingredient (Roots International Agro, Division) was purchased from local Market, Ferric Chloride (0.04 ml/L) solution was freshly prepared by dissolving 0.6 g in 100 ml of methanol. Tebuconazole stock solution (1000 $\mu\text{g/ml}$) was prepared by dissolving 0.1 mg of authentic standard reagent in 50 ml methanol and diluted upto 100 ml. Working standard of 100 and 10 $\mu\text{g/ml}$ solutions were prepared from stock solution by diluting with methanol. Sodium hydroxide solution of 0.87 mol/L was used.

Procedure

Working of standard solution containing tebuconazole in the range of 0.2-20 $\mu\text{g/ml}$ were transferred quantitatively into reaction flask followed by the addition of 0.4 ml Ferric chloride 5% and 0.4 ml of NaOH (5M) solutions. The absorbance of the coloured product was measured at 530 nm against reagent blank.

Determination of tebuconazole in commercial formulation

Known amount of commercial formulation containing tebuconazole was taken and diluted with methanol upto 50 mL and was filtered to remove any insoluble impurities. The solutions were then analyzed by the proposed method.

Results and Discussion


In the present study spectrophotometric determination of tebuconazole fungicide was carried out. For this purpose tebuconazole is first converted into a purple colored complex by reacting it with ferric chloride (III) in alkaline media. The proposed reaction mechanism is given as,

*Corresponding author: Kafeel Ahmad Khan, Bacha Khan University, Charsadda, 24420, KPK, Pakistan, E-mail: kafeelaps@gmail.com

Received June 10, 2016; Accepted June 25, 2016; Published June 30, 2016

Citation: Khan KA, Haider F (2016) Spectrophotometric Determination and Commercial Formulation of Tebuconazole Fungicide after Derivatization. Chem Sci J 7: 133. doi:10.4172/2150-3494.1000133

Copyright: © 2016 Khan KA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Scanning λ_{\max} for tebuconazole

The purple colored complex was scanned for various wavelengths (400-680 nm). As can be seen from Figure 1 the colored complex shows maximum absorbance at 530 nm. The obtained maximum wavelength (λ_{\max}) of the complex was used for further analysis of the tebuconazole fungicide.

Optimization of ferric (III) chloride

For the concentration optimization of ferric chloride six solutions of drug were prepared each contain different concentration of ferric chloride and their absorbance were measured at 530 nm, the absorbance gradually increases and reaches the maximum value at 5 mol/L beyond it the absorbance decreases. As the Figure 2, shows that the solution of drug having 5 mol/L concentration of ferric chloride have high degree of capability to convert maximum amount of drug into complex which gives maximum absorbance. It is worth mentioning that; identical plot was also obtained by Shah et al. [16] in their study, concerning the optimization of ferric (III) chloride; although they applied Fenoxaprop-ethyl under similar experimental conditions. However, similar work is also reported by Jan et al. [17] for absorption vs. concentration of pesticide carbofuran.

Volume optimization of ferric chloride

For the optimization of volume of ferric chloride seven solutions of drug was prepared in each sample various quantity (0.1-0.7 ml) of ferric chloride was added. The absorbance of these various complex solutions were measured at 530 nm, the solution having 0.5 ml of ferric chloride showed maximum absorbance as shown in Figure 3.

Optimization of sodium hydroxide

For the optimization of sodium hydroxide various concentrations ranging from 1-6 mol/L were prepared. Each of this sodium hydroxide was added to six different samples of tebuconazole complex. Initially

as the concentration of sodium hydroxide raises the absorbance of the complex increases up to 2 mol/L as shown in Figure 4, followed by decreases in absorbance significantly. This is optimum concentration of sodium hydroxide at which complete complexation of tebuconazole occurs and maximum absorbance was observed

Volume optimization of sodium hydroxide

For investigation of the optimum volume of sodium hydroxide at which maximum complexation takes place, various volume ranging 0.1- 0.7 ml were added to tebuconazole. In Figure 5 shows the optimum volume of sodium hydroxide (0.4 ml) at which tebuconazole exhibits maximum absorbance. The complex at this alkaline media shows maximum absorbance as this optimum volume provides most favorable condition for complexation [16].

Stability of colored complex

To check the stability of the final reaction product, the effect of time was studied upto 3 days (72 h) with the interval of 2 h. The colored reaction product was found to be stable for about 2 days (48 h).

Analytical characteristics

The absorbance concentration curve was found linear over the concentration of 0.2-20 $\mu\text{g mL}^{-1}$. The molar absorptivity of the resulting yellow coloured product was found to be $2.1 \times 10^4 \text{ L mol}^{-1} \text{ cm}^{-1}$ with limit of detection (3s/b) and quantification (10s/b) of 0.05 and 0.2 $\mu\text{g mL}^{-1}$ respectively, where s is the slope and b is the intercept. For more confirmation of the analytical applications of the proposed method, the standard addition method was applied to a commercial product. The recovery studied were carried out after adding known quantities of authentic standard to the pre-analyzed formulation. The percentage recoveries were in the range of 93.68-96.66%. The value obtained for commercial formulation was comparable with the labeled value.

Verification of Beer Lambert law

For the verification of Beer's Lambert law a calibration curve between concentration and absorbance was plotted in Figure 6. The figure shows a linear relation between concentration and absorbance till 20 $\mu\text{g mL}^{-1}$ concentration and beyond that concentration deviation was observed. In other words the Beer's Lambert's law was obeyed up to concentration 20 $\mu\text{g mL}^{-1}$ (Table 1).









Figure 1: Determination of λ_{\max} for tebuconazole complex.



Characteristics	Analysis
λ_{\max} (nm)	530
Colour	Purple
Calibration Curve Range ($\mu\text{g}/\text{mL}^{-1}$)	0.2-20
Slope	0.0287
Limit of detection ($\mu\text{g}/\text{mL}^{-1}$)	0.05
Limit of quantification ($\mu\text{g}/\text{mL}^{-1}$)	0.2
Molar absorptivity ($\text{L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$)	2.1×10^4

Table 1: The optimum conditions for the highest absorption due to maximum colored product formation were investigated.

Conclusion

The proposed method described for spectrophotometric determination of tebuconazole is simple, rapid and cost-effective as compared to the reported chromatographic methods for determination

of tebuconazole. The color of the final product is stable at room temperature for two days and the limit of quantification shows that the developed method can be used to determine tebuconazole at lower concentration in different samples. The optimum conditions for the highest absorption due to maximum colored product formation were investigated.

References

1. Farajzadeh MA, Mogaddam MRA, Ghorbanpour H (2014) Development of a new microextraction method based on elevated temperature dispersive liquid-liquid microextraction for determination of triazole pesticides residues. *J Chrom A* 1347: 8-16.
2. Sobrinho F, Lucas F, Fill TP, Filho ER, Mascaro LH, et al. (2015) Insights into electrodegradation mechanism of tebuconazole pesticide on Bi-doped PbO₂ electrodes. *Electrochim Acta* 154: 278-286.
3. Sancho E, Villarroel MJ, Fernandez C, Andreu E, Ferrando, et al. (2010) Short-term exposure to sublethal tebuconazole induces physiological impairment in male zebrafish (*Danio rerio*). *Enviro Safety* 73: 370-376.
4. Tentu NR, Tentu P, Patrudu TB (2012) Simultaneous determination of triazole fungicide residues in Papaya followed by a new validated HPLC-UV method. *Int J Res Pure Appl Chem* 1: 1-5.
5. Zhang Z, Jiang W, Jian Q, Song W, Zheng Z, et al. (2015) Residues and dissipation kinetics of triazole fungicides difenoconazole and propiconazole in wheat and soil in Chinese fields. *Food Chem* 168: 396-403.
6. Juan E, Otero R, Paradelo M, López-Periago JE (2011) Influence of the adjuvants in a commercial formulation of the fungicide "Switch" on the adsorption of their active ingredients: cyprodinil and fludioxonil, on soils devoted. *J Hazard Mater* 193: 288-295.
7. Gnanamangai B, Ponmurugan P (2012) Evaluation of various fungicides and microbial based biocontrol agents against bird's eye spot disease of tea plants. *Crop Prot* 32: 111-118.
8. Patyal SK, Sharma ID, Chandel RS, Dubey JK (2013) Dissipation kinetics of trifloxystrobin and tebuconazole on apple (*Malus domestica*) and soil—A multi location study from north western Himalayan region. *Chemosph* 92: 949-954.
9. Yang D, Wang N, Yan X, Shi J, Zhang M, et al. (2014) Microencapsulation of seed-coating tebuconazole and its effects on physiology and biochemistry of maize seedlings. *Biointerf* 114: 241-246.
10. Pereyra MA, Ballesteros FM, Creus CM, Sueldo RJ, Barassi CA (2009) Seedlings growth promotion by *Azospirillum brasilense* under normal and drought conditions remains unaltered in Tebuconazole-treated wheat seeds. *Europ J Soil Bio* 45: 20-27.
11. Ara B, Chen Z, Shah J, Jan MR, Ye LJ (2012) Preparation and characterization of uniform molecularly imprinted polymer beads for separation of triazine herbicides. *Appl Polym Sci* 126: 315-321.
12. Beltran J, Lopez FJ, Forcada M, Homandez F (1997) Microextraction procedures combined with large volume injection in capillary gas chromatography for the determination of pesticide residues in environmental. *Anal chem Acta* 356: 125-133.
13. Breton F, Euzet P, Piletaky SA, Giardi MT, Rouillon R (2006) Integration of photosynthetic biosensor with molecularly imprinted polymer-based solid phase extraction cartridge. *Anal Chim Acta* 569: 50-57.
14. Martinez RC, Gonzalo ER, Ruiz PR, Alvarez JD (2003) Solid-phase extraction and sample stacking—micellar electrokinetic capillary chromatography for the determination of multiresidues of herbicides and metabolites. *J Chroma A* 990: 291-302.
15. Huertas-pere JF, Iruck MO, Campana AMG, Casado AG, Navarro ASJ (2006) *Chroma A* 1102: 250-286.
16. Shah J, Jan MR, Muhammad M, Shehzad F, Un-Nisa (2010) Flow injection spectrophotometric determination of fenoxaprop-p-ethyl herbicide in different grain samples after derivatization. *J Braz Chem Soc* 21: 10.
17. Jan MR, Shah J, Khan H (2003) Investigation of new indirect spectrophotometric method for the determination of carbofuran in carbamate pesticides. *Chemosphere* 1: 1623-1626.