

# The Immunohistochemistry for HPV infection in High Grade Squamous cell lesions of Esophagus Diagnosed on Endoscopic Biopsies

Purvee Agrawal<sup>1</sup> and Arvind Bhake

Department of Pathology, Jawaharlal Nehru Medical College, DMIMS (DU), Sawangi, Wardha- 442001, M.S, India

## Abstract

**Background:** The recent report in the literatures refers to the role of HPV in pathogenesis of esophageal cancer with geographical biases. There were not many reports from India where in relationship between HPV and esophageal squamous cell high grade lesions were analyzed. Such studies are required to explore the role of HPV in esophageal Squamous cell carcinoma for its implication of prevention and treatment.

**Aim:** The aim of study was to observe the pathogenetic relationship between high grade squamous cell lesion and HPV detected on the tissue biopsies by immunohistochemistry. The another objective of the study was to study relationship of HPV with grades of SCC.

**Methods:** Fifty-six paraffin embedded blocks that were diagnosed as high grade squamous cell lesions were selected and the immunostaining for HPV was performed using mouse monoclonal Anti-Human Papillomavirus (HPV) antibodies (DAKO code M3528) against HPV.

**Results:** The study group 56 cases endoscopic biopsies of high grade SCC showed HPV positive IHC staining in total of 12 cases (21.42%) of which 1 case (1.79%) was of carcinoma in-situ and 11(20%) was of invasive SCC. All the 11 cases of invasive SCC were of well differentiated SCC which showed positive HPV IHC staining. The p-value for the results of HPV for high grade squamous cell lesions of esophagus was 0.0022 which was significant.

**Conclusion:** There exists the association between HPV and esophagus high grade squamous cell lesions. These aspects of pathogenicity are required to be propped further at large sample studies to be carried out in divergent populations.

## Keywords

Squamous cell carcinoma • Human papilloma virus • Immunohistochemistry

## Introduction

Globally cancer of esophagus is one of the leading causes of cancer mortality. The commonest subtype of esophageal cancer is Squamous cell carcinoma and the other common one is Adenocarcinoma which is known sequelae to barretts' esophagus. The esophageal SCC in past two decades has been the topic of interest for researchers for its evaluation for etiopathogenesis. To the contexts of Indian population esophageal SCC by the figures of ICMR cancer registry occupies 8th rank with age standardized incidence rate (ASR) of 6.5 in 100000 population for males and 4.2 per lakh population for females of the total body cancer. Since the introduction of endoscopy at diagnostic evaluation of lesions of gastrointestinal tract, the endoscopic biopsies of esophageal lesions became an integral part and have enhanced the cancer detection rate of the esophageal lesions. The one subtype that has found to be the commonest chunk of malignant neoplastic lesions diagnosed on esophageal biopsies is SCC. There are several risk factors listed for esophageal SCC including viral oncogenes

is which implicates Epstein- Barr virus, Herpes Simplex virus and Human papilloma virus. Of these risk factors, HPV have been cited in many studies to have the causal role in etiopathogenesis of esophageal cancer. The literature on these aspects of HPV is mostly drawn from China, where the evidences have been generated for detection of HPV on the tissue of esophageal SCC by adapting to by various methods of detection such as PCR, IHC, ISH. However, there are scarce studies in India at evaluation of role of HPV in genesis of esophageal cancer. There exists a gap of understanding as Indian medical literature do not have many publications which has evaluated the role of HPV and its association with high grade squamous cell lesions of esophagus (SCC in situ, micro invasive SCC, invasive SCC) [1, 2].

The aim of study was to observe the pathogenetic relationship between high grade squamous cell lesion and HPV detected on the tissue biopsies by IHC. The another objective of the study was to study relationship of HPV with grades of SCC.

## Materials and Methods

Fiftysix paraffin embedded blocks that were diagnosed as high grade squamous cell lesions on histopathology were selected from Jawaharlal Nehru Medical College Sawangi during the period from August 2017 to October 2019. Patient identification data were retrieved from patients file including age, sex, tumor grade, endoscopic findings, and history of

HPV-6,11,16,18,31,33,42,51,52,56 and 58 was performed using mouse monoclonal Anti-Human Papillomavirus (HPV) antibodies (DAKO code M3528) against HPV [3].

The immunohistochemical staining was carried out on 4 um thickness section from formalin fixed, paraffin embedded tumor were cut and mounted onto poly-lysine coated slides followed by deparaffinization in xylene, Rehydration through descending grades of alcohol and was placed in distill

\*Address for Correspondence: Purvee Agrawal, Department of Pathology, Jawaharlal Nehru Medical College, DMIMS (DU), Sawangi, Wardha- 442001, M. S, India; E-mail: purvee.angel14@gmail.com

Copyright: © 2020 Agrawal P, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Received September 03, 2020; Accepted September 15, 2020; Published September 22, 2020

water for 1 min. samples were steamed for antigen retrieval in pressure cooker done for 15 minutes, Washed with distill water by giving a single dip. Transfer the slides into buffer (Tris Buffer Solution) for at least 5 minutes at room temperature (3 times wash). Peroxidase activity was blocked for 30 minutes using 3, 5 Hydrogen Peroxide plus methanol. Wash with buffer 3 times for 5 minutes each. Apply mouse monoclonal anti HPV antibody for HPV at room temperature and kept for 1 hour. Wash with buffer thrice for 5 minutes each at room temperature. Envision to be undertaken by labeled polymer for 30 minutes at room temperature. Apply chromogen 3, 3'-Diaminobenzidine (DAB) for 15-20 minutes. (Working DAB solution: 1 ml DAB buffer + 25 µL DAB concentrate). Wash with buffer 3 times each for 5 minutes. Wash with buffer for 10 minutes and slides were counter stain with harris hematoxylin, wash with running tap water, mount with DPX.

**Interpretation of IHC**

The negative and positive IHC results were interpreted and were scored with the help of references available in literature. Distinct granular brown discoloration was generally considered as positive staining for IHC. The IHC was interpreted for its positive or negative results on following 3 parameters using light microscopy with minimum of 1000 cells and screened under high power objective- Nuclear/cytoplasmic staining, Granular staining/intensity of signal, No. of cells showing granular signals in cytoplasm and nucleus.

The following scoring system for the Interpretation of IHC results for high risk HPV were used- For area scoring: 0 indicated no positive staining; 1%-10% of tissue stained; 2, 11%-50% of tissue stained; 3, 51%-75% of tissue stained; and 4, .75% of tissue stained. Cytoplasmic signal staining in absence of nuclear staining too is considered positive for interpretation of IHC [4].

**Static analysis**

The data obtained was analyzed using the statistical programs software Statistical package for the social sciences version (22.0), Fischer Exact test and Chi square test were used to calculate the association between detection of HPV by IHC with the diagnosis of high grade squamous cell lesions of esophagus i.e. SCC in-situ, invasive SCC along with grades involved in the study. P-value less than 0.05 (<0.05) was considered

statistically significant.

**Patient characteristic**

56 subjects (26) out of them were males (46.2%) and 30 (53.58%) were females, with (0.8:1) male to female ratio, the patient age ranged between 31-90 years, With the mean age about 59±1.48 years. The majority of high grade squamous cell lesions patients were observed in the age range of 51-60 years as shown in Table 1.

**Histomorphological diagnosis**

Of total 56 subjects, out of which one case of carcinoma in situ and 55 were of invasive SCC. The frequency of invasive SCC were 30 (53.57%) well differentiated SCC, 9 (16.07%) moderately differentiated SCC, 16(28.57%) poorly differentiated SCC.

**HPV IHC status and histomorphological diagnosis**

The study group of 56 cases endoscopic biopsies of high grade SCC showed HPV positive IHC staining in total of 12 cases (21.42%) of which 1 case (1.79%) was of carcinoma in-situ and 11(20%) was of invasive SCC. There were scores of IHC staining for HPV and in 12 cases interpreted as positive revealed that all the had strong positive signals with (p=0.029) and was significant as shown in Table 2.

**Grading of invasive SCC**

Of the total 55 cases of invasive SCC, the maximum biopsies 30(53.57%) showed grade I well differentiated, 09(16.07%) grade II moderately differentiated and 16 (28.57%) grade III poorly differentiated, grade of invasive SCC [5, 6].

**HPV IHC status and grading of SCC**

The 11 cases of well differentiated SCC showed positive HPV IHC staining; no case from the category of moderate or poorly differentiated SCC showed positive results for IHC for HPV Figure 1. A single case of SCC Carcinoma in situ showed positive immunostaining for HPV IHC Figure 2, p-value was 0.0022 for this observation and was significant as shown in Table 3.

Age(years)	Male	Female	Total	Percentage %	
31-40	02	2	4	7.14%	0.75
41-50	01	10	11	19.64%	0.75
51-60	10	08	18	32.14%	0.75
61-70	10	06	16	28.57%	0.75
71-80	02	04	06	10.71%	0.75
81-90	01	0	01	1.79%	0.75
Total	26	30	56	100%	0.75


Table 1. Age and gender distribution of subjects (n=56).

Diagnosis	Total	HPV (IHC) status				Positive %
		Negative		Positive		
		Male	Female	Male	Female	
Study group (SCC)N=56	CIS	1	0	0	1	100%
	Invasive SCC	55	22	22	4	7
Control group N=20	26	26	26	26	26	26
P value	Fischer's Exact test p-value=0.029					


Table 2. IHC results for HPV in SCC cases (n=56) and control group (n=20).

HPV (IHC) status	Grades of SCC				Total	value 2x
	Carcinoma situ	Well differentiated	Moderately differentiated	Poorly differentiated		
Negative	0	19	9	16	44	55
					(78.57%)	
Positive	1	11	0	0	12	11.62
					(21.42%)	P=0.0022,S
Total	1	30	9	16	56	

**Table 3.** HPV (IHC) status and its relationship with broder's grades of SCC (n-56).



**Figure 1.** HPV positive (carcinoma in situ) shows brown discrete staining of cytoplasmic and occasional nuclei (anti HPV monoclonal antibody stain) intensity score 3+, 40X.



**Figure 2.** HPV positive (well differentiated SCC esophagus) shows brown discrete staining of cytoplasmic and occasional nuclei ( anti HPV monoclonal antibody stain) intensity score 3+, (40X).

## Results and Discussion

The distribution of patients of age ranges in the present study revealed that the age range of 41-50 years showed the maximum number of cases of SCC followed by 51-60 years. The similar observation for age for esophageal SCC have been observed in the studies Youngest patient in the study was 35 years and the incidence of SCC in esophagus before 4th decade of life was observed to be 7.2% in the present study. The SCC occurring in the lower age of 4th decade of life has also been reported in the studies.

Old age in itself is a risk factor for development of SCC has been quoted in the text. The present study observed 23 patients were older, that is past of 6th decade of life. Similar observations for occurrence of SCC in

old age has made in the studies, 45% (47/103) were the patients belonged to advanced age.

The gender bias has been recorded in most of the studies reviewed for the present work which observed that male suffered more of SCC than females and attributed the results for risk factor like tobacco user, alcohol and others. The study of these mentioned (M:F) ratio for SCC as (4:1), (2.3:1), (3:1), (4:1) for studies. The present study has contradictory observation to that of above studies the M: F ratio was almost approaching to equality (0.8:1).

There is an observation unusual for one of the female who was positive for HPV on IHC for esophageal SCC on biopsy and had simultaneous investigation of cervical cytology which shows HPV associated have screened women with esophageal SCC with positive IHC for HPV cervical

SCC too and have found to be concomitant existence of HPV.

The present study have concordant finding (21.43%) with the studies. The present study did not observe the very high positive % for HPV as has been documented in the few studies originating in China for the reason that the prevalence of HPV or viral antigenicity in the subjects within the present study from the central India probably has low endemicity of prevalence of HPV, too have reported low percentile of HPV antibody (3%) by IHC which is still lower for detection rate of HPV of the present study.

## Conclusion

The observation in the present study for association between HPV and squamous cell lesion including SCC implicates there causal relationship this aspect of pathogenicity is required to be further studied as a large sample at multiple center engaging the workup of esophageal SCC.

## References

1. Hardefeldt HA, Cox MR and Eslick GD. "Association between human papillomavirus (HPV) and oesophageal squamous cell carcinoma: a meta-analysis." *Epidemiol Infect* 142(2014):1119-1137.
2. Liyanage, Surabhi S, Segelov Eva, Garland Suzanne M and Tabrizi Sepehr N, et al. "Role of human papillomaviruses in esophageal squamous cell carcinoma: Role of HPV in esophageal SCC." *Asia-Pac J ClinOncol* (2013).
3. Ludmir, Ethan B, Stephens Sarah J, Palta Manisha and Willett Christopher G, et al. "HPV infections and oesophageal cancer." *J ClinPathol* (2002).
4. Mir, M Muzaffar and Dar Nazir Ahmad. "Esophageal Cancer in Kashmir (India): An Enigma for Researchers." *Int J Health Sci (Qassim)* 3(2009):71-85.
5. Shuyama K, Castillo A, Aguayo F and Sun Q, et al. "Human papillomavirus in high- and low-risk areas of oesophageal squamous cell carcinoma in China." *Br J Cancer* 96(2007):1554-1559.
6. Xi, Ruxing, Zhang Xiaozhi, Chen Xin and Pan Shupe, et al. "Human papillomavirus 16 infection predicts poor outcome in patients with esophageal squamous cell carcinoma." *Onco Targets Ther* 8(2015):573-581.

**How to cite this article:** Agarwal P and Bhake A. "The Immunohistochemistry for HPV infection in High Grade Squamous cell lesions of Esophagus Diagnosed on Endoscopic Biopsy." *J Surg Path Diag* 2 (2020).