Understanding Solvent Effects in the Solvolyses of 4-Fluorophenyl Chlorothionoformate

*MJ D'Souza¹, SM Hailey¹, BP Mahon¹, *DN Kevill²

¹ Department of Chemistry, Wesley College, 120 N. State Street, Dover, Delaware 19901-3875, USA.

² Department of Chemistry and Biochemistry, Northern Illinois University, DeKalb, Illinois 60115-2862, USA.

* Correspondence to: MJ D'Souza, dsouzama@wesley.edu; DN Kevill, dkevill@niu.edu

Accepted: April 8, 2011; Published: June 27, 2011

Abstract

The solvolyses of 4-fluorophenyl chlorothionoformate (1) are studied in fifteen binary aqueous organic mixtures of widely varying nucleophilicity and ionizing power values. The specific rates of solvolyses of 1 are plotted against the specific rates of solvolysis observed for phenyl chloroformate (2) and deviations from the line-of-best-fit are observed in some of the highly ionizing aqueous fluoroalcohol mixtures. An analysis of the solvolytic data accumulated using the extended (two-term) Grunwald-Winstein equation confirms this deviant behavior and shows that dual bimolecular addition-elimination and unimolecular ionization channels occur in the solvolyses of 1.

Keywords: Solvolysis; Grunwald-Winstein equations; nucleophilic solvation; chlorothionoformate.

1. Introduction

Compounds such as 4-fluorophenyl chlorothionoformate (1) that contain fluorophenyl groups are important in asymmetric synthesis in medicinal chemistry, as fluorine substitution in a drug molecule can alter the pharmacodynamic activity and toxicity of biologically active drug molecules [1]. Chlorothionofomate esters in general are also employed as useful reagents for the preparation of thiocarbonate esters, nitriles, and isonitriles [2,3].

As a result a number of research groups [4-19] are now involved in the investigation of the hydrolysis, aminolysis and alcoholysis of alkyl and aryl chlorothionoformates. Queen [4], in an analysis of the hydrolysis of a series of alkyl chlorothionoformates together with their activation parameters and solvent isotope effects, concluded that these compounds hydrolyze by S_N1 mechanisms. Lee et al. [5,6,13] favored a concerted process with a general base catalyzed S_N2 transition state for the solvolysis of alkyl and aryl chlorothionoformates, and a concerted mechanism [15] with a four-membered hydrogen bonded cyclic transition state for the aminolysis of aryl chlorothionoformates with anilines in acetonitrile. Castro, Santos and coworkers [7,8,11,14,16,17] proposed that the aminolysis of chlorothionoformates is a stepwise process with the formation of a zwitter ionic intermediate while the phenolysis is concerted.

We favor the use [19] of the extended Grunwald-Winstein equation (equation 1) [20] for solvolytic studies and have proposed that side-by-side addition-elimination $(A_N + D_N)$ and $S_N 1$ ionization $(D_N + A_N)$ mechanisms often occur during the solvolysis of chloro- (ROCOCI), chlorothio- (RSCOCI), chlorothiono- (ROCSCI), and chlorodithioformate esters (RSCSCI) [10,18,19,21,22].

$$\log (k/k_o) = IN + mY + c \tag{1}$$

In equation 1, k and k_o are the specific rates of solvolysis of a substrate in a given solvent and the standard solvent (80% ethanol) respectively, I is the sensitivity to changes in solvent nucleophilicity (M), m represents the sensitivity that controls the importance of the solvent ionizing power value (Y). N_T scales [23] based on the solvolyses of the S-methyldibenzothiophenium ion and Y_x values based on the solvolyses of 1- and 2-adamantyl

derivatives [24] have become the recognized standards for considerations of the Grunwald-Winstein linear free energy relationship (LFER).

Based on results obtained through Grunwald-Winstein (equation 1) analyses, we recently recommended [19] that the *I* (1.66) and *m* (0.56) values obtained for solvolyses of phenyl chloroformate (**2**), and the *I* = 0.69, *m* = 0.95 values obtained for phenyl chlorodithioformate (**3**), be taken as appropriate standards for the bimolecular addition-elimination ($A_N + D_N$) and unimolecular S_N 1 ionization ($D_N + A_N$) pathways respectively. The *I* value of 0.69 seen in the chlorodithioformate ester was said to represent a strong rear-side nucleophilic solvation of the developing thioacylium cation [19].

We also pointed out [19] that in RXCXCI (X = O or S) substrates superimposed mechanisms were observed and the ranges of dominance for each was dependent on the R group, if X = O or S, the solvent nucleophilicity value, and the solvent ionizing power value.

Lee [25] analyzing CNDO/2 studies on conformation and reactivity suggested that the most stable geometries for RXCXCI (where X = S or O) involve a configuration where the C=O or the C=S is *syn* with respect to R. Bentley and others [26] confirmed this conformation and in Figure 1, the molecular structures for 4-fluorophenyl chlorothionoformate (1), phenyl chloroformate (2), phenyl chlorodithioformate (3), phenyl chlorothionoformate (5), and phenyl fluorothionoformate (6) are shown where the halogen atom is in a *trans* position with respect to the aryl group i.e. the structures are drawn in a *syn* conformation.



Figure 1: Molecular structures of 4-fluorophenyl chlorothionoformate (1), phenyl chloroformate (2), phenyl chlorodithioformate (3), phenyl chlorothioformate (4), phenyl chlorothionoformate (5) and phenyl fluorothionoformate (6).

Kyong, and Kevill et al. [27] have recently showed that phenyl fluorothionoformate (**6**) has an *l/m* ratio of 3.38, a solvent deuterium isotope effect value for methanolysis (k_{MeOH}/k_{MeOD}) of 2.11, and entropy of activation range of -26.2 to -21.0 cal mol⁻¹K⁻¹. This data strongly indicated that **6** solvolyzes by a bimolecular pathway in all solvents studied with the addition step of the addition-elimination ($A_N + D_N$) reaction being rate-determining [27].

In the present paper, we have reported the first-order specific rate constants at 35.0°C for the solvolyses of **1** in ethanol and methanol and thirteen binary mixtures of aqueous ethanol (EtOH), aqueous methanol (MeOH), aqueous acetone, aqueous 2,2,2-trifluoroethanol (TFE), and aqueous 1,1,1,3,3,3-hexafluoro-2-propanol (HFIP). Applying the concept of a similarity model [28] we compare the specific rates of solvolysis of **1** against those of **2**, examine the solvent-deviations from the goodness-of-fit line, and justify the results obtained by employing correlation analyses using the two-term Grunwald-Winstein LFER (equation 1).

2. Methods

The 4-fluorophenyl chlorothionoformate (96%, TCI America) was used as received. Solvents were purified and the kinetic runs carried out as described previously [19]. A substrate concentration of approximately 0.005 M

in a variety of solvents was employed. For some of the runs, calculation of the specific rates of solvolysis (firstorder rate coefficients) was carried out by a process [19] in which the conventional Guggenheim treatment was modified so as to give an estimate of the infinity titer, which was then used to calculate for each run a series of integrated rate coefficients. The specific rates and associated standard deviations, as presented in Table 1, were obtained by averaging all of the values from, at least, duplicate runs.

Multiple regression analyses were carried out using the Excel 2007 package from the Microsoft Corporation, and the SigmaPlot 9.0 software version from Systat Software, Inc., San Jose, CA, was used for the Guggenheim treatments.

3. Results and Discussion

The first-order specific rates of solvolysis of **1** at 35.0 °C in fifteen pure and aqueous organic mixtures are reported in Table 1. The specific rates of **1** increase as the amount of water in the binary aqueous mixture increases. A small gradual increase in rate coefficients is also observed in the highly ionizing fluoroalcohols indicating that in these solvents, there is some nucleophilic participation of the solvent in the transition state.

Table 1: Specific rates of solvolysis (*k*) of **1**, in several binary solvents at 35.0 °C and literature values for N_7 and Y_{Cl} .

Solvent (%) ^a	1 @ 35.0 °C; 10 ⁵ k,	N _T ^c	Y _{CI} ^d
	3		
100% MeOH	55.3 ± 1.3	0.17	-1.2
90% MeOH	86.0 ± 1.3	-0.01	-0.20
80% MeOH	116 ± 6	-0.06	0.67
100% EtOH	20.0 ± 1.0	0.37	-2.50
90% EtOH	24.8 ± 0.5	0.16	-0.90
80% EtOH	28.2 ± 1.8	0.00	0.00
80% Acetone	3.44 ± 0.06	-0.35	-2.39
97% TFE (w/w)	0.467 ± 0.020	-3.30	2.83
90% TFE (w/w)	0.600 ± 0.010	-2.55	2.85
70% TFE (w/w)	2.08 ± 0.13	-1.98	2.96
80T-20E	0.272 <u>+</u> 0.010	-1.76	1.89
50T-50E	3.21 ± 0.14	-0.64	0.60
97%HFIP (w/w)	3.43 ± 0.09	-5.26	5.17
90%HFIP (w/w)	4.18 ± 0.08	-3.84	4.41
70%HFIP (w/w)	5.49 ± 0.07	-2.94	3.83

^a Substrate concentration of *ca.* 0.0052 M; binary solvents on a volume-volume basis at 25.0 °C, except for TFE-H₂O and HFIP-H₂O solvents which are on a weight-weight basis. T-E are TFE-ethanol mixtures. ^b With associated standard deviation. ^cRefs. 19,23. ^dRefs. 19,24.

A plot of log $(k/k_0)_1$ versus log $(k/k_0)_2$ shown in Figure 2, exhibits considerable deviations for some of the aqueous fluoroalcohol mixtures. Indeed this plot results in a very poor correlation coefficient (*R*) of 0.720, slope of 0.33 \pm 0.09, intercept (*c*) of -0.16 \pm 0.20, and *F*-test of 14.

Using all of the fifteen specific rates of solvolysis of **1** listed in Table 1 within the extended Grunwald-Winstein equation (equation 1) leads to a very poor correlation with a correlation coefficient (*R*) of 0.627, $I = 0.43 \pm 0.33$, $m = 0.11 \pm 0.25$, $c = -0.17 \pm 0.27$, and a very low *F*-test value of 4. This substandard correlation result expresses the possibility of the presence of dual mechanisms in the solvolyses of **1**.

A careful observation of Figure 2 indicates that 97%, 90%, and70% HFIP, and 97% and 90% TFE deviate significantly from the line of best fit. Removal of these five highly ionizing solvents for a correlation analysis of **1** against **2**, results in a significantly improved correlation coefficient (R) of 0.994, slope of 0.92 ± 0.04 , $c = 0.08 \pm 0.04$, and an *F*-test value of 622. This exceptional correlation suggests that the mechanism of **1** in the remaining ten more nucleophilic solvents is exactly the same as that proposed for **2**. The slope of 0.92

observed in the plot of log $(k/k_0)_1$ versus log $(k/k_0)_2$ in these ten solvents, is consistent with the proposal of a slightly earlier transition state for the solvolyses of **1** when compared to that seen for the solvolyses of **2**.



Figure 2: The plot of log (k/k_o) for 4-fluorophenyl chlorothionoformate (1) against log (k/k_o) for phenyl chloroformate (2) in the fifteen common pure and binary solvents studied.

In Table 2, we report the results obtained on the application of equation 1 to the specific rates of solvolyses of 1 using the ten more nucleophilic solvents. We obtain an *I* value of 1.76 ± 0.28 , an *m* value of 0.54 ± 0.15 , a *c* value of 0.34 ± 0.15 , an *F*-test value of 28, and an *R* value of 0.943. For **2** in the identical set of ten solvents we obtain, $I = 1.89 \pm 0.24$, $m = 0.55 \pm 0.13$, $c = 0.27 \pm 0.13$, *F*-test = 46, and R = 0.964. The similarities of the *I* and *m* strongly indicates that **1** and **2** solvolyze by the same mechanism in these ten solvents and the magnitude of the *I* and *m* values are consistent with the addition step of an addition-elimination ($A_N + D_N$) mechanism being rate determining (Scheme 1). The slightly earlier tetrahedral transition state (Scheme 1) favored for **1** (when compared to that of **2**) is due to the polarizability of the sulfur atom and its ability to distort the shape of its electron cloud.

		Giuni	valu-vvilistelli eq				
Substrate	n ^a	ľ	m ^D	C ^C	R ^a	F ^e	Mechanism
1	10	1.76 ± 0.28	0.54 ± 0.15	0.34 ± 0.15	0.943	28	A-E
	5	0.53 ± 0.18	0.89 ± 0.18	-2.66 ± 0.35	0.967	15	S _N 1
2 ^{<i>t</i>}	49	1.66 ± 0.05	0.56 ± 0.03	0.15 ± 0.07	0.980	568	A-E
3^{g}	31	0.69 ± 0.05	0.95 ± 0.03	0.18 ± 0.05	0.987	521	S _N 1
4 ^{<i>h</i>}	16	1.74 ± 0.17	0.48 ± 0.07	0.19 ± 0.23	0.946	55	A-E
	6	0.62 ± 0.08	0.92 ± 0.11	-2.29 ± 0.13	0.983	44	S _N 1
5 ^{<i>g</i>}	9	1.88 ± 0.28	0.56 ± 0.15	0.38 ± 0.15	0.950	28	A-E
	18	0.34 ± 0.05	0.93 ± 0.09	-2.54 ± 0.34	0.955	77	S _N 1
6'	22	1.32 ± 0.13	0.39 + 0.08	-0.02 ± 0.10	0.952	95	A-F

Table 2: Correlations of the specific rates of solvolysis of a variety of 1, 2, 3, 4, 5, and 6 using the extendedGrunwald-Winstein equation (equation 1).

^a *n* is the number of solvents. ^bWith associated standard error. ^cThe earlier values are accompanied by standard error of the estimate. ^dCorrelation coefficient. ^e*F*-test value. ^fValues taken from ref. 18. ^gValues taken from ref. 10. ^hValues taken from ref. 29. ⁱValues taken from ref. 27.



Scheme 1: Stepwise addition-elimination mechanism through a tetrahedral intermediate for chlorothionoformate esters.

At 25.0 °C, the rates of solvolysis in 100% EtOH for **1**, **2**, **3**, **4**, **5**, and **6** are $11.9 \pm 1.1 \times 10^{-5} s^{-1}$, $260 \pm 3 \times 10^{-5} s^{-1}$ [18], $0.996 \pm 0.02 \times 10^{-5} s^{-1}$ [10], $6.73 \pm 0.15 \times 10^{-5} s^{-1}$ [29], $3.15 \pm 0.04 \times 10^{-5} s^{-1}$ [10], and 1759 $\times 10^{-5} s^{-1}$ (estimated from the Arrhenius activation parameters reported in ref. 27) respectively. For substrates **1-6** it is proposed that the addition-elimination mechanism dominates in pure ethanol (Table 2). The corresponding ethanolysis rate order of $k_6 > k_2 > k_1 > k_4 > k_5 > k_3$ demonstrates the powerful inductive effects instigated by the presence of both a phenoxy group and a fluorine leaving group around the carbonyl carbon in **6** when compared to the electron-withdrawing effects witnessed in the tetrahedral addition-elimination transition-state of the other substrates. The observation that **1** is approximately four times faster than **5** in pure ethanol is consistent with a slightly positive Hammett σ_p value of + 0.06 [30] for *para*-F, implying a higher electron-withdrawing power of the *para*-substituted fluorine atom in **1** relative to the H in the *para* position in **5**.

A plot of log (k/k_o) for 4-fluorophenyl chlorothionoformate (1) against 1.76 N_T + 0.54 Y_{Cl} in the fifteen common pure and binary solvents studied is shown in Figure 3. The points for the 97, 90, 70 HFIP and 97, 90 TFE were not included in the correlation. They were added to show the extent of their deviation.

Application of equation 1 to the solvolysis of **1** in these five highly ionizing deviating fluoroalcohols results in $I = 0.53 \pm 0.18$, $m = 0.89 \pm 0.18$, $c = -2.66 \pm 0.35$, *F*-test = 15, and R = 0.967 (Table 2). A large negative *c* value is observed because the experimental k_0 value is the one applying to the other reaction channel. The large errors associated with the *I* and *m* values and the low *F*-test value are due to the very small number of solvents (five in this case) used for dual parameter analyses [19]. However, the ranges of these *I* and *m* values are very similar to those observed for the solvolyses of **3**, where an S_N1 ionization (D_N + A_N) mechanism with strong rear-side nucleophilic solvation was proposed.

We now assert that 1 undergoes solvolysis in 97%, 90%, and 70% HFIP, and 97% and 90% TFE, by an ionization $(S_N 1)$ mechanism with strong nucleophilic solvation of the resonance stabilized carbocation (Scheme 2). This proposal is in agreement with the slight increase in the specific rates of solvolysis that is observed as the water content increases (due to an increase in the nucleophilicity) in these mixtures.

On the other hand, it was shown [18] that **5** solvolyzes by the ionization pathway (Scheme 2) in a wider variety of ionizing solvents (when compared to the range of ionization observed in **1**) including the aqueous alcohols and water. This is due to the sulfur atom's lone-pair electron polarizability (in **5**) being more diffuse in the absence of the electron withdrawing fluorine atom in the *para* position, hence its ability to favor the resonance contributor 2b in Scheme 2.

4. Conclusion

4-Fluorophenyl chlorothionoformate (1) was shown to solvolyze by dual bimolecular addition-elimination ($A_N + D_N$) and unimolecular ionization ($D_N + A_N$) mechanisms whose dominance depended on the nucleophilicity and ionizing power of the solvent. The exact solvent compositions where the change in mechanism occurred were identified utilizing the similarity model concept and the extended Grunwald-Winstein equation.

5



Figure 3: The plot of log (k/k_o) for 4-fluorophenyl chlorothionoformate (1) against 1.76 N_T + 0.54 Y_{CI} in the fifteen common pure and binary solvents studied.



Scheme 2: Unimolecular step-wise solvolysis of chlorothionoformate esters.

Abbreviations

S _N 1	substitution nucleophilic unimolecular
S _N 2	substitution nucleophilic bimolecular
$A_N + D_N$	association-dissociation
$D_N + A_N$	dissociation-association
CNDO/2	"semi-empirical" all-valence electron-self-consistent-field theory
σ _p	Hammett sigma constant for <i>para</i> substituent
DE-INBRE	Delaware-IDeA Network of Biomedical Research Excellence

Competing Interests

The authors declare that they have no competing interests. **Authors' Contributions**

SMH and BPM completed this research under the direction of MJD as undergraduate research assistants in the DE-INBRE sponsored Wesley College Directed Research Program. DNK is a collaborator on this project.

Acknowledgement

This research was supported by grant number 2 P2O RR016472-011 from the National Center for Research Resources (NCRR), a component of the National Institutes of Health (NIH). This IDeA Network of Biomedical Research Excellence (INBRE) grant to the State of Delaware (DE) was obtained under the leadership of the University of Delaware, and the authors sincerely appreciate their efforts.

References

- Park K, Kitteringham NR, O'Neil PM, 2001. Metabolism of fluorine-containing drugs. Pharmacology and [1]. Toxicology, 41: 443-470.
- [2]. Bose DS, Goud, PR, 1999. Aryl chlorothionoformates: A new versatile reagent for the preparation of nitriles and isonitriles under mild conditions. Tetrahedron Letters, 40: 747-748.
- Rahmathalluah SM, Hall JE, Bender BC, McCurdy DR, Tidwell RR, Boykin DW, 1999. Prodrugs for [3]. Amidines: Synthesis and anti-pneumocystis carinii activity of carbamates of 2,5-bis(4amidinophenyl)furan. Journal of Medicinal Chemistry, 42: 3994-4000.
- McKinnon DM, Queen, A, 1972. Kinetics and mechanism for the hydrolysis of chlorothionoformates [4]. and chlorodithioformate esters in water and aqueous acetone. Canadian Journal of Chemistry, 50: 1401-1406.
- La S, Koh KS, Lee I, 1980. Nucleophilic substitution at a carbonyl carbon atom (XI). Solvolysis of methyl [5]. chloroformate and its thioanalogues in methanol, ethanol and ethanol-water mixtures. Journal of Korean Chemical Society, 24: 1-7.
- La S, Koh KS, Lee I, 1980. Nucleophilic substitution at a carbonyl carbon atom (XII). Solvolysis of methyl [6]. chloroformate and its thioanalogues in CH₃CN-H₂O and CH₃COCH₃-H₂O mixtures. Journal of Korean Chemical Society, 24: 8-14.
- [7]. Castro EA, Cubillos M, Santos JG, 1997. Kinetics and mechanism of the aminolysis of phenyl and 2nitrophenyl chlorothionoformates. Journal of Organic Chemistry, 62: 4395-4397.
- [8]. Castro EA, Cubillos M, Santos JG, 1998. Concerted mechanisms of the reactions of phenyl and 4nitrophenyl chlorothionoformates with substituted phenoxide ions. Journal of Organic Chemistry, 63: 6820-6823.
- Millan DS, Prager, RH, 1998. The dealkylation of tertiary aliphatic amines with phenyl [9]. chlorothionoformates. Tetrahedron Letters, 39: 4387-4390.
- Kevill DN, D'Souza MJ, 1999. Correlation of the rates of solvolysis of phenyl chlorothionoformate and [10]. phenyl chlorodithioformate. Canadian Journal of Chemistry, 77: 1118-1122.
- [11]. Castro EA, 1999. Kinetics and mechanisms of reactions of thiol, thiono, and dithio analogues of carboxylic esters with nucleophiles. Chemical Reviews, 99: 3505-3524.

- [12]. Millan DS, Prager, RH, 1999. Phenyl chloro(thionoformate): a new dealkylating agent of tertiary amines. Australian Journal of Chemistry, 52: 841-850.
- [13]. Koo IS, Yang K, Kang DH, Park HJ, Kang K, Lee I, 1999. Transition-state variation in the solvolyses of phenyl chlorothionoformate in alcohol-water mixtures. Bulletin of Korean Chemical Society, 20: 577-580.
- [14]. Castro EA, Cubillos M, Santos JG, 2004. Kinetics and mechanisms of the pyridinolysis of phenyl and 4nitrophenyl chlorothionoformates. Formation and hydrolysis of 1-(aryloxythiocarbonyl)pyridinium cations. Journal of Organic Chemistry, 69: 4802-4807.
- [15]. Oh HK, Ha JS, Sung DD, Lee I, 2004. Aminolysis of aryl chlorothionoformates with anilines in acetonitrile: Effects of amine nature and solvent on the mechanism. Journal of Organic Chemistry, 69: 8219-8223.
- [16]. Castro EA, Aliaga M, Campodonico PR, Leis JR, Garcia-Rio L, Santos JG, 2007. Reactions of aryl chlorothionoformates with quinuclidines. A kinetic study. Journal of Physical Organic Chemistry, 21: 102-107.
- [17]. Castro EA, 2007. Kinetics and mechanisms of reactions of thiol, thiono and dithio analogues of carboxylic esters with nucleophiles. An update. Journal of Sulfur Chemistry, 28: 407-435.
- [18]. Kevill DN, Koyoshi F, D'Souza MJ, 2007. Correlation of the specific rates of solvolysis of aromatic carbamoyl chlorides, chloroformates, chlorothionoformates, and chlorodithioformates revisited. International Journal of Molecular Sciences, 8: 346-352.
- [19]. Kevill DN, D'Souza MJ, 2008. Sixty years of the Grunwald-Winstein equation: Development and recent applications. Journal of Chemical Research, 61-66.
- [20]. Winstein S, Grunwald E, Jones HW, 1951. The correlation of solvolyses rates and the classification of solvolysis reactions into mechanistic categories. Journal of the American Chemical Society, 73: 2700-2707.
- [21]. D'Souza MJ, Hailey SM, Kevill DN, 2010. Use of empirical correlations to determine solvent effects in the solvolysis of s-methyl chlorothioformate. International Journal of Molecular Sciences, 11: 2253-2266.
- [22]. D'Souza MJ, Mahon BP, Kevill DN, 2010. Analysis of the nucleophilic solvation effects of isopropyl chlorothioformate solvolysis. International Journal of Molecular Sciences, 11: 2597-2611.
- [23]. Kevill DN, 1996. Development and uses of scales of solvent nucleophilicity. In *Advances in Quantitative Structure-Property Relationships; Volume 1*. Edited by Charton M. Greenwich, CT: JAI Press, 81-115.
- [24]. Bentley TW, Llewellyn G, 1990. Yx scales of solvent ionizing power. Progress in Physical Organic Chemistry, 17: 121-158.
- [25]. Lee I, 1972. Nucleophilic substitution at a carbonyl carbon. Part II. CNDO/2 studies on conformation and reactivity of the thio-analogues of methyl chloroformate. Journal of Korean Chemical Society, 16: 334-340.
- [26]. Bentley TW, 2008. Structural effects on the solvolytic reactivity of carboxylic and sulfonic acid chlorides. Comparisons with gas-phase data for cation formation. Journal of Organic Chemistry, 73: 6251-6257.
- [27]. Choi SH, Seong MH, Lee Y-W, Kyong JB, Kevill DN, 2011. Correlation of the rates of solvolysis of phenyl fluorothionoformate. Bulletin of Korean Chemical Society, 32: 1268-1272.

- [28]. Bentley TW, Harris HC, Zoon H-R, Gui TL, Dae DS, Szajda SR. 2005. Mechanisms of solvolyses of acid chlorides and chloroformates. Chloroacetyl and phenylacetyl chloride as similarity models. Journal of Organic Chemistry, 70: 8963–8970.
- [29]. Kevill DN, Bond MW, D'Souza MJ, 1997. Dual pathways in the solvolyses of phenyl chlorothioformate. Journal of Organic Chemistry, 62: 7869-7871.
- [30]. Hansch C, Leo A, 1979. Substituent constants for correlation analysis in chemistry and biology. New York: Wiley-Interscience.

http://astonjournals.com/csj