Understanding the Main Disorders of the Gastrointestinal Tract in Older Adults

Michele Gajewski*

Department of Internal Medicine, Northwestern University, Evanston, Chicago, USA

Introduction

The Gastrointestinal (GI) tract plays a crucial role in the digestion and absorption of nutrients, as well as in the overall health of individuals. However, as people age, the GI tract undergoes various changes that can predispose older adults to different disorders. These disorders can significantly impact their quality of life and overall health. In this article, we will explore the main disorders of the gastrointestinal tract in older people, their causes, symptoms, diagnosis and management. Gastro Esophageal Reflux Disease (GERD) occurs when the lower esophageal sphincter weakens or relaxes inappropriately, allowing stomach acid to flow back into the esophagus [1]. Factors such as obesity, hiatal hernia and certain medications can contribute to its development. Heartburn, regurgitation, chest pain, difficulty swallowing and chronic cough are common symptoms of GERD in older adults. Diagnosis involves a combination of patient history, symptom assessment, endoscopy and pH monitoring. Management strategies include lifestyle modifications, dietary changes, medications (e.g., proton pump inhibitors) and, in severe cases, surgery.

Description

Peptic ulcers are primarily caused by infection with Helicobacter pylori bacteria or the chronic use of Non-Steroidal Anti-Inflammatory Drugs (NSAIDs). Aging-related changes in the gastric mucosa may also contribute to ulcer formation. Symptoms include abdominal pain, bloating, nausea, vomiting and gastrointestinal bleeding (which can lead to anemia). Diagnosis involves endoscopy, biopsy and H. pylori testing. Treatment includes antibiotics to eradicate H. pylori, proton pump inhibitors and avoidance of NSAIDs. Diverticular disease is characterized by the formation of diverticula (small pouches) in the colon wall. Aging-related changes in the colonic wall, such as decreased elasticity and muscle tone, contribute to diverticula formation [2]. Many older adults with diverticular disease remain asymptomatic, but symptoms can include abdominal pain, bloating, changes in bowel habits and rectal bleeding. Diagnosis and Management: Diagnosis is typically made through colonoscopy, CT scan, or barium enema. Management includes dietary fiber supplementation, adequate fluid intake and, in severe cases, antibiotics or surgery to treat complications such as diverticulitis. Colorectal cancer is thought to arise from a combination of genetic, environmental and lifestyle factors. Age is a significant risk factor, with most cases diagnosed in individuals over 50. Symptoms can vary but may include rectal bleeding, changes in bowel habits, abdominal pain, weakness and unintended weight loss. Diagnosis and Management: Screening for colorectal cancer in older

*Address for Correspondence: Michele Gajewski, Department of Internal Medicine, Northwestern University, Evanston, Chicago, USA; E-mail: gajewskim42@yahoo.com

Copyright: © 2024 Gajewski M. This is an open-access article distributed under the terms of the creative commons attribution license which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Received: 19 January, 2024, Manuscript No. jdcm-24-129732; **Editor Assigned:** 22 January, 2024, PreQC No. P-129732; **Reviewed:** 05 February, 2024, QC No. Q-129732; **Revised:** 10 February, 2024, Manuscript No. R-129732; **Published:** 17 February, 2024, DOI: 10.37421/2475-3211.2024.9.248

adults is essential and may include colonoscopy, fecal occult blood testing and sigmoidoscopy. Treatment options depend on the stage of the cancer but may involve surgery, chemotherapy, radiation therapy, or a combination thereof. Older adults are more susceptible to gastrointestinal infections due to age-related changes in the immune system and underlying health conditions. Bacterial (e.g., Clostridium difficile), viral (e.g., norovirus) and parasitic infections can occur [3].

Symptoms include diarrhea, abdominal cramps, nausea, vomiting, fever and dehydration. Diagnosis involves stool testing to identify the causative agent. Treatment includes fluid and electrolyte replacement, antimicrobial therapy (if bacterial) and symptomatic relief. Constipation is common in older adults and can be caused by various factors, including decreased gastrointestinal motility, inadequate dietary fiber intake, dehydration, immobility and certain medications. Infrequent bowel movements, difficulty passing stools, straining, abdominal discomfort and bloating are characteristic symptoms. Diagnosis is based on clinical assessment, including stool frequency and consistency. Management involves dietary modifications (increased fiber intake), adequate hydration, regular exercise and, if necessary, laxatives or stool softeners [4,5].

Conclusion

The gastrointestinal tract undergoes significant changes as people age, predisposing older adults to various disorders such as GERD, peptic ulcer disease, diverticular disease, colorectal cancer, gastrointestinal infections and constipation. Early recognition and management of these disorders are crucial for maintaining the health and quality of life of older adults. Healthcare providers should be vigilant in screening, diagnosing and treating GI disorders in this vulnerable population, taking into account age-related physiological changes and comorbidities. Additionally, promoting healthy lifestyle habits and preventive measures can help reduce the risk of GI disorders in older adults.

Acknowledgement

None.

Conflict of Interest

None.

References

- Soenen, Stijn, Christopher K. Rayner, Karen L. Jones and Michael Horowitz. "The ageing gastrointestinal tract." *Curr Opin Clin Nutr Metab Care* 19 (2016): 12-18.
- Bhutto, Asif and John E. Morley. "The clinical significance of gastrointestinal changes with aging." Curr Opin Clin Nutr Metab Care 11 (2008): 651-660.
- Salles, N. "Basic mechanisms of the aging gastrointestinal tract." Dig Dis 25 (2007): 112-117.
- 4. Morley, John E. "The aging gut: Physiology." Clin Geriatr Med 23 (2007): 757-767.
- Besanko, Laura K., Carly M. Burgstad, Charles Cock and Richard Heddle, et al. "Changes in esophageal and lower esophageal sphincter motility with healthy aging." J Gastrointestin Liver Dis 23 (2014): 243-248.

How to cite this article: Gajewski, Michele. "Understanding the Main Disorders of the Gastrointestinal Tract in Older Adults." *J Diabetic Complications Med* 9 (2024): 248.