

Workplace Violence, Anxiety and Self-Esteem in Nursing Staff of Primary, Emergency and Intensive Care Units on the Island of Crete

Maria Maniou^{1*}, Sofia Zyga², Spyros Vliamos³, Panagiotis Prezerakos⁴, Katerina Flora⁵ and Constantinos Togas⁶

¹Faculty of Nursing, Technological Educational Institute of Crete, Iraklion, Greece

²Faculty of Nursing, University of Peloponnese, Sparta, Greece

³Neapolis University Pafos, Cyprus

⁴Department of Nursing, University of Peloponnese, Sparta, Greece

⁵Department of Clinical Psychology, Neapolis University Pafos, Cyprus

⁶Department of Clinical Psychology, Corinth Prison, Korinth, Greece

Abstract

Introduction: Mobbing in the workplace occurs when someone repeatedly, for a long time is exposed to negative behaviors and may have difficulty in defending himself. In the nursing profession, labor intimidation is common. It is now accepted that anxiety and self-esteem can play a particularly important role in mental and in physical health of nurses.

Purpose: The investigation of self-esteem, the existence of anxiety and the phenomena of workplace mobbing of nursing staff working in Primary Health Care, Intensive Care Unit and in the Emergency Department.

Methodology: An investigation was carried out in the area of Crete between August 2017 and January 2018 in fourteen Health Centers, two Primary National Health Networks, four Emergency Department, eleven Intensive Care Units. The study involved 213 nurses. The Leymann Inventory of Psychological Terror, the State-Trait Anxiety Inventory and the Culture-free Self-esteem Inventories has been used.

Results: The average age of the participants was 41.73. The 45.5% of the sample is working in Intensive Care Units, 24.9% in Emergency Departments and 29.6% in Primary Care Units. The mean value of trait anxiety was 40.82, state anxiety 39.03, and overall anxiety 79.85 for the overall study sample. They were exposed to at least one mobbing behavior in the past 12 months, and for at least once a week, almost daily or daily the 11.3%. These 24 nurses attribute this behavior to competition problems (60.9%) and jealousy (58.7%). The most nurses of the total sample (50.5%) had a middle self-esteem.

Conclusions: The evaluation of the results shows that the nursing staff of the overall sample of the study experiences mild anxiety symptoms. Mobbing seems to be at high rates. Early recognition of the phenomenon and its management as well as enhancing of the self-esteem should be the best practice of intervention to prevent it.

Keywords: Mobbing; Anxiety; Self-esteem; Nursing staff

Introduction

Heinz Leymann was the first who defined workplace bullying that has been a part of working life for centuries and denote a specific form of workplace aggression towards employees [1,2]. According to Leymann [1] the psychologically violent behaviours he saw in workplaces in Sweden were a kind of “workplace terrorism”, and determined it as “a type of psychological terrorism that arises in the form of directed, systematic, unethical communication and competitive behaviour by one or more persons towards one person”.

The following definition of mobbing see widely agreed upon and it is accepted for this study: Bullying at work means; “*harassing, offending, socially excluding someone or negatively affecting someone’s work tasks*”. In order for the label bullying (or mobbing) to be applied to a particular activity, interaction or process it has to occur repeatedly and regularly (e.g. at least once weekly) and over a period of time (e.g. about six months) [3-7].

Studies that took part in United Kingdom [8] and Finland [9] show prevalence rates of around 10%, whereas in the country of Austria reported results ranged from 7.8% to 26% [10]. The lack of a standardized definition and methodology applied to measure workplace bullying contribute probably to the difference in prevalence rates between populations, countries, and organizations [11].

Also, a survey conducted in nursing personnel of the Greek Emergency Departments showed that conflicts between colleagues related to harassment are 24%. A research work for the effect of

mobbing on the professional life of nurses in seven Hospitals of the 6th Health Region in Greece showed that nurses, men and women (71%), were victims of moral harassment during the past year and had psychosomatic symptoms (anxiety 54.3%, headaches 52%, atony 41.5%, denial of work 28%, depression 16.3% [12].

Violence in the workplace is expressed as anger, harassment, coercion, intimidation and insult [13]. It is mainly manifested by impacts such as cooperative difficulties, reduced resistance to rush and psychological reactions etc [14]. It is a chain of anti-social behaviors and an intense form of work-induced anxiety, and through this systematic and long-lasting process (systematic, repetitive, durable and progressive), the worker-victim is led to a psychological and labor extermination [15].

Anxiety was defined by Freud as “*something felt*”, “*an emotional state that included feelings of consciousness, intensity, nervousness and anxiety*”

***Corresponding author:** Maniou Maria, Faculty of Nursing, Technological Educational Institute of Crete, Iraklion, Greece; Tel: +030 2810 379-538; E-mail: mmaniou@hotmail.com

Received July 08, 2018; **Accepted** August 07, 2018; **Published** August 17, 2018

Citation: Maniou M, Zyga S, Vliamos S, Prezerakos P, Flora K, et al. (2018) Workplace Violence, Anxiety and Self-Esteem in Nursing Staff of Primary, Emergency and Intensive Care Units on the Island of Crete. J Nurs Care 7: 464. doi:10.4172/2167-1168.1000464

Copyright: © 2018 Maniou M, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

accompanied by normal stimulation." Parallel to and in conjunction with Darwin's evolutionary theory, Freud observed and noted that stress was adaptive to the motivation of behavior that prompted individuals to deal with threatening situations, and that intense anxiety prevailed in most psychiatric disorders [16]. It has also prevailed, the definition of anxiety as: "A normal physical or psychological reaction to external events that cause a strong emotional state to man. This is a long-standing difficulty or a serious personal event that lasts for at least four (4) weeks" [17].

According to Spielberger [16], transport concern refers to "subjective emotional state characterized by tension and asphyxia". Permanent anxiety refers to "the predisposition of the individual to perceive specific situations as threatening and to react with the anxiety to them".

Spielberger [16] about permanent and temporary anxiety considers that the recruitment of the individual from both internal and external stimuli is considered as threatening, resulting in reactions of temporary anxiety. Sensory and cognitive feedback mechanisms are the ones that cause high levels of transient stress to be recognized as unpleasant by the individual. The reaction of transient anxiety is proportional to the magnitude of the intensity of the threat.

The sensory and cognitive mechanisms of feedback are those that cause high levels of temporary stress to be recognized as unpleasant by humans. The reaction of temporary anxiety is proportional to the magnitude of the intensity of the threat. Correspondingly, the duration of tension affects the stability of the trait stress response. He also notes that people with a high level of permanent anxiety perceive more situations as threatening and react with more intense transient resistance reactions. Anxiety causes people to develop specific psychological protection mechanisms to achieve the reduction of temporary anxiety [18].

Nursing, as a profession, is described as a "high intensity profession", the reasons have already been extensively formulated. This in itself has negative effects on the self-esteem of the nurse, both as a person and as a professional. In Greece, a survey was conducted in three (3) public hospitals in the Prefecture of Ilia during the period 05-10/2008, with a sample of 167 nurses and nursing assistants aged 20-60 years old. The statistical analysis of the questionnaire showed that 56.3% of the sample of nurses has a "modest" emotional state, which affects their self-esteem [19].

There are no surveys in Greece related to the correlation of work harassment with permanent, temporary anxiety and self-esteem in nursing staff. The first aim of this study was to measure the prevalence and forms of workplace bullying among the nursing personnel working in the Primary Health Care, Emergency Departments (ED) and Intensive Care Units (ICU) of Crete. The first approximation comprises the measurement of the frequency and duration of exposure to one or more of the 45 forms of bullying [20] and the second approximation include a self-reporting question relevant to bullying within the past 12 months based on a definition according to Heinz Leymann [6]. The second aim was to measure the existence of state and trait anxiety with two questionnaires [21] and the third aim was to measure the self-esteem of the nursing personnel.

Aim

The aim of this study was to investigate the state and trait anxiety and work violence in nursing staff in Primary Health Care, in Emergency Departments (ED) and in Intensive Care Units (ICU) of five major hospitals and 11 Health Centres in the prefecture of Crete.

Methodology

The study involved 213 nurses from five hospitals in the region of

Crete. Of these, 45.5% were working in Intensive Care Units while the other 24.9% were working at the Emergency Departments and 29.6% in Primary Health Care. The sample selection was made by the method of random sampling. The inclusion criteria were: i) be nursing staff; ii) be nursing staff coming from all ICU, ED and Primary Health Care centres of Crete and the exclusion criteria was not to be a student nurse

The survey was conducted from August 2017 to January 2018 and included the voluntary and anonymous participation of nursing staff. The psychometric tools included in the study are presented below:

Instruments

For the purpose of the present research, the French version of "Leymann's Inventory of Psychological Terror" (LIPT) instrument [22], the Greek Version of "Leymann Inventory of Psychological Terror" Instrument were applied [11], The State-Trait Anxiety Inventory (STAI) [20,21] and the Culture-free Self-esteem Inventories [23].

Leymann's Inventory of Psychological Terror consists of 45 items, each item measuring the exposure to workplace bullying the last 12 months with two response options (yes or no) [11]. In addition, two questions on the frequency were included (monthly basis, weekly, or daily) as well as the duration of bullying (years and months). In five sections are grouped the 45 bullying behaviors (1) social relationships at work (criticism, no possibility to communicate, and indifference and verbal aggression), (2) exclusion (isolation, avoidance and rejection), (3) job tasks (too many tasks, no tasks, uninteresting tasks, humiliating tasks, tasks superior, or inferior to skills), (4) personal attacks (attacks on origins or opinion, rumors, ridicule and gossiping), and (5) physical violence (physical threats and sexual annoyance). According to Leymann, those who report exposure to at least one of the 45 behaviors that concerns bullying the last 12 months, weekly or more, and for six months or longer are defined as victims of bullying.

Moreover, in the present study will be used questions included in the French version of LIPT instrument. The instrument includes the definition of workplace bullying: "Bullying may be defined by a situation in which someone is exposed to a hostile behavior on the part of one or more persons in the work environment that aim continually and repeatedly to offend, oppress, maltreat or to exclude or isolate over a long period of time" [4]. A) Nurses are asked whether they aware themselves as being victims of bullying within the previous year. For those considering themselves as being victims, contributing factors are further researched. B) In addition, nurses are asked whether they witnessed bullying at the current work to another employee during the past 12 months.

The State-Trait Anxiety Inventory (STAI) is a psychological inventory based on a 4-point Likert scale. It consists of 40 questions. The STAI measures two type of anxiety-state anxiety, or anxiety about an event and trait anxiety or anxiety level as personal characteristics. Higher scores are positively correlated with higher levels of anxiety.

State anxiety (S-anxiety) can be defined as discomfort, fear, nervousness, etc. and the arousal of the autonomic nervous system induced by different situations that are apprehend as dangerous and is considered temporary. Trait anxiety (T-anxiety) can be defined as feelings of worry, stress discomfort, etc. that one experiences and how people feel across typical situation daily. The State-Trait Anxiety Inventory assess both state and trait anxiety separately. Each type of anxiety has its own scale of 20 questions and the scores range from 20 to 80, with higher scores correlating with higher levels anxiety. Each scale asks twenty questions each and based on a 4-point Likert scale. Low scores show a mild form of anxiety. Median scores indicate a moderate

form of anxiety and high scores shows a severe form of anxiety. Anxiety absent questions impersonate the absence of anxiety in a statement like, “I feel secure.” Anxiety declares questions represent the presence of anxiety e.g. “I feel worried.” More examples from the STAI on anxiety absent and present questions are below. The 4-point scale for S-anxiety is as follows: (1) not at all, (2) somewhat, (3) moderately so, (4) very much so and for the 4-point scale for T-anxiety is: (1) almost never, (2) sometimes, (3) often, (4) almost always [21].

The Culture-free Self-esteem Inventories is a self-referencing questionnaire, which includes (without the lie scale) 32 statements. These statements seek to measure the general, personal and social perception of themselves (self-perception) and are divided into two groups: those that are high and those showing low self-esteem. Each question answers either with “yes” or with “no”. Three self-assessment sub-scales are included: general (16 statements), social (8 statements) and personal (8 statements). There is also a lie scale o (8 statements) [23].

Permissions were obtained from the developers of the French version of LIPT instrument, the Greek version of LIPT instrument, the State-Trait Anxiety Inventory (STAI) and Culture-free Self-esteem Inventories. The time needed to fill out the questionnaire was 8-10 min.

Study Population

The present study conducted among nurses working in primary health-care setting with the participation of 14 Health-care Centres and 5 Hospitals in the island of Crete. A total of 213 nurses take part in the study. The study was performed in the following departments: 14 Health-care Centers, 4 Emergency Departments (ED) and 11 Intensive Care Units (ICU). The study was performed during August 2017 to January 2018.

Statistical Analysis

With respect to the statistical analysis that follows, the quantitative variables are reported based on the mean ± standard deviation (mean ± SD) as well as the median and the Interquartile Range (IQR), while for the qualitative variables we have the corresponding frequencies and percentages. Depending on the appropriate statistical and/or graphic controls, it is recommended that median and the Interquartile Range (median, IQR) are used as representative descriptive measures.

The Shapiro-Wilk statistical control was used to check the regularity of the quantitative variables in the questionnaire. The existence of regularity has also been confirmed or rejected by the visual overview of the corresponding histograms, the normal Q-Q plots and box-plots of the variables. At the same time, the appropriate parametric and non-parametric statistical examinations were conducted to investigate any differences between the three structures (ICU, KY/PEDY and TEI) and the scales under study, determining the level of significance at 0.05. In all cases it was necessary to use accurate tests and/or model-carousel simulation (10000 samples). Also, where necessary, the Levene test was used for the homogeneity of the difference.

Finally, reliability and internal consistency was assessed by internal consistency using Cronbach’s alpha test. Statistical analysis was performed using the statistical software IBM SPSS statistics (version 21.0). A p value<0.05 was considered statistically significant.

Ethics Approval

The researcher submitted the research protocol for obtaining the necessary written permissions from the Research and Ethics Committees of the University General Hospital of Heraklion, General

Hospital of Heraklion “Venizelio Pananio”, General Hospital of Chania, General Hospital of Agios Nikolaos, General Hospital of Rethymno and the 7th Health District of Crete. The permissions of the above organizations were given provided that the results of the study will be announced in the Administration of the 7th Health District of Crete. After the permissions of the research were given, the participants were approached. Written consent was given by all participants to participate in the study.

Results

Reliability of Leymann’s Inventory of Psychological Terror (LIPT) instrument

The reliability of LIPT instrument expressed by Cronbach α was 0.938 suggesting high internal consistency.

Characteristics of the study sample

In this study, the participants were nursing staff (n=213) and the majority of the participants 89.2% of the sample were women and ICU nurses represented 45.5% of the study population. Mean age for the total sample was 41.73 (SD=7.33 years). The 75.1% (n=160) of the sample were married and followed by 23.0% (n=23%) that were unmarried. The majority of the sample 64.8% (n=138) were graduates of Technological Educational Institute (TEI) also, 8.9% of the sample (n=18) had a master’s degree and one person had a PhD. Demographic characteristics of the research population are shown in Table 1. In regard to the mean length of employment was 15.78 (SD=8.49 years). The average of work in the current department was 8.00 years (SD=10.50 years) as shown in below Table 2.

Characteristics	n (%)
Departments	
	Intensive Care Unit 97 (45.5)
	Primary Care 63 (29.6)
	Emergency Department 53 (24.9)
Gender	
	Man 23 (10.8)
	Women 190 (89.2)
Marital Status	
	Married 160 (75.1)
	Unmarried 49 (23.0)
	Widowed/ Divorced 4 (1.9)
Educational Level	
	Secondary School 43 (20.2)
	Technological Educational Institute 138 (64.8)
	University 13 (6.1)
	Master degree 18 (8.4)
	PhD 1 (0.5)

Table 1: Characteristics of the study population (N=213).

Characteristics	
Age (years)	41.73 ± 7.33*
Years of work	15.78 ± 8.49*
Years of work in the present department	8.00 (10.50)**

* mean ± sd
**median (IQR)

Table 2: Descriptive characteristics among the study population (N=213).

Prevalence of workplace bullying among nurses

According to Leymann, [1] mobbing is defined as the worker's exposure at least once a week for the past 12 months. Based on this definition, we have the following results. Among the 213 nurses that consisted the total sample of the study, 95 nurses (44.6%) were exposed to at least one bullying behavior at work within the last 12 months, whereas 24 nurses (32.43%) were exposed to at least one bullying behavior or more at least once weekly the last 12 months as shown in Table 3. There is a statistically significant difference between the three

	Yes (%)
Industrial Relations	
Being silenced by superior	8.5
Being continuously interrupted	10.3
Being silenced by others	14.6
Being scolded and yelled	11.3
Being criticized regarding work assignments	9.4
Private life being criticized by others	5.2
Being terrorized by means of phone calls	5.2
Receiving verbal threats	7.5
Receiving written threats	2.3
Being exposed to irritating gestures/looks	14.6
Physical presence ignored, addressing only others	9.4
They isolate you systematically	
Not being talked to	5.6
Not being allowed to physically contact others	3.3
Being isolated from others at work	3.3
Conversation with colleagues is forbidden	1.9
Physical presence being ignored among others	5.2
Being addressed only in written ways	0.5
Your professional duties have been amended as a punishment	
They do not assign you jobs, you have no job	0.9
Being given meaningless work assignments	7.5
Being given work assignments far below capacity	6.1
Continuously being given new work assignments	8.5
Being given humiliating work assignments	2.8
Being given difficult work assignments far above capacity	7.5
Attacks on your face	
Being gossiped	15.0
Being exposed to slanders and lies	12.2
Being ridiculed	4.7
Being said to have a mental illness	1.9
Being forced to go through psychiatric exams	1.9
Being mocked due to a handicap that you have	4.2
Voice, gestures, and way of moving are imitated to tease	5.6
Suffering verbal attacks regarding political and religious beliefs	3.3
Being teased due to ethnic background	3.8
Being forced to do work assignments which are against your conscious	8.9
Being judged for your work in an injustice and humiliating way	11.3
Your decisions are questioned by others	8.9
Being reviled using obscene or degrading terms	2.8
Being sexually threaten	2.3
Violence and threats of violence	
Being given dangerous work assignments for your health	4.7
Despite your bad health you are forced to do work assignments that heart your health	7.0
Being physically threaten	0.9
Being physically threatened in the form of mild violence as a warning	2.8
Being physically attacked	1.4
Being forced to spend big sums of money	0.5
Workplace or home is damaged by others	1.9
Being sexually attacked	0.0

Table 3: Prevalence of exposure to each of the 45 mobbing behaviors among the study population during the last 12 months (N=213).

structures (p=0.016) in the percentages of those who said they were exposed to at least one mobbing (out of 45) in the last 12 months. We observe that the highest rate of exposure to at least one moral harassment behavior is reported in the Emergency Department (58.5%), while the lowest percentage in Primary Care (31.7%) as shown in Table 4. With regard to the frequency of the mobbing in the last 12 months and the answers "rarely" or "at least once a month" and "at least once a week" or "almost daily" or "daily", compared to the structures, there is no statistically significant difference, as shown in Table 5. The percentage of the nurses still continuing to experience similar situations is 69.23% (n=54). The median of the time period for which someone with these conditions had come was 12 months (IQR=51.75). The prevalence of the victims of workplace bullying among nurses working at the Intensive Care Units was 10 (41.7%) whereas among those working at the Emergency Departments (ED) was 10(41.7%), and among nurses working at the Primary Health Care was 4 (16.6%).

According to Niedhammer et al. [20] definition 16.9% of the total sample reported exposure in bullying behaviors at their workplace the last 12 months. Between them 16.5%, 9.5% and 26.4% of the nurses worked at the ICU, Primary Health Care and Emergency Department respectively. Finally, based on the definition of psychological violence at work, there was a statistically significant difference between the three departments (p=0.047), with the highest percentage reported by Emergency Departments (26.4%).

			Exposure to at least one moral harassment behavior		Total
			No	Yes	
Departments	ICU	N	53	44	97
		%	54.6%	45.4%	100.0%
	Primary Care	N	43	20	63
		%	68.3%	31.7%	100.0%
	ED	N	22	31	53
		%	41.5%	58.5%	100.0%
Total		N	118	95	213
		%	55.4%	44.6%	100.0%

$\chi^2(2)=8.374$; p=0.016

Table 4: Mobbing in departments among the study population during the last 12 months (N=213).

			How often have you been confronted with one or more of the above situations in the last 12 months?		Total
			Rarely/At least once a month	At least once a week/ Almost daily/ Everyday	
Departments	ICU	n	23	10	33
		%	69.7%	30.3%	100.0%
	Primary Care	n	12	4	16
		%	75.0%	25.0%	100.0%
	ED	n	15	10	25
		%	60.0%	40.0%	100.0%
Total		n	50	24	74
		%	67.6%	32.4%	100.0%

$\chi^2(2)=1.125$; p=0.625

Table 5: Frequency of mobbing in departments.

Moreover colleagues, chiefs or persons holding higher rank in hierarchy were most commonly recognized as mobbers between participants (43.9%, 43.9% respectively). With regard to the sex of the persons who have practiced psychological violence, 43.31% were women, 13.58% were men and 43.31% were both of them (women and men). The median of the number of people who were against those who received psychological violence during this period was 2 (IQR=2).

Bullying behaviors and frequency of exposure to workplace bullying

The most common mobbing behaviors reported among the participants were the following: “Being badly criticized behind your back” (15.0%); with the same percentage appears “being continuously interrupted by others” and “being accepted contemptuous glances and/or contemptuous gestures” (14.6%); “being exposed to slanders and lies” (12.2%); and with the same percentage appears “being vilified and shouted” and “being criticized regarding work assignments” (11.3%).

According to the definition of mobbing as proposed by Niedhammer et al. [20], 29.3% of the study participants answered positively regarding exposure to hostile behaviors within the last 12 months (45.4% of those working at the ICU, 31.7% of those working in Primary Care and 58.5 of those working in Emergency Department respectively). There is a statistically significant result ($p=0.016$). We observe that the highest rate of exposure to at least one behavior of moral harassment is reported in Emergency Department (58.5%), while the lowest rate in Primary Care (31.7%).

When applying the definition of mobbing as proposed by Niedhammer et al. [20], 36 nurses (16.9%) of the study participants answered that having been subjected to psychological violence at work in the last 12 months (16, 0.5% of those working at the ICU, 9.5% of those working at the Primary Health Care and 26.4% of those working at the Emergency Department, respectively). The most common causes responsible for the exposure to hostile behaviors for the total sample were “competitive behaviors between employees” (60.9%) and “jealousy” (58.7%), followed by “a generally bad working environment” (17.0%) and “problems in the management and professional position” (14.2%). The 30.0% of the total sample of the individuals reported that they

became observers of mobbing behaviors against another employee during the previous year.

Reliability of the State-Trait Anxiety Inventory (STAI)

The reliability of STAI instrument expressed by Cronbach α was 0.940 suggesting high internal consistency (State anxiety: $\alpha=0.895$, Trait anxiety: $\alpha=0.907$).

Prevalence of state-trait anxiety

The mean value of trait anxiety was 40.82, the mean value of state anxiety was 39.03 and the total anxiety was 79.85 (Table 6).

Concerning the first subscale for the emotional state (trait anxiety) of the interviewee at the time of completing the questionnaire we have the following results. Based on ANOVA, there is no statistically significant difference in the mean score of Spielberger [1], between the three structures: F (2,210)=1.991, $p=0.139$, Intensive Care Units (ICU): 41.74 ± 11.59 , Primary Care: 35.57 ± 9.62 , Emergency Department (ED): 41.81 ± 10.04 . Based on the Tukey HSD test for comparisons between two structures, there was no statistically significant difference between ICU-Primary Care ($p=0.160$), nor between ICU/ED ($p=0.236$) nor between Primary Care-ED ($p=0.236$).

Regarding the second sub-scale for the moral status (state anxiety) of the persons how they feels general in their lives, based on ANOVA, there is no statistically significant difference in the mean score of this unit of the three structures: F (2,210)=2.752, $p=0.066$, ICU: 40.28 ± 10.58 , Primary Care: 36.59 ± 8.50 , ED: 39.66 ± 10.44 . Based on the Tukey HSD for comparisons between two structures, there was no statistically significant difference between the ICU-Primary Care ($p=0.060$), neither between the ICU-ED ($p=0.930$) nor between the Primary Care-ED ($p=0.226$).

Regarding Spielberger’s overall scale and overall score, we have the following results. First of all, the homogeneity of the dispersion did not apply, so we proceeded to the Welch test, which resulted in a statistically significant difference between the three groups, $p=0.039$, ICU: 82.02 ± 21.16 , Primary Care: 75.16 ± 15.53 , 19.22. However, based on the two-time comparisons and on the basis of the Tamhane and Dunnett T3

Scales	n (%)	Mean	SD	Median	IQR	Range
State-Trait Anxiety Inventory (STAI)						
STAI-Part 1		40.82	10.72	39.00	16.00	23-68
STAI-Part 2		39.03	10.06	38.00	13.50	22-70
STAI-(Part 1 & Part 2)		79.85	19.32	77.00	25.00	47-138
Culture-free Self-esteem Inventories (James Battle),						
General self-esteem*		12.27	2.89	13.00	3.50	2-16
Social self-esteem*		6.35	1.38	7.00	1.00	1-8
Personal self-esteem*		3.86	2.16	4.00	3.50	0-8
Scale of lie		4.77	1.56	5.00	2.00	0-8
Total self-esteem		22.47	5.28	23.00	7.00	4-32
Too low	17 (8.0)					
Low	37 (17.4)					
Middle	110 (51.6)					
Hugh	32 (15.0)					
Very Hugh	17 (8.0)					

*(IQRs) as representative descriptive measures for these scales

Table 6: Descriptive characteristics of the scales among the study population (N=213).

controls, there was no statistically significant result between the ICU-Primary Care (p=0.057), nor between the ICU-ED (p=0.998) between Primary Care-ED (p=0.162).

Reliability of Culture-free Self-esteem Inventories (James Battle)

The reliability of Culture-free Self-esteem Inventories (James Battle) expressed by Cronbach α was 0.763 suggesting high internal consistency (General self-esteem: $\alpha=0.737$, Social self-esteem: $\alpha=0.442$, Personal self-esteem: $\alpha=0.724$, Lie scale: $\alpha=0.525$, General self-esteem: $\alpha=0.763$).

Valuation of self-esteem

From the analysis of the results of the Culture-free Self-esteem Inventories (James Battle), it was found that 51.6% of the nurses who participated in the study had a middle self-esteem, while the lowest percentage of the total sample of 8.0% showed very high self-esteem. Also, it was found that 8.0% had very low self-esteem, 17.4% had low self-esteem and 15% had high self-esteem. Also, the results showed that the General self-esteem was 12.27 (mean), the Social self-esteem 6.35 (mean), the Personal self-esteem was 3.86 (mean) and the scale of lie was 4.77 Table 6. Regarding the Culture-free Self-esteem Inventories and based on the chi-square test was applied. There are no statistically significant differences between the three structures (ICU, ED, and Primary Care) compared to the level of self-esteem, despite the fairly large difference in many percentages. Something that is probably due to the existence of many categories at the level of self-esteem. With regard to the level of self-esteem for the three structures (ICU, ED, Primary Care) we had the following results: ICU: very low self-esteem: (8.2%), low self-esteem (17.5%), middle self-esteem (50.5%), high self-esteem (13.4%), very high self-esteem (10.3%), ED: very low self-esteem: (11.3%), low self-esteem (26.4%), middle self-esteem (37.7%), high self-esteem (20.8%), very high self-esteem (3.8%), Primary Care: very low self-esteem: (4.8%), low self-esteem (9.5%), middle self-esteem (65.1%), high self-esteem (12.7%), very high self-esteem (7.9%) (Table 7).

Discussion

The present study investigated the anxiety levels, work violence and self-esteem in a sample of 213 nursing staff members in five major hospitals and eleven Health Centres in the prefecture of Crete.

In the present study, they have been exposed to at least one mobbing behavior in the last 12 months and at least once a week, almost daily or daily the 11.3% of the total sample (n=24), and these people, according to Leymann [1] are the victims of mobbing. In a relevant study carried out in Cyprus in 2014 and the study sample was made up of 136 health professionals (44 general practitioners, 50 nurses, 42 other health professionals) working in Primary Health Care in Nicosia

and was used the same questionnaire of the Leymann Inventory of Psychological Terrorization (LIPT) as used in the present study, the prevalence of “*mobbing syndrome*” among health professionals according to Leymann’s official definition was 8.8% of the total sample reported exposure at least one behavioral psychological violence in the last 12 months, while 11% reported exposure to at least one “*mobbing behaviour*” at least once a week. In another study when the Leymann definition was applied, 5.9% of the participants reported exposure to at least one bullying behavior at least once a week and for at least 6 months [11].

The analysis of the data from the present study showed that the most common “*mobbing*” behaviors reported among the participants were: “*they speak badly for you behind your back*” and the same percentage follows “*they interrupt you while you express*” 15.0%), “*receive scornful eyes and/or scornful gestures*” (14.6%), “*spread false rumors about you*”, and with the same percentage they appear to “*revile you and shout*” (12.2%) and finally, they “*they are constantly criticizing your work*” (11.3%).

In the present study, the less frequent “*mobbing behaviors*” identified among nursing staff were “*sexual assault*” (0.0%), “*they are only addressed to you in writing*” (0.5%), and “*they are causing you damage by trying to hurts you financially* (0.5%)”.

On the other hand, the less frequent “*mobbing behaviors*” identified among health professionals in Primary Health Care was: “*you’ve been sexually attacked*,” “*psychological violence*,” “*physical violence*,” “*oral or sexual suggestions*,” “*want to force you to undergo a psychiatric examination*” and “*accept written threats*” (0% for all of the statements above) [24].

The 63.5% of the nurses who participated in the survey stated that they had received psychological violence by a “*colleague*” (this is why the total figure exceeds 100%), which accounts for 43.9% of all the answers and with the same percentage of “*people with a higher hierarchical position or superior*”. Also, with regard to the gender that commit psychological violence, the 43.2% of the nurses answered that were “*women*”, with the same percentage being “*both men and women*” and the median price of the number of people who commit psychological violence during this period was 2 (IQR=2). According to Zachariadou et al. [11] states that senior officials were identified as those who practiced psychological violence (57.5%). Health professionals working in public hospitals and Primary Health Care reported having faced hostile attitudes mainly by their superiors (58.3% vs. 55.9%, respectively). As for the sex of perpetrators, women were most often identified as those who had commit psychological violence compared to men, but there was no statistical significance (69.1% vs. 19.8% p=0.147) [25]. Another research in Turkey by Picakciefe et al. [25] to people working in the Primary Care, 70.3% of the victims was superiors. In the present study

			Level of self-esteem					Total
			Very Low	Low	Middle	High	Very high	
Department	ICU	N	8	17	49	13	10	97
		%	8.2%	17.5%	50.5%	13.4%	10.3%	100.0%
	Primary Care	N	3	6	41	8	5	63
		%	4.8%	9.5%	65.1%	12.7%	7.9%	100.0%
	ED	N	6	14	20	11	2	53
		%	11.3%	26.4%	37.7%	20.8%	3.8%	100.0%
Total		N	17	37	110	32	17	213
		%	8.0%	17.4%	51.6%	15.0%	8.0%	100.0%

$\chi^2(8)=13.898$; p=0.080

Table 7: Level of self-esteem among the study population (N=213).

the participants stated that they discussed the problems they faced in their workplace related with violence and 68.3% of the respondents said that among other things they had spoken with their colleague. Also, only 2 of respondents said they did not have a person they could contact but they would like to have, while 3 people said they did not have a person to contact or they did not need. In a study of Zachariadou et al. [11] it is reported that health professionals working at major public hospitals in Cyprus have discussed their exposure to mobbing behavior at their workplace with a colleague of 67.7%, while the 30.6% with a member of his family.

By applying the definition “Psychological violence at work can be defined as the situation in which an individual is subjected to hostile attitudes by one or more individuals in his working environment which in a continuous and repeated manner seek to destroy him, to oppress him, to abuse him or even to exclude or isolate them for a long time”. 36 nurses (16.9%) said they have been subjected to psychological violence in their work in the last 12 months.

This behavior was attributed to “competition problems between individuals” (60.9%) and “jealousy” (58.7%). According to the Chira et al. [24] study, 30 workers (26.3%) stated that they had been subjected to psychological violence in the last year according to this definition and this behavior was attributed to “poor organization of work” (43.3%) followed by “problems in administration” (40%). According to Zachariadou et al. [11], 29.3% of respondents responded positively to exposure to hostile behavior over the last 12 months according to the definition (31.4% of those working in the large public hospitals in Cyprus and 26.3% of those working in Primary Health Care respectively). The most common causes of exposure to hostile attitudes towards health professionals working at major public hospitals in Cyprus were “management/ placement problems at work 33.3%” followed by “generally poor working atmosphere 31.3%”, “poor organization 29.2%” and “jealousy 29.2%”. In Primary Health Care, the most common causes were “poor organization 43.3%”, followed by “problems in the administration/ placement at work 40%” and “poor working atmosphere 33.3%” and with the same percentage “competition problems among individuals”.

To the question: “You have perceived psychological violence against someone else in your workplace in the last 12 months,” 30.0% of nursing staff in the sample of this survey reported that they have perceived such a situation. A similar study reported 38 workers (31.7%) who responded positively to the question. In another survey, respondents reported that they had notice (43.4%) bullying behavior against another employee in the previous year. Health professionals working in hospitals in Cyprus noticed that one or more of their colleagues were harassed more frequently than primary care workers in Cyprus (52.2% vs. 31.4%), which was statistically significant ($p=0.001$). In England, Quine [26], a health care practitioner, it was found that 42% of the participants had witnessed intimidation behaviors against their colleagues.

From the analysis of the results of the STAI questionnaire it was found that the mean value of the trait anxiety was 40.82 while the mean value of the state anxiety was 39.03 and the mean value of the total anxiety was 79.85. A study carried out in 2010 in nursing personnel in the sixth Healthcare Region of Greece working in nephrologic centers showed that trait anxiety was 41.4 and state anxiety 38.6 [27]. Finally, in a research that investigated the trait and state anxiety in nursing personnel working at Thessaly Health Centres, they showed high rates of trait anxiety, which exceeded the corresponding rates for the Greek population [28].

In the present study, the results of the Battle Questionnaire for Self-Esteem showed that 51.6% of the nursing staff involved in the study had

a middle self-esteem, while the lowest percentage of the total sample (8.0%) had a very high self-esteem. Researchers conducted a study in Wales on mental health community nurses who found that most nurses had middle self-esteem [29]. According to Randle [30], the result of continuous exposure to bullying includes the reduced of self-esteem .

Conclusions

The evaluation of the results shows that the nursing staff of the overall sample of the study experiences mild anxiety symptoms, which necessitate the training of nursing staff in symptom management to avoid increasing their intensity. Mobbing seems to be at high rates. Prevalence of mobbing is an existing reality in the health sector of Crete. Early recognition of the phenomenon and its management as well as enhancing of the self-esteem should be the best practice of intervention to prevent it.

References

1. Leymann H (1990) Mobbing and psychological terror at workplaces. *Violence Vict* 5: 119-126.
2. Davenport N, Schwartz RD, Elliott GP (1999) *Mobbing: Emotional abuse in the American Workplace*. Civil Society Publishing
3. Einarsen, S, Helge H, Dieter Z, Carly C (2003) *Bullying and emotional abuse in the workplace: International Perspectives in Research and Practice*, CRC Press.
4. Einarsen S (2000) Harassment and bullying at work; A review of the Scandinavian approach. *Aggression and Violent Behavior* 5: 379-401.
5. Einarsen S, Skogstad A (1996) Bullying at work: Epidemiological findings in public and private organizations. *Eur J Work Organ Psychol* 5: 185-201.
6. Leymann H (1996) The content and development of mobbing at work. *Eur J Work Org Psychol* 2: 165e84.
7. Zapf D (1999) Organisational, work group related and personal causes of mobbing/bullying at work. *Int J Manpower* 20: 70-85.
8. Hoel H, Cooper C (2000) *Destructive conflict and bullying at work*. Manchester School of Management. University of Manchester Institute of Science and Technology, Manchester.
9. Vartia M. (1996) The sources of bullying: psychological work environment and organizational climate. *Eur J Work Organ Psychol*, 5: 203-214.
10. Niedl K (1996) Mobbing and well-being: economic and personnel development implications. *Eur J Work Organ Psychol* 5: 239-249.
11. Zachariadou T, Zannetos S, Chira S, Gregoriou S, Pavlakis A (2017) Prevalence and forms of workplace bullying among health-care professionals in Cyprus: Greek Version of “Leymann inventory of psychological terror” instrument. *Saf HealthWork*: 1-8.
12. Siousioura D (2010) *The Effect of group psycho-educational intervention on patients with Type I diabetes dissertation*. Rhodes: University of the Aegean.
13. Papadopoulou D (2009) Violence in the workplace of nursing staff. *Nursing* 48: 379-387.
14. Koutrouvides P (2013) Ministry of education and religious affairs. General secretariat for lifelong learning. Time management. Educational material for Lifelong Learning Centers.
15. Tsiama M.H (2013) *The phenomenon of moral/psychological harassment in the workplace: Conceptual determinations*. National Labor and Human Resources Foundation.
16. Spielberger CD (2010) *State-trait anxiety inventory*.
17. Siousioura D (2010) *The effect of group psycho-educational intervention on patients with Type I diabetes mellitus*. Rhodes: University of the Aegean.
18. Mitrousi, S (2014) *Effect of anxiety on the physical and mental health of nurses*. Doctoral Thesis. Sparta.
19. Staurianopoulou T, Stamati S, Gevrekli E, Gourvelou O, Papadimitriou M (2011) Level of illness, fatigue, satisfaction and self-esteem of nursing staff in public hospitals in Ilia. *Rostrum of Asclepius* 10.

20. Niedhammer I, David S, Degioanni S, Drummond A, Philip P (2009) Workplace bullying and sleep disturbances: findings from a large scale cross-sectional survey in the French working population. *Sleep* 32: 1211-1219.
21. Spielberger CD, Sydeman SJ (1994) State-Trait anxiety inventory and state-trait anger expression inventory. The use of psychological testing for treatment planning and outcome assessment. Hillsdale, Lawrence Erlbaum Associates: 292-321.
22. Niedhammer I, David S, Degioanni S, du travail M (2006) The french version of the Leymann's questionnaire on work place bullying: the Leymann inventory of Psychological Terror (LIPT). *J Epidemiol Public Health* 54: 245-262.
23. Stalikas A, Triliva S, Rousi, P (2002) Psychometric instruments in Greece. *Greek Letters Athens*: 240-241.
24. Chira S (2014) Prevalence of mobbing syndrome between healthcare professionals working at the primary healthcare setting of Cyprus. Postgraduate thesis. Open University of Cyprus.
25. Picakciefe M, Acar G, Colak Z, Kilic I (2012) The Relationship between socio-demographic characteristics, work conditions and level "mobbing" of health workers in primary health care. *J Interpers Violence*: 18-23.
26. Quine L (1999) Workplace bullying in NHS community trust: staff questionnaire survey. *BMJ* 318: 228-232.
27. Marneras Ch, Theodorakopouloy G, Albani E, Gouva M, Dimopoulou E, et al. (2010) Satisfaction from work and anxiety level in nurses working in kidney centers. *Nosileftiki* 49: 83-90.
28. Faki M, Gouva M, Papathanasiou I, Prapas P, Pasxou D, et al. (2009) Resistance to workers in primary health care structures. *Interscientific Health Care* 1: 9-15.
29. Fothergill A, Edwards D, Hannigan B, Burnard P, Coyle D (2000) Self-esteem in community mental health nurses: findings from the all Wales stress study. *J Psychiatr Ment Health Nurs* 7: 315-321.
30. Randle J (2003) Bullying in the nursing profession. *J Adv Nurs* 43: 395-401.