

# Heavy Metals Toxicity and the Environment Protection

Om Prakash Meena\*

Department of Chemistry, MSJ Govt. PG. College, Bharatpur, Rajasthan, India

## Abstract

Pollution of environment is one of the most horrible ecological crisis to which we are subjected today. One of the main sources of pollution in the environments is metallic compounds. Metals and metalloids have long been mined and used in numerous applications. This has led to a significant increase of metal pollutions. Metals can accumulate in all environmental matrices at either high or trace levels of concentration. Heavy metals are naturally occurring elements that have a high atomic weight and a density. Therefore amount of various kinds of metals are present in soil, plants, air, lakes, animals, oceanic regions, even in foodstuffs and human beings. Their widespread distribution, especially heavy metals, became serious problems because of their toxicities for animals, human health and the environment. Their toxicity of heavy metals depends on several factors including the dose, route of exposure and chemical species, as well as the age, gender, genetics and nutritional status of exposed individuals. Because of their high degree of toxicity, lead, cadmium, chromium, zinc, nickel, arsenic and mercury rank among the priority metals that are of public health significance. Metals generally enter in the ecosystem in a relatively non-toxic form and generally become intrinsic components of the environment in such a way that it is difficult to remove them from the environment. Some of them are converted into toxic forms through the environmental reactions involving various micro-organisms and non-biological pathways. For example, methylated compounds like dimethyl mercury,  $(\text{CH}_3)_2\text{Hg}$ , are more toxic than their inorganic forms. In the present investigation more attention has been given to heavy metals like lead, cadmium, nickel and zinc. Although, the term "heavy metals" refer to any metallic element that has a relatively high density and is toxic or poisonous at low concentrations. Examples of heavy metals include Pb, Cd, Hg, As, Cr and Ti etc. This review provides an analysis of their environmental occurrence, production and use, potential for human exposure and molecular mechanisms of toxicity, genotoxicity and carcinogenicity.

**Keywords:** Heavy Metals • Environmental • Human Health • Toxicity • Carcinogenicity

## Introduction

### Environmental occurrence, industrial production and use

Heavy metals are defined as metallic elements that have a relatively high density compared to water. According to United Nation Development Programme (UNDP), human development report there was about 185.4 million people during 1988-93, who were not having access of safe drinking water. According to a survey published in "The Hindu" (Nov. 23, 2002), at present about 15,000 habitations in the country have no potable drinking water, while two lakhs are partially covered and 2-17 lakh villages are identified with problem with quality of water. Lakes like Dal and Nagin in Kashmir, Loktak in Manipur and Hussain Sagar in Hyderabad have seriously choked by aquatic weeds due to eutrophication. Sukhtal and Sariatal represent high pollution level in Kumaun region. In urban sector also, the drinking piped water is not safe. Many experiments found industrial pollutants [1-4], e.g. lead, mercury, chemical wastes and other toxic substance in piped water. Arsenic species are determined in water, soil and plants [5]. Mercury pollution was found in the Tapajos River basin Amazon. Aluminium, Pb, Cd has been determined in water samples [6].

## Literature Review

The heavy metals like lead, mercury, synthetic chemicals and some hazardous wastes dissolve in water in a manner that it is quite difficult to detect and separate them in the purification process. Thus problem of deteriorating

*\*Address for Correspondence:* Meena Om Prakash, Assistant Professor, Department of Chemistry, MSJ Govt. PG. College, Bharatpur, Rajasthan, India, E-mail: [omprakashmeena50278@gmail.com](mailto:omprakashmeena50278@gmail.com)

**Copyright:** © 2021 Prakash OM. This is an open-access article distributed under the terms of the creative commons attribution license which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

**Received** 06 July, 2021; **Accepted** 23 August, 2021; **Published** 30 August, 2021

water quality and thus aquatic ecosystem is worst where rivers pass through large cities or areas having industrial establishment which discharge effluents, solid wastes, particles and other hazardous wastes, which account for one-third of the total water pollution. In rural and semi-urban sectors having less number of industrial establishments, water pollution is mostly due to human wastes and agricultural run-off. Utilization of contaminated water causes many water borne diseases. The tendency of heavy metals to concentrate in the sediments may result in a persistent source of the contamination to various tropic levels in the aquatic environment.

### Lead

Lead is a relatively abundant metal in nature, occurring in lead minerals. In the atmosphere, it is relatively more abundant than other heavy metals. The major source of air borne Pb is the combustion of leaded petrol/gasoline. Pb is added in the form of tetra alkyl lead, primarily  $\text{Pb}(\text{CH}_3)_4$  and  $\text{Pb}(\text{C}_2\text{H}_5)_4$ , together with 1,2-Dibromoethane. The major biochemical effect of Pb is its interference with Heme synthesis, which leads to hematological damage. Pb inhibits several of the key enzymes involved in the overall process of Heme synthesis whereby the metabolic intermediates accumulate. One such intermediate is delta-amino levulinic acid. An important phase of heme synthesis is the conversion of delta-aminolevulinic acid to porphobilinogen. The Pb inhibits the porphobilinogen synthase enzyme so that it cannot proceed further to form porphobilinogen. The overall effect is the disruption of the synthesis of hemoglobin as well as other respiratory pigments, such as cytochromes; which require heme. Pb doesn't permit utilization of  $\text{O}_2$  and glucose for life sustaining energy production. At higher levels of Pb in the blood (more than 0.8 ppm) there will be symptoms of anemia due to deficiency of hemoglobin. Elevated Pb levels (more than 0.5-0.8 ppm) in the blood cause kidney dysfunction and brain damage, CNS-syndrome that may result to coma and death. Lead poisoning also causes constipation, abdominal pain and abnormalities in fertility and pregnancy.

The study of effect of lead intoxication on peripheral blood and hematopoietic bone marrow has been done which shows that the percent of reticulocyte in the peripheral blood was 10 times more elevated than the controls. It delays the maturation of erythrocyte. Lead poisoning can be cured by treatment with chelating agents which strongly bind  $\text{Pb}^{2+}$ . Thus calcium chelate in solution is fed to the victim of lead poisoning;  $\text{Pb}^{2+}$  displaces  $\text{Ca}^{2+}$  from

the chelate and the resulting  $Pb^{2+}$  chelate is rapidly excreted in the urine. Long term exposure to lead has been shown to produce behavioral disturbances in human and animal models which are associated with the alternations in cholinergic and dopaminergic neurotransmission in the central nervous system [7]. Lead poisoning is also observed in wild waterfowl in Japan [8] and wood peckers in Sweden [9].

## Cadmium

Cadmium occurs in low concentration in all natural materials. It is a by-product of zinc and lead smelting. Cd mainly finds use in cathode material of alkaline batteries, electroplating and galvanizing stabilizer PVC plastics, silver solders and as colour pigment in paint. Some of the important sources of exposure of Cd are air pollution, cigarette smoke, fruits, grains and vegetables grown in Cd-laden soil, meat (kidney, liver and poultry), fresh water fish, fungicides, incinerators, mining, Ni-Cd batteries, oxide dust, paint, phosphate fertilizers, power plants, sea foods, smelting plants, tobacco etc [10]. Cereals and vegetables account for about 50% of the Cd intake, which are normally about 2-25  $\mu\text{g}/\text{day}$  in children and 10-50  $\mu\text{g}/\text{day}$  in adults. Sea food like mollusks and crustaceans also contribute to daily Cd intake. Hyper accumulation of cadmium by hairy roots of *Thalpi caeruleus* and its content in herbal raw material is determined [11]. It is also determined in Chinese ceramic by using Oscillopolarography. Long term exposure to Cd from industrial or other sources may cause chronic poisoning and thus result in disorders of various systems, e.g. renal diseases, bone diseases, pulmonary damage, anemia, hypertension, yellow discoloration of the front teeth near gum line and possibly cancer (prostate, renal and lung). Even low levels of environmental cadmium exposure may cause bone demineralization leading to increased bone fragility and increased risk of fractures [12].

## Nickel

Nickel is an essential trace metal for animal nutrition. The nutritional essentiality of Ni was established when it was found that rats fed on diet deficient in nickel had retarded body growth associated with reduction of blood hemoglobin concentration, hematocyte and erythrocytes count. However, the ingestion of Ni above the tolerable limits over long period of time causes harmful health effects. The various sources of exposure of Ni are: - Ceramics, cocoa, hair dye, cooking utensils, cosmetics, coins, dental materials, food (chocolates, hydrogenated oils, nuts, food grown near industrial areas), hair spray, industrial waste, jewellery, medicinal plants, metal refineries, metal tools, nickel-cadmium batteries, shampoo, soil waste incinerators, stainless steel kitchen utensils, tap water, tobacco, water pipes [13] etc. Copper and nickel has been determined in the crude oil simultaneously with single sweep polarography [14].

The dietary intake of nickel is highly variable and depends on the Ni content of foodstuffs from various regions. In general food of animal origin has lower Ni content compared to that of plant origin. Bread, cereals, beverages contribute to half of the daily intake of Ni. Based on certain human experiments where skin reaction to Hg was observed after oral dose of 600  $\mu\text{g}$  [15] nickel (as nickel sulphate), it appears that the safe threshold level for Ni Toxicity may be much lower than 600  $\mu\text{g}$ . Ecotoxicity of nickel to *Epsenia fetida*, *Enchytraeus*, *Folsomia candida* is observed [16,17].

## Zinc

The total content of zinc in an adult body is about 2.0 g. Prostate gland is very rich in Zn (100  $\mu\text{g}/\text{g}$ ). Zinc is mainly an intracellular element.

## Biochemical functions

- Zn is an essential component of several enzymes e.g. carbonic anhydrase, alcoholic dehydrogenase, alkaline phosphatase, carboxypeptidase and superoxide dismutase.
- The storage and secretion of insulin from the beta-cells of pancreas requires Zn.
- Zn is necessary to maintain the normal levels of vitamin A in serum.
- It is required for wound healing.

- Gusten, a zinc containing protein of the saliva, is important for taste sensation.

Besides all above heavy metals, arsenic is posing harmful effects on human health in present time [18]. A review of as exposure through the ages shows that metals has poisoned and killed more people than any other toxin known to human kind [19]. Humans may come into contact with and may be exposed to chemicals in soils via a number of pathways, including inhalation, dermal absorption and ingestion. In most cases, the significant pathway in regard to exposure to metal-contaminated soils is ingestion. Incidental soil ingestion by children is an important pathway in assessing public health risks associated with exposure to arsenic contaminated soils. Hair arsenic levels are the useful indicators of chronic arsenic poisoning [20]. Similarly excessive intake of selenium can cause selenosis which includes weight loss, emotional disturbance, diarrhoea and hair loss and garlic odour in breath. In recent years Rajasthan has taken big strides in agricultural production. This has become possible mainly because of irrigation facilities and use of pesticides and fertilizers.

Large number of analytical techniques such as X-ray fluorescence [21], Atomic absorption spectrophotometry [22-27], Flame photometry [28], Chronopotentiometry [29-31], Neutron activation analysis [32-34], are available today for trace analysis purposes but factors such as high cost of instrumentation, much time consumption, lower sensitivity and lesser reproducibility reduce the efficacy of these techniques in trace analysis. Voltammetric methods have proved to be one of the most sensitive and reproducible methods available today for carrying out trace analysis in diverse matrices like food [35-40], drugs [41-53], plants [54,55], soil [56] water [57-62], etc.

## Discussion and Conclusion

In the last three decades, classical d.c. Polarography which was the best analytical technique has been modified to give birth to a number of highly sensitive and reproducible techniques like Normal pulse Polarography, Differential pulse Polarography, Linear sweep voltammetry, Square wave Polarography, Stripping voltammetry, Cyclic voltammetry etc. These techniques are among the best analytical techniques available today for any type of analysis at microgram to nanogram levels. The techniques are so versatile that 80% elements of the periodic table can be analyzed using them. Moreover, instrumentation for these techniques is very simple. This review provides an analysis of their environmental occurrence, production and use, potential for human exposure and molecular mechanisms of toxicity, genotoxicity and carcinogenicity.

## References

1. Dhanalakshmi Krishnamoorthy, Bhagwati Prasad, Neigapula Venkata Swamy Naidu and Suresh Kumar, et al. "Voltammetry of Potassium Octyl Xanthate at Mercury Electrode." *Anal Lett* 41 (2008): 437-443.
2. Passow Ho, Arnold Rothstein and Thomas W Clarkson. "The general pharmacology of the heavy metals." *Pharmacol Revs* 13 (1961).
3. Kirkpatrick L Robin, Di R Giulio and Scanlon F Peter. "Presented at USA-Czechoslovak Seminar on the chronic effects of toxic substances of wildlife, Strbske pleso, Czecho-Slovakia." 1983.
4. Nieboer Evert and David HS Richardson. "The replacement of the nondescript term 'heavy metals' by a biologically and chemically significant classification of metal ions." *Environ Pollut* 1 (1980): 3-26.
5. Mattusch Jurgen, Rainer Wennrich, AC Schmidt and Werner Reisser, et al. "Determination of arsenic species in water, soils and plants." *Fresenius J Anal Chem* 366 (2000): 200-203.
6. Bhakuni DS and DS Rawat. "Bioactive Marine Peptides." *Bioact Mar Nat Prod* (2005): 278-328.
7. Nehru Bimla and Parminder Sidhu. "Behavior and neurotoxic

- consequences of lead on rat brain followed by recovery." *Biol Trace Elem Res* 84 (2001): 113-121.
8. Ochiai Kazuhiko, Tatsuaki Kimura, Kota Uematsu and Takashi Umemura, et al. "Lead poisoning in wild waterfowl in Japan." *J Wildl Dis* 35 (1999): 766-769.
  9. Mörner Torsten and Lars Petersson. "Lead poisoning in woodpeckers in Sweden." *J Wildl Dis* 35 (1999): 763-765.
  10. World Health Organization. "Trace elements in human nutrition and health". *World Health Organization* 1996.
  11. Gnusowski Bogusław. "Zawartosc metali ciezkich w surowcach zielarskich." *Plant Health Prog* 38 (1998): 610-613.
  12. Staessen Jan, Harry A Roels, Dmitri Emelianov and Tatyana Kuznetsova, et al. "Environmental exposure to cadmium, forearm bone density and risk of fractures: A prospective population study." *Lancet* 353 (1999): 1140-1144.
  13. George A Smart and Jack C Sherlock. "Nickel in foods and the diet." *Food Addit Contam* 4 (1987): 61-71.
  14. Gao Manglai, Zheng Gu and Zhongxin Luo. "One-layer-only molecular deposition filming flooding: A review." *Colloids Surf A Physicochem Eng Asp* 572 (2019): 182-196.
  15. William F Sunderman, Anterior Aitio and Lindsay G Morgan, et al. "Biological monitoring of nickel." *Toxicol Ind Health* 2 (1986): 17-78.
  16. Lock Koen and Colin R Janssen. "Ecotoxicity of nickel to *Eisenia fetida*, *Enchytraeus albidus* and *Folsomia candida*." *Chemosphere* 46 (2002): 197-200.
  17. Lock Koen and Colin R Janssen. "Comparative toxicity of a zinc salt, zinc powder and zinc oxide to *Eisenia fetida*, *Enchytraeus albidus* and *Folsomia candida*." *Chemosphere* 53 (2003): 851-856.
  18. Ross H Anderson, Knobeloch Lynda and Warzeche Charlie. "Arsenic Exposure Health Eff." *In Proc Int Conf* (1998): 367-372.
  19. Nriagu Jerome. "Arsenic poisoning through the ages." *Environ Chem Arsenic* 1 (2002): 1-26.
  20. Hindmarsh J Thomas, Diane Dekerkhove, Geoffrey Grime and Jonathan Powell, et al. "Hair arsenic as an index of toxicity." *Arsenic Expos Health Eff* (1999): 41-49.
  21. Environmental US. "What is a pesticide." *EPA Gov* (2007): 15.
  22. Gilden C Robyn, Katie Huffling and Barbara Sattler. "Pesticides and health risks." *J Obstet Gynecol Neonatal Nurs* 39 (2010): 103-110.
  23. Koçyiğit Hasan and Firdevs Sinanoğlu. "Method validation for the analysis of pesticide residue in aqueous environment." *Environ Monit Assess* 192 (2020): 1-13.
  24. Rajesh Sharma, Rajesh Bahadur Thapa, Hira K Manandhar and Sachin Mulmi Shrestha, et al. "Use of pesticides in Nepal and impacts on human health and environment." *J Agric Environ* 13 (2012): 67-74.
  25. Vinod Kumar and Piyush Kumar. "Pesticides in agriculture and environment: Impacts on human health." *J Environ Sci Health* 1 (2019): 76.
  26. Kamrin A Michael. *Pesticide profiles: toxicity, environmental impact and fate*. CRC Press, 1997.
  27. Alert Eurek. "New 'green'pesticides are first to exploit plant defenses in battle of the fungi." *The benefits of pesticides: A story worth telling. Purdue. Edu* (2009): 25-28.
  28. Sangita Dhara and Nand Lal Mishra. "Elemental characterization of nuclear materials using total reflection X-ray fluorescence spectrometry." *Trends Analyt Chem* 116 (2019): 31-43.
  29. Mannweiler Ulrich. "Accuracy and Precision of the Spectrometric Analysis of Al and Its Alloys." *Erzmetall* 28 (1975): 386-390.
  30. Maghrabi A Ibrahim. "Determination of some mineral and heavy metals in Saudi Arabia popular herbal drugs using modern techniques." *Afr J Pharm Pharmacol* 8 (2014): 893-898.
  31. Khemnani Stuti, Aswani B Nair, Ashish Arora and Raj Singh Sindal, et al. "Detection of heavy metal contents in the seed oil of *Solanum malongena* (eggplant) of arid zone." *Int J Bas Appl Chem Sci* 2 (2012): 59-65.
  32. Adepoju-Bello, Omar Abu Issa, Oluwafemi Omoniyi Oguntibeju and GA Ayoola, et al. "Analysis of some selected toxic metals in registered herbal products manufactured in Nigeria." *Afr J Biotechnol* 11 (2012): 6918-6922.
  33. Sarojanni S Mohammed, Batu and Mohammed Bappah Mohammed. "Analysis of Cr in dumpsite soil samples using AAS and EDXRF techniques." *Res J Chem Sci* 2 (2012): 46-51.
  34. Peter J Loveland and Peter Digby. "The extraction of Fe and Al by 0.1 M pyrophosphate solutions: A comparison of some techniques." *J Soil Sci* 35 (1984): 243-250.
  35. Bakanov Vyacheslav Ivanovich and Matvei Safonovich Zakharov. "Inversion amalgam chronopotentiometry." *Russ Chem Rev* 45 (1976): 1.
  36. Duyckaerts george, De Graeve and Massaux J. *Actual Chim Anal Org Pharm Bromatol* 22 (1973): 39-68.
  37. Hutton A Emily, Samo B Hočevar, Božidar Ogorevc and Malcolm R Smyth. "Bismuth film electrode for simultaneous adsorptive stripping analysis of trace cobalt and nickel using constant current chronopotentiometric and voltammetric protocol." *Electrochem Commun* 5 (2003): 765-769.
  38. Mohamed Ahmad. "Determination of trace elements in sugar cane refuse by instrumental neutron activation analysis." *J Radioanal Nucl Chem* 107 (1986): 121-128.
  39. Gómez-Aracena Jorge, José M Martin-Moreno, Rudolph A Riemersma and Peter Bode, et al. "Association between toenail scandium levels and risk of acute myocardial infarction in European men: the EURAMIC and Heavy Metals Study." *Toxicol Ind Health* 18 (2002): 353-360.
  40. Chatt Amares, Hai-Son Dang, Byron B. Fong and Channa K Jayawickreme, et al. "Determination of trace elements in food by neutron activation analysis." *J Radioanal Nucl Chem* 124 (1988): 65-77.
  41. Giroussi Stella, Anastasios Voulgaropoulos, Aristomenis K Ayiannidis and Marcin Janicki. "Determination of cobalt in vegetable animal foodstuffs by differential pulse adsorptive voltammetry using  $\alpha$ -benzil dioxime." *Sci Total Environ* 176 (1995): 135-139.
  42. Honfo G Fernande, Noel Akissoe, Anita R Linnemann and Mohamed Soumanou, et al. "Nutritional composition of shea products and chemical properties of shea butter: A review." *Crit Rev Food Sci Nutr* 54 (2014): 673-686.
  43. Holak Walter and John J Specchio. "Interference by tin in AOAC dry-ash voltammetric method for determination of lead and cadmium in canned foods." *J Assoc Off Anal Chem* 71 (1988): 857-859.
  44. Hannisdal Atle and Knut H Schröder. "Differential pulse polarographic determination of the artificial sweeteners acesulfame-K and saccharin in beverages." *Electroanalysis* 5 (1993): 183-185.
  45. Khoulif Zidane, Claude Jambon, Michelle Chatelut and Olivier Vittori, et al. "Determination of heavy metals in concentrated refined sugar and raw syrups with differential pulse polarography and anodic stripping voltammetry." *Electroanalysis* 5 (1993): 339-342.
  46. Locatelli Clinio and Giancarlo Torsi. "Heavy metal determination in aquatic species for food purposes." *Anna Chim* 91 (2001): 65-72.
  47. Volke Jiri "Polarographic and voltammetric methods in pharmaceutical chemistry and pharmacology." *J Electroanal Chem Interf Electrochem* 155 (1983): 7-23.

48. Belal F Yousif. "The polarographic behavior of the antitumor agent taumustine and assay of its tablets." *Electroanalysis* 5 (1993): 605-609.
49. Chatt Amares, Hai-Son Dang, Byron B. Fong and Channa K Jayawickreme, et al. "Determination of trace elements in food by neutron activation analysis." *J Radioanal Nucl Chem* 124 (1988): 52-58.
50. Joerstad Kari and Brit Salbu. "Determination of trace elements in seawater by neutron activation analysis and electrochemical separation." *Analy Chem* 52 (1980): 672-676.
51. Pasquale Avino, Emiliano Santoro and Fabrizia Sarto. "Neutron activation analysis for investigating purity grade of copper, nickel and palladium thin films used in cold fusion experiments." *J Radioanal Nucl Chem* 290 (2011): 427-436.
52. Drabaek Iver, Van Carlsen and Lothar Just. "Routine determination of mercury, arsenic and selenium by radiochemical neutron activation analysis." *J Radioanal Nucl Chem* 103 (1986): 249-260.
53. Alghamdi H Ahmad. "Applications of stripping voltammetric techniques in food analysis." *Arab J Chem* 3 (2010): 1-7.
54. Miroslav Rievaj, Štefan Mesároš, Alica Brunova and Dušan Bustin, et al. "Application of Voltammetric Microelectrodes for Mercury Trace Analysis in Food Products-Fruit Juices and Beer." *Chem Pap* 51 (1997): 11-14.
55. Zhang Yongchao. Amperometric DNA sensing using wired enzyme based electrodes. The University of Texas at Austin, USA, 2003.
56. Wang Joseph, Den Bai Luo, Percio Ahmad M Farias and Jawad S Mahmoud, et al. "Adsorptive stripping voltammetry of riboflavin and other flavin analogs at the static mercury drop electrode." *Anal Chem* 57 (1985): 158-162.
57. Villar C Juan, Agustín Graciela and Bruno Nicolau Paulino. "Adsorptive stripping voltammetric behaviour of mitoxantrone on mercury electrodes." *Talanta* 40 (1993): 333-339.
58. Mercan Hülya and Recai İnam. "Differential pulse polarographic behaviour of thiazopyr herbicide and application to its determination in fruit juice and soil samples." *Int J Environ Anal Chem* 88 (2008): 879-890.
59. Gissel-Nielsen, Umesh C Gupta, Michel Lamand and Tuomas Westermarck, et al. "Selenium in soils and plants and its importance in livestock and human nutrition." *Adv Agron* 37 (1984): 397-460.
60. Ji-Wen Teng. "Geophysical Investigations Of The Earth's Crust And Upper Mantle In China [J]." *Chinese J Sinica* 4 (1979).
61. Zhao Zaofan, Jianhong Pei, Xiujing Zhang and Xingyao Zhou, et al. "Adsorptive stripping voltammetry determination of molybdenum (VI) in water and soil." *Talanta* 37 (1990): 1007-1010.
62. Kamenev Alexey, Viter, Gorshkova and Gus' Kov, et al. "Inversion voltammetric analysis of heavy metal ions in water." *Gig Sanit* 11 (1990): 93-94.

**How to cite this article:** Om Prakash Meena. "Heavy Metals Toxicity and the Environment Protection". *J Environ Anal Chem* 8 (2021): 326