

Percutaneous Anthropometric Dimensions of the Upper Arm and Forearm Bones: Determination of Sex and Stature in Adult Nigerian Population in Lagos

Robert Smithwick*

Department of Anatomical Sciences, School of Medicine, Jiroft University of Medical Sciences, Jiroft, Iran

Abstract

Background: Stature, like other phenotypic characteristics, is a crucial identification indicator that is influenced by both genetic and environmental factors. One of the most important and useful anthropometric parameters that determine a person's physical identity is stature, or body height. This is because of the alarming rise in the frequency of road, flood, intentional mutilation, and natural disasters. In forensic medical examination or anthropological studies, it is also regarded as one of the important and significant parameters for establishing personal identity.

Objectives: To measure the percutaneous length of the arm and forearm bones in an adult Nigerian population to determine stature and gender.

Methods: The sample group for this study consisted of 222 people between the ages of 18 and 65, split evenly between 115 men and 107 women. The workers, students, and volunteers at the University of Lagos made up the sample group. A wide range of anthropological instruments, including the Stadiometer, an anthropometric instrument made in Germany by SECA alpha®. A centimetre-calibrated calliper, weighing balance, and tape were used for the measurement.

Conclusion and Findings: When logistic regression was used to make a sex prediction, the intercondylar breadth had the highest value for statistical significance. The ulnar measurement had the strongest correlation with stature, while the intercondylar measurement had the weakest correlation across all parameters. Percutaneous measurement of arm and forearm lengths is available for both men and women, and it is reliable for estimating stature and determining sex. There is a stronger correlation between sexual dimorphism and the intercondylar. Straightforward and various direct relapses exhibited that utilizing the foot length is the most dependable procedure to anticipate and appraise height.

Keywords: Arm and forearm length • Stature • Sex • Correlation coefficient • Simple • Multiple linear regressions

Introduction

Stature In forensic medicine, the distance between a person's vertex and the standing surface is thought to be a crucial characteristic for personal identification. This parameter is influenced by a variety of variables, including genetics, nutrition, environment, gender, age, and physical activity. From one race, ethnic origin, and geographic area to another, the stature and bone length, as well as the factors affecting stature and bone length, vary greatly. Furthermore, stature has been implicated as a measure of development and growth. It has been used in nutrition and health research in clinical settings. According to studies, stature plays a crucial role in estimating nutrient needs as well as basal energy expenditure, body mass index, basal metabolic rate, body composition, and vital capacity. Even among the same subjects, anthropometric measurements have been reported to vary with populations global. Environmental and genetic influences are thought to be responsible for these variances. It is normal practise to utilise anthropometric methods to estimate skeletal length and stature from a subject's skeleton for over a

century, anthropologists, physicians, and anatomists have studied various body sections one hundred years [1-3].

Anatomical and mathematical techniques are the two main ways to estimate a person's height from long bones and other body parts. The anatomical method is thought to give the closest estimate of stature, but its fundamental drawback is that it needs the entire skeleton. While the mathematical approach just needs one bone or body part, it is less accurate. Regression analysis is thought to be a superior and more trustworthy mathematical method, nonetheless. In order to rebuild the deceased's biological profile, including age, sex, race, and stature, anthropometric measurements are more important than multiplication analysis. Estimating stature is one of these "big four" forensic anthropology's primary criteria for identifying a person during a forensic investigation. The biological basis for determining sex based on the physical and behavioural distinctions between males and females is known as sexual dimorphism. Sexual differences that affect bone size, shape, and appearance typically develop during growth as a result of unique genetic markers and a person's response to sex hormones during puberty. In both sexes, bone formation is influenced by a mix of hereditary factors and hormone exposure. There are a number of population-specific genetic and environmental factors that influence the age at which these sex-specific morphological alterations first manifest. Sex estimation standards must be population-specific since the degree of sexual dimorphism and the age at which it manifests in males and females differ between various communities. Since anthropologist has over the years concentrated on how to determine stature from skeletal remains, the current study was unable to find any studies that analyse the determination of stature and gender from percutaneous bones of arm and forearm. However, this study will be conducted on live subjects to estimate stature from the length of the arm and forearm percutaneous bones, and the findings will be used to teach anthropologists in this area how to estimate stature from partially skeletonized

*Address for Correspondence: Robert Smithwick, Department of Anatomical Sciences, School of Medicine, Jiroft University of Medical Sciences, Jiroft, Iran, E-mail: smithwick_r@gmail.com

Copyright: © 2022 Smithwick R. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Received: 01 November, 2022, Manuscript No. jfr-22-84252; **Editor assigned:** 03 November, 2022, PreQC No. P-84252; **Reviewed:** 15 November, 2022, QC No. Q-84252; **Revised:** 22 November, 2022, Manuscript No. R-84252; **Published:** 29 November, 2022, DOI: 10.37421/2157-7145.2022.13.523

bodies where the tissues are still required for additional forensic investigation [4-8].

Discussion

Study design Subjects were informed of the objectives of the study, Stature, hand anthropometric dimensions, clothing requirements, measurement procedures and freedom to withdraw were let known to the participants who met inclusion and exclusion criteria. A sample size of 230 convenient sample participants (100 males and 130 females) who were all Nigerian medical students (undergraduates and postgraduates), ages between 18 through 36 years were invited to the department of Anatomy, College of medicine, University of Lagos for measurement [9].

Percutaneous anthropometry

The measurement was taken as the maximum vertical distance from the floor to the vertex of the head using Stadiometer for the measurement. Protocol: Each participant was asked to stand barefooted on the flat platform, maintaining an erect position with their heels, buttocks, upper back with the head in contact with the wall. The arms were placed on the sides of the thigh and instructed to inhale and hold their breath. Their heads were held in the Frankfort horizontal plane. This position is achieved when the line joining the orbitale to the tragion is horizontal or at right angles to the long axis of the body. The horizontal sliding bar is then positioned on the contact point of the vertex of the head and stature is recorded in centimeters [10].

Conclusion

The present study will provide a database on an adult Nigerian population for the prediction of stature from hand dimensions. This could lead to the development of a standard for such data on young adult Nigerians. The comparison made with the other studies on different world population could contribute to the understanding of the relative status of Nigerian population in the context of the anthropometric variations around the world. Hence, the regression formulas derived from hand variables, may be of help when reconstructing stature from dismembered body parts, but it is only peculiar to a Nigerian population.

Acknowledgement

None.

Conflict of Interest

None.

References

1. Matsunobu, Yusuke, Junji Morishita, Yosuke Usumoto and Miki Okumura, et al. "Bone comparison identification method based on chest computed tomography imaging." *Leg Med* 29 (2017): 1-5.
2. Kawazoe, Yusuke, Junji Morishita, Yusuke Matsunobu and Miki Okumura, et al. "A simple method for semi-automatic readjustment for positioning in post-mortem head computed tomography imaging." *J Forensic Radiol Imaging* 16 (2019): 57-64.
3. Morishita, Junji, Shigehiko Katsuragawa, Keisuke Kondo and Kunio Doi, et al. "An automated patient recognition method based on an image-matching technique using previous chest radiographs in the picture archiving and communication system environment." *Med Phys* 28 (2001): 1093-1097.
4. Ueda, Yasuyuki, Junji Morishita, Shohei Kudomi and Katsuhiko Ueda, et al. "Usefulness of biological fingerprint in magnetic resonance imaging for patient verification." *Med Biol Eng Comput* 54 (2016): 1341-1351.
5. Sonoda, Minoru, Masao Takano, Junji Miyahara and Hisatoyo Kato. "Computed radiography utilizing scanning laser stimulated luminescence." *Radiol* 148 (1983): 833-838.
6. Nam-Goong, Il Seong, Ha Young Kim, Gyungyub Gong and Ho Kyu Lee. "ultrasonography-guided fine-needle aspiration of thyroid incidentaloma: Correlation with pathological findings." *Clin Endocrinol* 60 (2004): 21-28.
7. Stavros A Thomas, David Thickman, Cynthia L Rapp and Mark A. Dennis. "Solid breast nodules: Use of sonography to distinguish between benign and malignant lesions." *Radiology* 196 (1995): 134.
8. Agnihotri, Arun Kumar, Smriti Agnihotri, Nilima Jeebun and Krishna Googoolye. "Prediction of stature using hand dimensions." *J Forensic Leg Med* 15 (2008): 479-482.
9. Akhlaghi, Mitra, Marzieh Hajibeygi, Nasim Zamani and Behzad Moradi. "Estimation of stature from upper limb anthropometry in Iranian population." *J Forensic Leg Med* 19 (2012): 280-284.
10. Papini Enrico, Rinaldo Guglielmi, Antonio Bianchini and Anna Crescenzi. "Risk of malignancy in nonpalpable thyroid nodules: Predictive value of ultrasound and color-Doppler Features." *J Clin Endocrinol Metabol* 87 (2002): 1941-1946.

How to cite this article: Smithwick, Robert. "Percutaneous Anthropometric Dimensions of the Upper Arm and Forearm Bones: Determination of Sex and Stature in Adult Nigerian Population in Lagos." *J Forensic Res* 13 (2022): 523.